

THE ALGEBRA OF REGULAR EXPRESSIONS

Reminder of Basic Definitions and Some Basic Proofs

(1) For languages $L, M \subseteq \Sigma^*$; $L+M$, $L.M$ and L^* are interpreted as follows :

$L+M = L \cup M$; $L.M = \{w \mid w = u.v; u \in L; v \in M\}$; $L^* = \bigcup_{i=0,+\infty} L^i$ where $L^i := L.L \dots L$ (i times)

(2) $(L+M)^* = (L^*.M^*)^*$

Proof of (2) :

Let $u \in (L+M)^*$ then by definition $u = u_1.u_2 \dots u_k$ for some integer $k \geq 0$ where for each j

$u_j \in L+M$. But $L \subseteq L^* \subseteq L^*$, $e \subseteq L^*$, $M^* \subseteq e.M^* \subseteq L^*.M^*$ and thus

$u_j \in L^*.M^* + L^*.M^* = L^*.M^*$ and therefore $(L+M)^* \subseteq (L^*.M^*)^*$

Conversely let $u \in (L^*.M^*)^*$ then by definition $u = u_1.u_2 \dots u_k$ where $u_j \in L^*.M^*$ hence

$u_j = v_j^1.v_j^2 \dots v_j^{l(j)}.w_j^1.w_j^2 \dots w_j^{p(j)}$ where $v_j^m \in L$ and $w_j^m \in M$. Hence

$u = z_1.z_2 \dots z_q$ where $q = \sum_{j=1,k} l(j)+p(j)$ and each $z_j \in L+M$. This proves that $(L^*.M^*)^* \subseteq (L+M)^*$

which proves that $(L+M)^* = (L^*.M^*)^*$

(3) $(L+M)^* = (L^*+M^*)^*$

Proof of (3) :

Since $L \subseteq L^*$ and $M \subseteq M^*$ it follows that $(L+M)^* \subseteq (L^*+M^*)^*$.

Conversely let $u \in (L^*+M^*)^*$ then $u = (v_1+w_1) \dots (v_k+w_k)$ where for each j

$v_j \in L^*$ and $w_j \in M^*$. We show that $u \in (L^*.M^*)^*$ by using induction on k . For $k=1$

$v_1 \in L^* \subseteq L^*$, $e \subseteq L^*.M^* \subseteq (L^*.M^*)^*$ similarly $w_1 \in M^* \subseteq e.M^* \subseteq L^*.M^* \subseteq (L^*.M^*)^*$ hence

$v_1+w_1 \in (L^*.M^*)^*$. Now assume statement holds for $k-1$, hence

$z := (v_1+w_1) \dots (v_{k-1}+w_{k-1}) \in (L^*.M^*)^*$. But using the above reasoning for v_k+w_k it follows that

$v_k+w_k \in (L^*.M^*)^*$ and therefore $u = z.(v_k+w_k) \in (L^*.M^*)^*$. $(L^*.M^*)^* = (L^*.M^*)^*$ using the

obvious identity $K^*.K^* = K^*$ for any language K . This proves that $(L^*+M^*)^* \subseteq (L^*.M^*)^*$, but by (2)

$(L+M)^* = (L^*.M^*)^*$ hence $(L^*+M^*)^* \subseteq (L+M)^*$ and (3) is proved.

(4) $(L.M)^* \subseteq (L^*.M^*)^*$ and $(L.M)^* = (L^*.M^*)^*$ iff $e \in L$ and $e \in M$

Proof of (4) :

First statement is obvious using $L \subseteq L^*$ and $M \subseteq M^*$.

To prove the second one assume $e \in L$ and $e \in M$

and let $u \in (L^*.M^*)^*$ then $u = v_1.w_1 \dots v_k.w_k$ where $v_j \in L^*$ and $w_j \in M^*$ therefore

$v_j = v_j^1 \dots v_j^{l(j)}$ and $w_j = z_j^1 \dots z_j^{p(j)}$ with $v_j^m \in L$ and $z_j^m \in M$. Hence

$u = q_1 \dots q_r$ where $r = \sum_{j=1,k} (l(j)+p(j))$ where each $q_i \in L$ or $q_i \in M$. Using the assumption we can

write $u = q'_1 . \dots . q'_r$, by adding an empty string in between the q_j strings, if necessary, so that we have for each $j=1, \dots, r$, $q'_j \in L$ and $q'_{j+1} \in M$. This proves that $u \in (L.M)^*$. To prove the converse result we present counter-examples that violate the assumption $e \in L$ and $e \in M$. Suppose $e \notin L$ choose $L = 0.0^*$ and $M = 1^*$ then $1 \in (L.M)^*$ whereas $1 \notin (L.M)^*$; alternatively if $e \notin M$ choose $L = 0^*$ and $M = 1.1^*$ then $0 \in (L.M)^*$ whereas $0 \notin (L.M)^*$.

Homework #2 due October 18, 2016, Tuesday

(1) Using either the results or the techniques used above try to simplify the following expressions and prove your simplification.

- (i) $(0+1)^*.1.(0+1) + (0+1)^*.1.(0+1)$
- (ii) $((0^*.1^*)+1)^*(0+1)^*$
- (iii) $(L+M^*)^*$
- (iv) $(L.M^*)^*$

(2) Convert the regular expression $((0.0)^*. (1.1))^* . 01$ into an ϵ -NFA

(3) Problems from the textbook

- 3.1.1 (b) and (c)
- 3.1.4 (b) and (c)
- 3.2.1 (c) and (d)
- 3.2.3