

If $A = UDV^T$, then $A^T A$ is:

(A) UD^2U^T

(B) VD^2V^T

(C) A square matrix

(D) is always full rank

(E) none of the above

Consider a set of general vectors $\mathbf{a}_i \in \mathbb{R}^d$. (assume all elements are some random numbers in the range of $[0, 1]$) \mathbf{b} is another such vector. Consider the matrix:

$$\mathbf{A} = \sum_{i=1}^k \mathbf{a}_i \mathbf{a}_i^T + \sum_{i=k+1}^d \mathbf{b} \mathbf{b}^T$$

What is the effective rank of \mathbf{A}

- (A) k
- (B) 1
- (C) none of the above
- (D) $k + 1$
- (E) d

Consider a set of general vectors $\mathbf{a}_i \in \mathbb{R}^d$. (assume all elements are some random numbers in the range of $[0, 1]$) \mathbf{b} is another such vector. Consider the matrix:

$$\mathbf{A} = \sum_{i=1}^k 10^{-k} \mathbf{a}_i \mathbf{a}_i^T + \sum_{i=k+1}^d \mathbf{b} \mathbf{b}^T$$

What is the effective rank of \mathbf{A}

- (A) $k + 1$
- (B) k
- (C) d
- (D) 1
- (E) none of the above

Consider a set of general vectors $\mathbf{a}_i \in \mathbb{R}^d$. (assume all elements are some random numbers in the range of $[0, 1]$) \mathbf{b} is another such vector. Consider the matrix:

$$\mathbf{A} = \sum_{i=1}^k 10^i \mathbf{a}_i \mathbf{a}_i^T + \sum_{i=k+1}^d 10^{-i} \mathbf{b} \mathbf{b}^T$$

What is the effective rank of \mathbf{A}

- (A) 1
- (B) k
- (C) $k + 1$
- (D) d
- (E) none of the above

Consider a set of general vectors $\mathbf{a}_i \in \mathbb{R}^d$. (assume all elements are some random numbers in the range of $[0, 1]$) \mathbf{b} is another such vector. Consider the matrix:

$$\mathbf{A} = \sum_{i=1}^k 10^{-i} \mathbf{a}_i \mathbf{a}_i^T + \sum_{i=k+1}^d 10^i \mathbf{b} \mathbf{b}^T$$

What is the effective rank of \mathbf{A}

- (A) 1
- (B) none of the above
- (C) k
- (D) d
- (E) $k + 1$