

Topics in Applied Optimization

Optimization Algorithms for ML and Data Sciences

Pawan Kumar

CSTAR, IIIT-H

Convex Optimization Problems

Convex Optimization Problems

Optimization problem:

$$\begin{array}{ll}\text{minimize} & f_0(x) \\ \text{subject to} & f_i(x) \leq 0, \quad i = 1, \dots, m \\ & h_i(x) = 0, \quad i = 1, \dots, p\end{array}$$

Convex Optimization Problems

Optimization problem:

$$\begin{array}{ll}\text{minimize} & f_0(x) \\ \text{subject to} & f_i(x) \leq 0, \quad i = 1, \dots, m \\ & h_i(x) = 0, \quad i = 1, \dots, p\end{array}$$

- Find an x that **minimizes** $f_0(x)$ among all x that satisfy
 - $f_i(x) \leq 0, i = 1, \dots, m$
 - $h_i(x) = 0, i = 1, \dots, p$

Convex Optimization Problems

Optimization problem:

$$\begin{array}{ll}\text{minimize} & f_0(x) \\ \text{subject to} & f_i(x) \leq 0, \quad i = 1, \dots, m \\ & h_i(x) = 0, \quad i = 1, \dots, p\end{array}$$

- Find an x that **minimizes** $f_0(x)$ among all x that satisfy
 - $f_i(x) \leq 0, i = 1, \dots, m$
 - $h_i(x) = 0, i = 1, \dots, p$
- x is called **optimization variable**

Convex Optimization Problems

Optimization problem:

$$\begin{array}{ll}\text{minimize} & f_0(x) \\ \text{subject to} & f_i(x) \leq 0, \quad i = 1, \dots, m \\ & h_i(x) = 0, \quad i = 1, \dots, p\end{array}$$

- Find an x that **minimizes** $f_0(x)$ among all x that satisfy
 - $f_i(x) \leq 0, i = 1, \dots, m$
 - $h_i(x) = 0, i = 1, \dots, p$
- x is called **optimization variable**
- $f_0 : \mathbb{R}^n \rightarrow \mathbb{R}$ is called **objective function or cost function**

Convex Optimization Problems

Optimization problem:

$$\begin{aligned} & \text{minimize} && f_0(x) \\ & \text{subject to} && f_i(x) \leq 0, \quad i = 1, \dots, m \\ & && h_i(x) = 0, \quad i = 1, \dots, p \end{aligned}$$

- Find an x that **minimizes** $f_0(x)$ among all x that satisfy
 - $f_i(x) \leq 0, i = 1, \dots, m$
 - $h_i(x) = 0, i = 1, \dots, p$
- x is called **optimization variable**
- $f_0 : \mathbb{R}^n \rightarrow \mathbb{R}$ is called **objective function or cost function**
- The inequalities $f_i(x) \leq 0$ are called **inequality** constraints

Convex Optimization Problems

Optimization problem:

$$\begin{aligned} & \text{minimize} && f_0(x) \\ & \text{subject to} && f_i(x) \leq 0, \quad i = 1, \dots, m \\ & && h_i(x) = 0, \quad i = 1, \dots, p \end{aligned}$$

- Find an x that **minimizes** $f_0(x)$ among all x that satisfy
 - $f_i(x) \leq 0, i = 1, \dots, m$
 - $h_i(x) = 0, i = 1, \dots, p$
- x is called **optimization variable**
- $f_0 : \mathbb{R}^n \rightarrow \mathbb{R}$ is called **objective function or cost function**
- The inequalities $f_i(x) \leq 0$ are called **inequality constraints**
- $f_i : \mathbb{R}^n \rightarrow \mathbb{R}$ are called **inequality constraint functions**

Convex Optimization Problems

Optimization problem:

$$\begin{aligned} & \text{minimize} && f_0(x) \\ & \text{subject to} && f_i(x) \leq 0, \quad i = 1, \dots, m \\ & && h_i(x) = 0, \quad i = 1, \dots, p \end{aligned}$$

- Find an x that **minimizes** $f_0(x)$ among all x that satisfy
 - $f_i(x) \leq 0, i = 1, \dots, m$
 - $h_i(x) = 0, i = 1, \dots, p$
- x is called **optimization variable**
- $f_0 : \mathbb{R}^n \rightarrow \mathbb{R}$ is called **objective function or cost function**
- The inequalities $f_i(x) \leq 0$ are called **inequality constraints**
- $f_i : \mathbb{R}^n \rightarrow \mathbb{R}$ are called **inequality constraint functions**
- The equations $h_i(x)$ are called **equality constraints**

Convex Optimization Problems

Optimization problem:

$$\begin{aligned} & \text{minimize} && f_0(x) \\ & \text{subject to} && f_i(x) \leq 0, \quad i = 1, \dots, m \\ & && h_i(x) = 0, \quad i = 1, \dots, p \end{aligned}$$

- Find an x that **minimizes** $f_0(x)$ among all x that satisfy
 - $f_i(x) \leq 0, i = 1, \dots, m$
 - $h_i(x) = 0, i = 1, \dots, p$
- x is called **optimization variable**
- $f_0 : \mathbb{R}^n \rightarrow \mathbb{R}$ is called **objective function or cost function**
- The inequalities $f_i(x) \leq 0$ are called **inequality constraints**
- $f_i : \mathbb{R}^n \rightarrow \mathbb{R}$ are called **inequality constraint functions**
- The equations $h_i(x)$ are called **equality constraints**
- The functions $h_i : \mathbb{R}^n \rightarrow \mathbb{R}$ are called **equality constraint functions**

Convex Optimization Problems

Optimization problem:

$$\text{minimize } f_0(x) \tag{1}$$

$$\text{subject to } f_i(x) \leq 0, i = 1, \dots, m \tag{2}$$

$$h_i(x) = 0, i = 1, \dots, p \tag{3}$$

Convex Optimization Problems

Optimization problem:

$$\text{minimize } f_0(x) \quad (1)$$

$$\text{subject to } f_i(x) \leq 0, i = 1, \dots, m \quad (2)$$

$$h_i(x) = 0, i = 1, \dots, p \quad (3)$$

- If **no constraints**, i.e., $m = p = 0$, then called **unconstrained problem**

Convex Optimization Problems

Optimization problem:

$$\text{minimize } f_0(x) \quad (1)$$

$$\text{subject to } f_i(x) \leq 0, i = 1, \dots, m \quad (2)$$

$$h_i(x) = 0, i = 1, \dots, p \quad (3)$$

- If **no constraints**, i.e., $m = p = 0$, then called **unconstrained problem**
- **Domain of opt. problem**: where objective and constraint are defined

$$\mathcal{D} = \bigcap_{i=0}^m \text{dom } f_i \cap \bigcap_{i=1}^p \text{dom } h_i$$

Handwritten orange note: f_0 with an arrow pointing to the $i=0$ term in the first intersection.

Convex Optimization Problems

Optimization problem:

$$\text{minimize } f_0(x) \quad (1)$$

$$\text{subject to } f_i(x) \leq 0, i = 1, \dots, m \quad (2)$$

$$h_i(x) = 0, i = 1, \dots, p \quad (3)$$

- If **no constraints**, i.e., $m = p = 0$, then called **unconstrained problem**
- **Domain of opt. problem**: where objective and constraint are defined

$$\mathcal{D} = \bigcap_{i=0}^m \text{dom } f_i \quad \cap \quad \bigcap_{i=1}^p \text{dom } h_i$$

- A point x is called **feasible** if it satisfies the constraints

$$f_i(x) \leq 0, \quad i = 1, \dots, m$$

$$h_i(x) = 0, \quad i = 1, \dots, p$$

Convex Optimization Problems

Optimization problem:

$$\text{minimize } f_0(x) \quad (1)$$

$$\text{subject to } f_i(x) \leq 0, i = 1, \dots, m \quad (2)$$

$$h_i(x) = 0, i = 1, \dots, p \quad (3)$$

- If **no constraints**, i.e., $m = p = 0$, then called **unconstrained problem**
- **Domain of opt. problem**: where objective and constraint are defined

$$\mathcal{D} = \bigcap_{i=0}^m \text{dom } f_i \quad \cap \quad \bigcap_{i=1}^p \text{dom } h_i$$

- A point x is called **feasible** if it satisfies the constraints

$$f_i(x) \leq 0, \quad i = 1, \dots, m$$

$$h_i(x) = 0, \quad i = 1, \dots, p$$

- The optimization problem is called **feasible** if there exists atleast one feasible point

Convex Optimization Problems

Optimization problem:

$$\begin{array}{ll}\text{minimize} & f_0(x) \\ \text{subject to} & f_i(x) \leq 0, i = 1, \dots, m \\ & h_i(x) = 0, i = 1, \dots, p\end{array}$$

Convex Optimization Problems

Optimization problem:

$$\begin{array}{ll}\text{minimize} & f_0(x) \\ \text{subject to} & f_i(x) \leq 0, i = 1, \dots, m \\ & h_i(x) = 0, i = 1, \dots, p\end{array}$$

- Feasible set or constraint set: set of all feasible points

Convex Optimization Problems

Optimization problem:

$$\begin{aligned} & \text{minimize} && f_0(x) \\ & \text{subject to} && f_i(x) \leq 0, i = 1, \dots, m \\ & && h_i(x) = 0, i = 1, \dots, p \end{aligned}$$

- Feasible set or constraint set: set of all feasible points
- Optimal Value: The optimal value p^* defined as

$$p^* = \inf \{ f_0(x) \mid f_i(x) \leq 0, i = 1, \dots, m, h_i(x) = 0, i = 1, \dots, p \}$$

Convex Optimization Problems

Optimization problem:

$$\begin{array}{ll}\text{minimize} & f_0(x) \\ \text{subject to} & f_i(x) \leq 0, i = 1, \dots, m \\ & h_i(x) = 0, i = 1, \dots, p\end{array}$$

- Feasible set or constraint set: set of all feasible points
- Optimal Value: The optimal value p^* defined as

$$p^* = \inf \{f_0(x) \mid f_i(x) \leq 0, i = 1, \dots, m, h_i(x) = 0, i = 1, \dots, p\}$$

- p^* is allowed to take extended values $\pm\infty$

Convex Optimization Problems

Optimization problem:

$$\begin{aligned} & \text{minimize} && f_0(x) \\ & \text{subject to} && f_i(x) \leq 0, i = 1, \dots, m \\ & && h_i(x) = 0, i = 1, \dots, p \end{aligned}$$

- **Feasible set or constraint set:** set of all feasible points
- **Optimal Value:** The **optimal** value p^* defined as

$$p^* = \inf \{ f_0(x) \mid f_i(x) \leq 0, i = 1, \dots, m, h_i(x) = 0, i = 1, \dots, p \}$$

- p^* is allowed to take **extended** values $\pm\infty$
- **Infeasible problem:** problem is called **infeasible** when $p^* = \infty$

inf $\{\emptyset\}$
↓
 ∞

Convex Optimization Problems

Optimization problem:

$$\begin{aligned} & \text{minimize} && f_0(x) \\ & \text{subject to} && f_i(x) \leq 0, i = 1, \dots, m \\ & && h_i(x) = 0, i = 1, \dots, p \end{aligned}$$

- **Feasible set or constraint set:** set of all feasible points
- **Optimal Value:** The **optimal** value p^* defined as

$$p^* = \text{inf} \{ f_0(x) \mid f_i(x) \leq 0, i = 1, \dots, m, h_i(x) = 0, i = 1, \dots, p \}$$

- p^* is allowed to take **extended** values $\pm\infty$
- **Infeasible problem:** problem is called **infeasible** when $p^* = \infty$
 - Note: we used the fact that $\inf \phi = \infty$

Convex Optimization Problems

Optimization problem:

$$\begin{aligned} & \text{minimize} && f_0(x) \\ & \text{subject to} && f_i(x) \leq 0, i = 1, \dots, m \\ & && h_i(x) = 0, i = 1, \dots, p \end{aligned}$$

- **Feasible set or constraint set:** set of all feasible points
- **Optimal Value:** The **optimal** value p^* defined as

$$p^* = \text{inf} \{ f_0(x) \mid f_i(x) \leq 0, i = 1, \dots, m, h_i(x) = 0, i = 1, \dots, p \}$$

- p^* is allowed to take **extended** values $\pm\infty$
- **Infeasible problem:** problem is called **infeasible** when $p^* = \infty$
 - Note: we used the fact that $\inf \phi = \infty$
- **Unbounded below:** Problem is **unbounded below** if $f_0(x_k) \rightarrow -\infty$ as $k \rightarrow \infty$

Optimal and locally optimal

Optimization problem:

$$\begin{array}{ll}\text{minimize} & f_0(x) \\ \text{subject to} & f_i(x) \leq 0, i = 1, \dots, m \\ & h_i(x) = 0, i = 1, \dots, p\end{array}$$

Optimal and locally optimal

Optimization problem:

$$\begin{array}{ll}\text{minimize} & f_0(x) \\ \text{subject to} & f_i(x) \leq 0, i = 1, \dots, m \\ & h_i(x) = 0, i = 1, \dots, p\end{array}$$

- **Optimal Point:** x^* is called **optimal** if it solves the given optimization problem, i.e.,

Optimal and locally optimal

Optimization problem:

$$\begin{array}{ll}\text{minimize} & f_0(x) \\ \text{subject to} & f_i(x) \leq 0, i = 1, \dots, m \\ & h_i(x) = 0, i = 1, \dots, p\end{array}$$

- **Optimal Point:** x^* is called **optimal** if it solves the given optimization problem, i.e.,
 - x^* is **feasible** point

Optimal and locally optimal

Optimization problem:

$$\begin{aligned} & \text{minimize} && f_0(x) \\ & \text{subject to} && f_i(x) \leq 0, i = 1, \dots, m \\ & && h_i(x) = 0, i = 1, \dots, p \end{aligned}$$

- **Optimal Point:** x^* is called **optimal** if it solves the given optimization problem, i.e.,
 - x^* is **feasible** point
 - $f(x^*) = p^*$, that is, at x^* **optimal value** p^* is obtained

Optimal and locally optimal

Optimization problem:

$$\begin{array}{ll}\text{minimize} & f_0(x) \\ \text{subject to} & f_i(x) \leq 0, i = 1, \dots, m \\ & h_i(x) = 0, i = 1, \dots, p\end{array}$$

- **Optimal Point:** x^* is called **optimal** if it solves the given optimization problem, i.e.,
 - x^* is **feasible** point
 - $f(x^*) = p^*$, that is, at x^* **optimal value** p^* is obtained
- **Optimal Set:** The set of **all** optimal points is called **optimal set**

Optimal and locally optimal

Optimization problem:

$$\begin{array}{ll}\text{minimize} & f_0(x) \\ \text{subject to} & f_i(x) \leq 0, i = 1, \dots, m \\ & h_i(x) = 0, i = 1, \dots, p\end{array}$$

- **Optimal Point:** x^* is called **optimal** if it solves the given optimization problem, i.e.,
 - x^* is **feasible** point
 - $f(x^*) = p^*$, that is, at x^* **optimal value** p^* is obtained
- **Optimal Set:** The set of **all** optimal points is called **optimal set**
- If there exists an optimal point, then we say that optimal value is **achieved** and the problem is **solvable**

Optimal and locally optimal

Optimization problem:

$$\begin{aligned} & \text{minimize} && f_0(x) \\ & \text{subject to} && f_i(x) \leq 0, i = 1, \dots, m \\ & && h_i(x) = 0, i = 1, \dots, p \end{aligned}$$

- **Optimal Point:** x^* is called **optimal** if it solves the given optimization problem, i.e.,
 - x^* is **feasible** point
 - $f(x^*) = p^*$, that is, at x^* **optimal value** p^* is obtained
- **Optimal Set:** The set of **all** optimal points is called **optimal set**
- If there exists an optimal point, then we say that optimal value is **achieved** and the problem is **solvable**
- If the optimal set is **empty**, then we say that optimal value is **not** attained

Optimal and Locally Optimal Points

Optimization problem:

$$\begin{array}{ll}\text{minimize} & f_0(x) \\ \text{subject to} & f_i(x) \leq 0, i = 1, \dots, m \\ & h_i(x) = 0, i = 1, \dots, p\end{array}$$

Optimal and Locally Optimal Points

Optimization problem:

$$\begin{aligned} & \text{minimize} && f_0(x) \\ & \text{subject to} && f_i(x) \leq 0, i = 1, \dots, m \\ & && h_i(x) = 0, i = 1, \dots, p \end{aligned}$$

- ϵ -suboptimal: \bar{x} is ϵ -suboptimal if $f(\bar{x}) \leq p^* + \epsilon$, $\epsilon > 0$

Optimal and Locally Optimal Points

Optimization problem:

$$\begin{aligned} & \text{minimize} && f_0(x) \\ & \text{subject to} && f_i(x) \leq 0, i = 1, \dots, m \\ & && h_i(x) = 0, i = 1, \dots, p \end{aligned}$$

- ϵ -suboptimal: \bar{x} is ϵ -suboptimal if $f(\bar{x}) \leq p^* + \epsilon$, $\epsilon > 0$
 - \bar{x} is just ϵ more than optimal value p^*

Optimal and Locally Optimal Points

Optimization problem:

$$\begin{array}{ll}\text{minimize} & f_0(x) \\ \text{subject to} & f_i(x) \leq 0, i = 1, \dots, m \\ & h_i(x) = 0, i = 1, \dots, p\end{array}$$

- **ϵ —suboptimal:** \bar{x} is ϵ —suboptimal if $f(\bar{x}) \leq p^* + \epsilon$, $\epsilon > 0$
 - \bar{x} is **just** ϵ more than optimal value p^*
- **ϵ —suboptimal set:** set of all ϵ —suboptimal points

Optimal and Locally Optimal Points

Optimization problem:


$$\begin{array}{ll}\text{minimize} & f_0(x) \\ \text{subject to} & f_i(x) \leq 0, i = 1, \dots, m \\ & h_i(x) = 0, i = 1, \dots, p\end{array}$$



- **ϵ -suboptimal:** \bar{x} is ϵ -suboptimal if $f(\bar{x}) \leq p^* + \epsilon$, $\epsilon > 0$
 - \bar{x} is **just** ϵ more than optimal value p^*
- **ϵ -suboptimal set:** set of all ϵ -suboptimal points
- **Active constraint:** If x is feasible and $f_i(x) = 0$, then i th inequality is **active**

Optimal and Locally Optimal Points

Optimization problem:


$$\begin{aligned} & \text{minimize} && f_0(x) \\ & \text{subject to} && f_i(x) \leq 0, i = 1, \dots, m \\ & && h_i(x) = 0, i = 1, \dots, p \end{aligned}$$

- **ϵ -suboptimal:** \bar{x} is ϵ -suboptimal if $f(\bar{x}) \leq p^* + \epsilon$, $\epsilon > 0$
 - \bar{x} is **just** ϵ more than optimal value p^*
- **ϵ -suboptimal set:** set of all ϵ -suboptimal points
- **Active constraint:** If x is feasible and $f_i(x) = 0$, then i th inequality is **active**
- **Inactive constraint:** If x is **feasible** and $f_i(x) < 0$, then this constraint is **inactive**

Optimal and Locally Optimal Points

Optimization problem:

$$\begin{array}{ll}\text{minimize} & f_0(x) \\ \text{subject to} & f_i(x) \leq 0, i = 1, \dots, m \\ & h_i(x) = 0, i = 1, \dots, p\end{array}$$

- **ϵ —suboptimal:** \bar{x} is ϵ —suboptimal if $f(\bar{x}) \leq p^* + \epsilon$, $\epsilon > 0$
 - \bar{x} is **just** ϵ more than optimal value p^*
- **ϵ —suboptimal set:** set of all ϵ —suboptimal points
- **Active constraint:** If x is feasible and $f_i(x) = 0$, then i th inequality is **active**
- **Inactive constraint:** If x is **feasible** and $f_i(x) < 0$, then this constraint is **inactive**
- **Redundant constraint:** A constraint is **redundant** if removing it **does not change** the feasible set

Optimization problem:

$$\begin{array}{ll}\text{minimize} & f_0(x) \\ \text{subject to} & f_i(x) \leq 0, i = 1, \dots, m \\ & h_i(x) = 0, i = 1, \dots, p\end{array}$$

Optimization problem:

$$\begin{array}{ll}\text{minimize} & f_0(x) \\ \text{subject to} & f_i(x) \leq 0, i = 1, \dots, m \\ & h_i(x) = 0, i = 1, \dots, p\end{array}$$

Define the feasible set

$$\Omega = \{x \mid \underline{f_i(x) \leq 0}, \quad i = 1, \dots, m, \quad \underline{h_i(x) = 0} \quad i = 1, \dots, p\}$$


Optimization problem:

$$\begin{array}{ll}\text{minimize} & f_0(x) \\ \text{subject to} & f_i(x) \leq 0, i = 1, \dots, m \\ & h_i(x) = 0, i = 1, \dots, p\end{array}$$

Define the **feasible set**

$$\Omega = \{x \mid f_i(x) \leq 0, \quad i = 1, \dots, m, \quad h_i(x) = 0 \quad i = 1, \dots, p\}$$

More **compactly**, we we can write:



Examples: $1/x$

Consider the optimization problem:

$$\begin{aligned} &\text{minimize } f_0(x) = 1/x, \\ &\text{subject to } x \in \mathbb{R} \end{aligned}$$


where $f_0 : \mathbb{R}_{++} \rightarrow \mathbb{R}$

Quiz: What is feasible set?

Quiz: What is p^* ?

Quiz: Is the optimal value achieved?

Figure 1: Plot of $1/x$



Examples: $-\log x$

Consider the optimization problem:

$$\begin{aligned} &\text{minimize } f_0(x) = -\log x, \\ &\text{subject to } x \in \mathbb{R} \end{aligned}$$

where $f_0 : \mathbb{R}_{++} \rightarrow \mathbb{R}$


Quiz: What is feasible set?

Quiz: What is p^* ?

Quiz: Is the optimal value achieved?

Quiz: Is this problem bounded below?

Figure 2: Plot of $-\log x$



Examples: $x \log x$

Consider the optimization problem:

$$\begin{aligned} &\text{minimize } f_0(x) = x \log x, \\ &\text{subject to } x \in \mathbb{R} \end{aligned}$$

where $f_0 : \mathbb{R}_{++} \rightarrow \mathbb{R}$

Quiz: What is feasible set?


Quiz: What is p^* ?

Quiz: Is the optimal value achieved?


Quiz: Is this problem bounded below?

Quiz: What is optimal point?

Figure 3: Plot of $x \log x$



Examples: Graphically




Expressing Problems in Standard Form

Optimization problem (Standard Form):

$$\begin{array}{ll}\text{minimize} & f_0(x) \\ \text{subject to} & f_i(x) \leq 0, i = 1, \dots, m \\ & h_i(x) = 0, i = 1, \dots, p\end{array}$$

Expressing Problems in Standard Form

Optimization problem (Standard Form):

$$\begin{array}{ll}\text{minimize} & f_0(x) \\ \text{subject to} & f_i(x) \leq 0, i = 1, \dots, m \\ & h_i(x) = 0, i = 1, \dots, p\end{array}$$


Convention for standard form:

- **righthand** side of the inequality and equality constraints **are zero**

Expressing Problems in Standard Form

Optimization problem (Standard Form):

$$\begin{aligned} & \text{minimize} && f_0(x) \\ & \text{subject to} && f_i(x) \leq 0, i = 1, \dots, m \\ & && h_i(x) = 0, i = 1, \dots, p \end{aligned}$$

Convention for standard form:

- **righthand** side of the inequality and equality constraints **are zero**
 - For example, $\underbrace{g_i(x) = \tilde{g}_i(x)}$ can be written as $h_i(x) = 0$

hi: $g_i(x) - \tilde{g}_i(x) = 0$

Expressing Problems in Standard Form

Optimization problem (Standard Form):

$$\begin{array}{ll}\text{minimize} & f_0(x) \\ \text{subject to} & f_i(x) \leq 0, i = 1, \dots, m \\ & h_i(x) = 0, i = 1, \dots, p\end{array}$$

only \leq is allowed

Convention for standard form:


- **righthand** side of the inequality and equality constraints **are zero**
 - For example, $g_i(x) = \tilde{g}_i(x)$ can be written as $h_i(x) = 0$
- $f_i(x) \geq 0$ as $-f_i(x) \leq 0$

~~~~~

(Box Constraints). Consider the following

$$\begin{array}{ll}\text{minimize} & f_0(x) \\ \text{subject to} & \underline{l_i} \leq x_i \leq \underline{u_i}, \quad i = 1, \dots, n\end{array}$$

- constraints here are called **variable bounds** or **box constraints**





(Box Constraints). Consider the following

$$\begin{array}{ll}\text{minimize} & f_0(x) \\ \text{subject to} & l_i \leq x_i \leq u_i, \quad i = 1, \dots, n\end{array}$$

- constraints here are called **variable bounds** or **box constraints**

The problem above **can** be expressed in **standard form**:

$$\begin{array}{ll}\text{minimize} & f_0(x) \\ \text{subject to} & \underline{l_i} - x_i \leq 0, \quad i = 1, \dots, n \\ & x_i - \underline{u_i} \leq 0, \quad i = 1, \dots, n\end{array}$$

(Box Constraints). Consider the following

$$\begin{array}{ll}\text{minimize} & f_0(x) \\ \text{subject to} & l_i \leq x_i \leq u_i, \quad i = 1, \dots, n\end{array}$$

- constraints here are called **variable bounds** or **box constraints**

The problem above **can** be expressed in **standard form**:

$$\begin{array}{ll}\text{minimize} & f_0(x) \\ \text{subject to} & l_i - x_i \leq 0, \quad i = 1, \dots, n \\ & x_i - u_i \leq 0, \quad i = 1, \dots, n\end{array}$$

$$\begin{array}{ll}\text{minimize} & f_0(x) \\ \text{subject to} & f_i(x) \leq 0, \quad i = 1, \dots, 2n\end{array}$$

where  $f_i(x) = l_i - x_i$ ,  $i = 1, \dots, n$  and  $f_i(x) = x_{i-n} - u_{i-n}$ ,  $i = n+1, \dots, 2n$

## Maximization Problems Seen as Minimization Problems

**Note:** Maximization problem can be solved by minimization. Consider

$$\begin{array}{ll} \text{maximize} & f_0(x) \\ \text{subject to} & f_i(x) \leq 0, \quad i = 1, \dots, m \\ & h_i(x) = 0, \quad i = 1, \dots, p, \end{array}$$

It can be solved by solving a corresponding minimization problem with  $-f_0(x)$

## Maximization Problems Seen as Minimization Problems

**Note:** Maximization problem can be solved by minimization. Consider

$$\begin{array}{ll}\text{maximize} & f_0(x) \\ \text{subject to} & f_i(x) \leq 0, \quad i = 1, \dots, m \\ & h_i(x) = 0, \quad i = 1, \dots, p,\end{array}$$

It can be solved by solving a corresponding minimization problem with  $-f_0(x)$

Optimization problem (Standard Form):

$$\begin{array}{ll}\text{minimize} & -f_0(x) \\ \text{subject to} & f_i(x) \leq 0, \quad i = 1, \dots, m \\ & h_i(x) = 0, \quad i = 1, \dots, p\end{array}$$

## Maximization Problems Seen as Minimization Problems

**Note:** Maximization problem can be solved by minimization. Consider

$$\begin{aligned} &\text{maximize} && f_0(x) \\ &\text{subject to} && f_i(x) \leq 0, \quad i = 1, \dots, m \\ & && h_i(x) = 0, \quad i = 1, \dots, p, \end{aligned}$$

It can be solved by solving a corresponding minimization problem with  $-f_0(x)$

Optimization problem (Standard Form):

$$\begin{aligned} &\text{minimize} && -f_0(x) \\ &\text{subject to} && f_i(x) \leq 0, \quad i = 1, \dots, m \\ & && h_i(x) = 0, \quad i = 1, \dots, p \end{aligned}$$

- Note the constraints remain same!

## Maximization Problems Seen as Minimization Problems

**Note:** Maximization problem can be solved by minimization. Consider

$$\begin{aligned} & \text{maximize} \quad f_0(x) \\ & \text{subject to} \quad f_i(x) \leq 0, \quad i = 1, \dots, m \\ & \quad \quad \quad h_i(x) = 0, \quad i = 1, \dots, p, \end{aligned}$$

It can be solved by solving a corresponding minimization problem with  $-f_0(x)$

Optimization problem (Standard Form):

$$\begin{aligned} & \text{minimize} \quad -f_0(x) \\ & \text{subject to} \quad f_i(x) \leq 0, \quad i = 1, \dots, m \\ & \quad \quad \quad h_i(x) = 0, \quad i = 1, \dots, p \end{aligned}$$


- Note the constraints remain **same**!
- Obviously, the **optimal value**  $p^*$  is

$$p^* = \sup \{ f_0(x) \mid f_i(x) \leq 0, \quad i = 1, \dots, m, \quad h_i(x) = 0, \quad i = 1, \dots, p \}$$

*Handwritten notes:* An orange arrow points from the word "sup" in the equation above to the word "sup" in the equation below. Another orange arrow points from the expression  $-f_0(x)$  in the minimization problem above to the expression  $f_0(x)$  in the set of the equation below. A large orange curly brace is drawn to the right of the constraints in the minimization problem above.

## Equivalent Optimization Problems

Two problems are **equivalent** if from a solution of the one, a solution of the other can be found



## Equivalent Optimization Problems

Two problems are **equivalent** if from a solution of the one, a solution of the other can be found

**Example:** Consider the following problem

$$\begin{aligned} &\text{minimize} && \alpha_0 f_0(x) \\ &\text{subject to} && \alpha_i f_i(x) \leq 0, \quad i = 1, \dots, m \\ & && \beta_i h_i(x) = 0, \quad i = 1, \dots, p \end{aligned}$$

where  $\alpha_i > 0, i = 0, \dots, m, \quad \beta_i \neq 0, i = 1, \dots, p.$

$\alpha_0 > 0$

$\alpha_i f_i(x) \leq 0$

$\Rightarrow f_i(x) \leq 0$

$\beta_i h_i(x) = 0$

$\Rightarrow h_i(x) = 0$

$\Rightarrow x^*$  is opt val  
 $p^* = \alpha_0 f_0(x^*)$   
don't forget

Change?

min  $\alpha_0 f_0(x)$

$\Rightarrow$  min  $f_0(x)$

Only optimal value.  
Optimal pt remains same.



## Equivalent Optimization Problems

Two problems are **equivalent** if from a solution of the one, a solution of the other can be found

**Example:** Consider the following problem

$$\begin{aligned} & \text{minimize} && \alpha_0 f_0(x) \\ & \text{subject to} && \alpha_i f_i(x) \leq 0, \quad i = 1, \dots, m \\ & && \beta_i h_i(x) = 0, \quad i = 1, \dots, p \end{aligned}$$

where  $\alpha_i > 0, i = 0, \dots, m, \quad \beta_i \neq 0, i = 1, \dots, p$ .

**Note:**  $\alpha_i$  can be removed from inequality, and  $\beta_i$  can be removed from equality constraint. Minimizing  $\alpha_0 f_0(x)$  is same as minimizing  $f_0(x)$ . Hence it is **equivalent** to

## Equivalent Optimization Problems

Two problems are **equivalent** if from a solution of the one, a solution of the other can be found


**Example:** Consider the following problem

$$\begin{aligned} &\text{minimize} && \alpha_0 f_0(x) \\ &\text{subject to} && \alpha_i f_i(x) \leq 0, \quad i = 1, \dots, m \\ & && \beta_i h_i(x) = 0, \quad i = 1, \dots, p \end{aligned}$$

where  $\alpha_i > 0, i = 0, \dots, m, \quad \beta_i \neq 0, i = 1, \dots, p$ .

**Note:**  $\alpha_i$  can be removed from inequality, and  $\beta_i$  can be removed from equality constraint. Minimizing  $\alpha_0 f_0(x)$  is same as minimizing  $f_0(x)$ . Hence it is **equivalent** to

$$\begin{aligned} &\text{minimize} && f_0(x) \\ &\text{subject to} && f_i(x) \leq 0, \quad i = 1, \dots, m \\ & && h_i(x) = 0, \quad i = 1, \dots, p \end{aligned}$$



## Equivalent Problems: Change of Variables


Suppose  $\phi : \mathbb{R}^n \rightarrow \mathbb{R}^n$  is one-one.

## Equivalent Problems: Change of Variables

Suppose  $\phi : \mathbb{R}^n \rightarrow \mathbb{R}^n$  is one-one.

- Define  $\tilde{f}_i$  as

$$\tilde{f}_i(z) = f_i(\phi(z)), \quad i = 0, \dots, m$$



## Equivalent Problems: Change of Variables


Suppose  $\phi : \mathbb{R}^n \rightarrow \mathbb{R}^n$  is one-one.

- Define  $\tilde{f}_i$  as

$$\tilde{f}_i(z) = f_i(\phi(z)), \quad i = 0, \dots, m$$

- Define  $\tilde{h}_i$  as

$$\tilde{h}_i(z) = h_i(\phi(z)), \quad i = 1, \dots, p$$



## Equivalent Problems: Change of Variables

Suppose  $\phi : \mathbb{R}^n \rightarrow \mathbb{R}^n$  is one-one.

- Define  $\tilde{f}_i$  as


$$\tilde{f}_i(z) = f_i(\phi(z)), \quad i = 0, \dots, m$$

- Define  $\tilde{h}_i$  as

$$\tilde{h}_i(z) = h_i(\phi(z)), \quad i = 1, \dots, p$$

Consider

$$\begin{aligned} & \text{minimize } \tilde{f}_0(z) \\ & \text{subject to } \tilde{f}_i(z) \leq 0, \quad i = 1, \dots, m \\ & \quad \quad \quad \tilde{h}_i(z) = 0, \quad i = 1, \dots, p \end{aligned}$$



## Equivalent Problems: Change of Variables

Suppose  $\phi : \mathbb{R}^n \rightarrow \mathbb{R}^n$  is one-one.

- Define  $\tilde{f}_i$  as

$$\tilde{f}_i(z) = f_i(\phi(z)), \quad i = 0, \dots, m$$


- Define  $\tilde{h}_i$  as

$$\tilde{h}_i(z) = h_i(\phi(z)), \quad i = 1, \dots, p$$

Consider

$$\begin{aligned} &\text{minimize} && \tilde{f}_0(z) \\ &\text{subject to} && \tilde{f}_i(z) \leq 0, \quad i = 1, \dots, m \\ &&& \tilde{h}_i(z) = 0, \quad i = 1, \dots, p \end{aligned}$$

- If  $z$  solves above, then  $x = \phi(z)$  solves the standard optimization problem
- Similarly, if  $x$  solves original opt problem, then  $z = \phi^{-1}(x)$  solves above



## Equivalent Problems: Transformation of objective and constraint function

- Suppose  $\psi_0 : \mathbb{R} \rightarrow \mathbb{R}$  is monotone increasing



## Equivalent Problems: Transformation of objective and constraint function

- Suppose  $\psi_0: \mathbb{R} \rightarrow \mathbb{R}$  is monotone increasing
- $\psi_1, \dots, \psi_m: \mathbb{R} \rightarrow \mathbb{R}$  satisfy

## Equivalent Problems: Transformation of objective and constraint function

- Suppose  $\psi_0 : \mathbb{R} \rightarrow \mathbb{R}$  is **monotone increasing**
- $\psi_1, \dots, \psi_m : \mathbb{R} \rightarrow \mathbb{R}$  satisfy

$$\psi_i(u) \leq 0, \quad \text{if and only if } u \leq 0$$

## Equivalent Problems: Transformation of objective and constraint function

- Suppose  $\psi_0 : \mathbb{R} \rightarrow \mathbb{R}$  is **monotone increasing**
- $\psi_1, \dots, \psi_m : \mathbb{R} \rightarrow \mathbb{R}$  satisfy

$$\psi_i(u) \leq 0, \quad \text{if and only if } u \leq 0$$

- $\psi_{m+1}, \dots, \psi_{m+p} : \mathbb{R} \rightarrow \mathbb{R}$  satisfy

## Equivalent Problems: Transformation of objective and constraint function

- Suppose  $\psi_0 : \mathbb{R} \rightarrow \mathbb{R}$  is **monotone increasing**
- $\psi_1, \dots, \psi_m : \mathbb{R} \rightarrow \mathbb{R}$  satisfy

$$\psi_i(u) \leq 0, \quad \text{if and only if } u \leq 0$$

- $\psi_{m+1}, \dots, \psi_{m+p} : \mathbb{R} \rightarrow \mathbb{R}$  satisfy

$$\psi_i(u) = 0, \quad \text{if and only if } u = 0$$

- Define  $\tilde{f}_i$  as

## Equivalent Problems: Transformation of objective and constraint function

- Suppose  $\psi_0 : \mathbb{R} \rightarrow \mathbb{R}$  is **monotone increasing**
- $\psi_1, \dots, \psi_m : \mathbb{R} \rightarrow \mathbb{R}$  satisfy

$$\psi_i(u) \leq 0, \quad \text{if and only if } u \leq 0$$

- $\psi_{m+1}, \dots, \psi_{m+p} : \mathbb{R} \rightarrow \mathbb{R}$  satisfy

$$\psi_i(u) = 0, \quad \text{if and only if } u = 0$$

- Define  $\tilde{f}_i$  as

$$\tilde{f}_i(x) = \psi_i(f_i(x)), \quad i = 0, \dots, m$$

- Define  $\tilde{h}_i$  as

## Equivalent Problems: Transformation of objective and constraint function

- Suppose  $\psi_0 : \mathbb{R} \rightarrow \mathbb{R}$  is **monotone increasing**
- $\psi_1, \dots, \psi_m : \mathbb{R} \rightarrow \mathbb{R}$  satisfy

$$\psi_i(u) \leq 0, \quad \text{if and only if } u \leq 0$$

- $\psi_{m+1}, \dots, \psi_{m+p} : \mathbb{R} \rightarrow \mathbb{R}$  satisfy

$$\psi_i(u) = 0, \quad \text{if and only if } u = 0$$

- Define  $\tilde{f}_i$  as

$$\tilde{f}_i(x) = \psi_i(f_i(x)), \quad i = 0, \dots, m$$

- Define  $\tilde{h}_i$  as

$$\tilde{h}_i(x) = \psi_{m+i}(h_i(x)), \quad i = 1, \dots, p$$

The **associated problem** is

## Equivalent Problems: Transformation of objective and constraint function

- Suppose  $\psi_0 : \mathbb{R} \rightarrow \mathbb{R}$  is **monotone increasing**
- $\psi_1, \dots, \psi_m : \mathbb{R} \rightarrow \mathbb{R}$  satisfy

$$\psi_i(u) \leq 0, \quad \text{if and only if } u \leq 0$$

- $\psi_{m+1}, \dots, \psi_{m+p} : \mathbb{R} \rightarrow \mathbb{R}$  satisfy

$$\psi_i(u) = 0, \quad \text{if and only if } u = 0$$

- Define  $\tilde{f}_i$  as

$$\tilde{f}_i(x) = \psi_i(f_i(x)), \quad i = 0, \dots, m$$

- Define  $\tilde{h}_i$  as

$$\tilde{h}_i(x) = \psi_{m+i}(h_i(x)), \quad i = 1, \dots, p$$

The **associated problem** is

$$\begin{aligned} & \text{minimize} && \tilde{f}_0(x) \\ & \text{subject to} && \tilde{f}_i(x) \leq 0, \quad i = 1, \dots, m \\ & && \tilde{h}_i(x) = 0, \quad i = 1, \dots, p \end{aligned}$$

# Equivalent Problems: Transformation of objective and constraint function

- Suppose  $\psi_0 : \mathbb{R} \rightarrow \mathbb{R}$  is **monotone increasing**
- $\psi_1, \dots, \psi_m : \mathbb{R} \rightarrow \mathbb{R}$  satisfy

$$\psi_i(u) \leq 0, \quad \text{if and only if } u \leq 0$$

- $\psi_{m+1}, \dots, \psi_{m+p} : \mathbb{R} \rightarrow \mathbb{R}$  satisfy

$$\psi_i(u) = 0, \quad \text{if and only if } u = 0$$

- Define  $\tilde{f}_i$  as

$$\tilde{f}_i(x) = \psi_i(f_i(x)), \quad i = 0, \dots, m$$

- Define  $\tilde{h}_i$  as

$$\tilde{h}_i(x) = \psi_{m+i}(h_i(x)), \quad i = 1, \dots, p$$

The **associated problem** is

$$\begin{cases} \text{minimize} & \tilde{f}_0(x) \\ \text{subject to} & \tilde{f}_i(x) \leq 0, \quad i = 1, \dots, m \\ & \tilde{h}_i(x) = 0, \quad i = 1, \dots, p \end{cases}$$

Handwritten notes:

- $\min_{x \geq 0} f_0(x)$
- s.t.  $f_1(x) \leq 0$
- $f_0 = \frac{1}{2}(x-2)^2$
- Std. Opt. Prob.
- $f^* = f_0(x^*)$

Handwritten notes:

- $\min_x f_0(x)$
- s.t.  $f_i \leq 0$
- $f_i = 0$

Handwritten notes:

- $\psi_i(f_i) \leq 0$
- $p_1$
- $p_2$
- $p_3$



## Equivalent Problems: Slack Variables

Given the optimization problem in standard form

Optimization problem (Standard Form):

$$\begin{aligned} & \text{minimize} && f_0(x) \\ & \text{subject to} && f_i(x) \leq 0, i = 1, \dots, m \\ & && h_i(x) = 0, i = 1, \dots, p \end{aligned}$$

equality constraint and  $\geq 0$

$$-f_i(x) \geq 0$$

Quiz: Is it possible to replace inequality constraints by equality constraints and non-negativity  $s_i(x)$  constraints?

$$f_i + s_i = 0$$

$$s_i \geq 0$$

$$h_i(x) = 0$$

$$f_i(x) \leq 0$$

## Equivalent Problems: Slack Variables

Given the optimization problem in standard form

Optimization problem (Standard Form):

$$\begin{aligned} & \text{minimize} && f_0(x) \\ & \text{subject to} && f_i(x) \leq 0, i = 1, \dots, m \\ & && h_i(x) = 0, i = 1, \dots, p \end{aligned}$$

**Quiz:** Is it possible to replace inequality constraints by equality constraints and **non-negativity** constraints?

**Ans:** Yes. **Key observation** is:  $f_i(x) \leq 0$ , **if and only if** there is an  $s_i \geq 0$  such that  $f_i(x) + s_i = 0$

## Equivalent Problems: Slack Variables

Given the optimization problem in standard form

Optimization problem (Standard Form):

$$\begin{aligned} & \text{minimize} && f_0(x) \\ & \text{subject to} && f_i(x) \leq 0, i = 1, \dots, m \\ & && h_i(x) = 0, i = 1, \dots, p \end{aligned}$$

Quiz: Is it possible to replace inequality constraints by equality constraints and non-negativity constraints?

Ans: Yes. Key observation is:  $f_i(x) \leq 0$ , if and only if there is an  $s_i \geq 0$  such that  $f_i(x) + s_i = 0$

$$\begin{aligned} & \text{minimize} && f_0(x) \\ & \text{subject to} && s_i \geq 0, \quad i = 1, \dots, m \\ & && f_i(x) + s_i = 0, \quad i = 1, \dots, m \\ & && h_i(x) = 0, \quad i = 1, \dots, p \end{aligned}$$

Note: Here  $s_i$  are called slack variable. Is this equivalent?

## Convex Optimization Problem in Standard Form

### Convex Optimization Problem (Standard Form):

$$\begin{aligned} & \text{minimize} && f_0(x) \\ & \text{subject to} && \underline{f_i(x)} \leq 0, \quad i = 1, \dots, m \\ & && \underline{a_i^T x} = b_i, \quad i = 1, \dots, p \end{aligned}$$

where  $f_0, \dots, f_m$  are **convex** functions.

*Opt. Book*

*$h_i(x)$  is affine  
 $\Rightarrow$   $h_i$  is conv*

Comparing this with the standard form


### Optimization problem (Standard Form):

$$\begin{aligned} & \text{minimize} && f_0(x) \\ & \text{subject to} && f_i(x) \leq 0, \quad i = 1, \dots, m \\ & && h_i(x) = 0, \quad i = 1, \dots, p \end{aligned}$$

- objective function **must be convex**
- inequality constraint functions **must be convex**
- equality constraint functions  $h_i(x) = a_i^T x - b_i$  **must be affine**

## Convex Optimization Problem

Consider the following optimization problem with  $x \in \mathbb{R}^2$

$$\begin{array}{ll}\text{maximize} & f_0(x) = x_1^2 + x_2^2 \\ \text{subject to} & f_1(x) = x_1/(1 + x_2^2) \leq 0 \\ & h_1(x) = (x_1 + x_2)^2 = 0,\end{array}$$


which is in standard form.

# Convex Optimization Problem

Consider the following optimization problem with  $x \in \mathbb{R}^2$

$$\begin{aligned} &\text{maximize} && f_0(x) = x_1^2 + x_2^2 \\ &\text{subject to} && f_1(x) = x_1/(1+x_2^2) \leq 0 \\ &&& h_1(x) = (x_1+x_2)^2 = 0, \end{aligned}$$

which is in standard form.

**Quiz:** Is this problem a convex optimization problem?

$$\begin{aligned} (x_1+x_2)^2 &= 0 \\ \iff \\ x_1+x_2 &= 0 \end{aligned}$$

$$\begin{aligned} \frac{x_1}{1+x_2^2} &\leq 0 \\ \iff \\ \frac{x_1}{1+x_2^2} &\geq 0 \\ \iff \\ x_1 &\leq 0 \end{aligned}$$

Observation

- ①  $h_1(x)$  is not affine
  - ② Is  $f_0$  conv?
  - ③ Is  $f_1$  conv?
- Construct Hessian  
& check  
aTH  $x \geq 0$   
 $x_2$

## Convex Optimization Problem

Consider the following optimization problem with  $x \in \mathbb{R}^2$

$$\begin{aligned} &\text{maximize} && f_0(x) = x_1^2 + x_2^2 \\ &\text{subject to} && f_1(x) = x_1/(1 + x_2^2) \leq 0 \\ &&& h_1(x) = (x_1 + x_2)^2 = 0, \end{aligned}$$

which is in standard form.

Quiz: Is this problem a convex optimization problem? **Ans: No**

Quiz: Can you rewrite this in convex optimization problem?

## Convex Optimization Problem

Consider the following optimization problem with  $x \in \mathbb{R}^2$

$$\begin{aligned} &\text{maximize} && f_0(x) = x_1^2 + x_2^2 \\ &\text{subject to} && f_1(x) = x_1/(1 + x_2^2) \leq 0 \\ &&& h_1(x) = (x_1 + x_2)^2 = 0, \end{aligned}$$

which is in standard form.

Quiz: Is this problem a convex optimization problem? Ans: No

Quiz: Can you rewrite this in convex optimization problem? Ans: Yes

$$\begin{aligned} &\text{maximize} && f_0(x) = x_1^2 + x_2^2 \\ &\text{subject to} && f_1(x) = x_1 \leq 0 \\ &&& h_i(x) = x_1 + x_2 = 0, \end{aligned}$$

Note: This is now a convex optimization problem



## Convex Optimization Problem: Local Optima = Global Optima


**Fact:** For a convex optimization problem, any local optima is a global optima

**Ans:** Proof on chalkboard!

**Scratch Space** : For Convex opt : Local opt = Global opt.

Suppose that  $x$  is locally optimal  $\Rightarrow x$  is feasible and  $B_R(x)$

$$f_0(x) = \inf \{ f_0(z) \mid z \text{ is feasible, } \|z-x\|_2 \leq R \}$$



for some  $R > 0$ .

Assume on contrary: Spse  $x$  is not globally optimally optimal,

i.e., there is a feasible  $y$  for which  $f_0(y) < f_0(x)$

Evidently,  $\|y-x\|_2 > R \rightarrow \text{#}$

Consider the point  $z$  given by  $\theta = \frac{R}{2\|y-x\|_2} < \frac{1}{2}$

$$z = (1-\theta)x + \theta y$$

## Scratch Space

Look at  $z - x$ :  $= -\theta x + \theta y$

We have  $\|z - x\|_2^2 = (z - x)^T (z - x) = (-\theta x + \theta y)^T (-\theta x + \theta y) = \theta^2 (y - x)^T (y - x)$

$$= \theta^2 \|y - x\|_2^2 = \frac{R^2}{2^2 \|y - x\|_2^2} \cdot \|y - x\|_2^2 = \frac{R}{2} < \underline{R} \Rightarrow z \in B_R(x)$$

Feasible set convex  $\Rightarrow z = (1 - \theta)x + \theta y \in \text{feasible set}.$

$$f_0(z) = f_0((1 - \theta)x + \theta y) \leq (1 - \theta)f_0(x) + \theta \underline{f_0(y)}$$
$$\leq (1 - \theta)f_0(x) + \theta \underline{f_0(x)}$$

$$\Rightarrow z \in B_R(x), \quad z \text{ is feasible, } f_0(z) \leq f_0(x)$$

$\Rightarrow$  it contradicts that  $x$  was locally optimal

## Scratch Space

Hence our assumpt. that  $x$  is not globally opt. is wrong, i.e.,  $x$  is globally optimal.

$f$  diff &  $C \nabla x$

$$f(y) \geq f(x) + \nabla f(x)(y-x) \rightarrow \textcircled{1}$$

$\forall x, y$

If  $x^*$  is locally optimal

$$\Rightarrow \nabla f(x^*) = 0$$

from  $\textcircled{1} \Rightarrow$

$\Rightarrow f(y) \geq f(x^*) \quad \forall y \in \text{feasible set}$   
 $\Rightarrow x^*$  is globally optimal.

## Convex Optimization Problem: Optimality Criteria

**Fact:** If  $f_0$  in a convex optimization problem is differentiable, then the point  $x$  is optimal if  $f$

$$\nabla f_0(x)^T (y - x) \geq 0 \quad \text{for all } y \in X$$

**Proof:** On chalkboard!

$$f(y) \geq f(x) + \underbrace{\nabla f(x)^T (y - x)}_{\forall y}$$

$$\text{If } \underline{x \text{ is optimal}} \Rightarrow \underline{f(y) \geq f(x)} \quad \forall y$$

$$b \geq c -$$

## Scratch Space

$X = \text{feasible set}$

Spse  $x \in X$ . and satisfies  $\nabla f_0(x)^T(y-x) \geq 0$

Since  $f$  is cvx & diff: 1st order.

$$f_0(y) \geq f_0(x) + \underbrace{\nabla f_0(x)^T(y-x)}_{\geq 0}$$

$$f_0(y) \geq f_0(x) \quad ?$$

dropping a positive term keeps the ineq.

$\Rightarrow x$  is optimal.