

Motivation

Problem: Find a **linear** decision surface (Hyperplane) that can separate two classes, and has the largest distance between border line classes.

- borderline class (sample) = **support vectors**
- largest distance or gap = **margin**

¹Lemma-1 in <https://www.svms.org/tutorials/Burges1998.pdf>

Motivation

Problem: Find a **linear** decision surface (Hyperplane) that can separate two classes, and has the largest distance between border line classes.

- borderline class (sample) = **support vectors**
- largest distance or gap = **margin**

- Assume that a linear classifier exists
- If the convex hull of two classes form non-intersecting convex sets, then there exists a separating hyperplane¹
- What if the data is **not** linearly separable?


¹Lemma-1 in <https://www.svms.org/tutorials/Burges1998.pdf>

Motivation

Problem: Find a **linear** decision surface (Hyperplane) that can separate two classes, and has the largest distance between border line classes.


- borderline class (sample) = **support vectors**
- largest distance or gap = **margin**

- Assume that a linear classifier exists
- If the convex hull of two classes form non-intersecting convex sets, then there exists a separating hyperplane¹
- What if the data is **not** linearly separable?




¹Lemma-1 in <https://www.svms.org/tutorials/Burges1998.pdf>

When the Data is not linearly separable




When the Data is not linearly separable



- Data is **not** linearly separable. That is, there **does not exist** a hyperplane that **separates** two classes


When the Data is not linearly separable



- Data is **not** linearly separable. That is, there **does not exist** a hyperplane that **separates** two classes

- After mapping the data is now **linearly separable**!

When the Data is not linearly separable



- Data is **not** linearly separable. That is, there **does not exist** a hyperplane that **separates** two classes
- Such a mapping is achieved using clever so-called **Kernel trick**!
- Note dimension of the space increases: **price to pay for linear separation**
- After mapping the data is now **linearly separable**!

History of SVM

- SVM have long history since 1960

History of SVM

- SVM have long history since 1960
- Use of SVMs have grown since...

History of SVM

- SVM have long history since 1960
- Use of SVMs have grown since...
- Main paper: "A training algorithm for optimal margin classifiers", by Boser, Guyon, Vapnik

History of SVM


- SVM have long history since 1960
- Use of SVMs have grown since...
- Main paper: "A training algorithm for optimal margin classifiers", by Boser, Guyon, Vapnik
- It was discovered in 1960s, but its use started only around 1980s when Vapnik moved to USA

History of SVM

- SVM have long history since 1960
- Use of SVMs have grown since...
- Main paper: "A training algorithm for optimal margin classifiers", by Boser, Guyon, Vapnik
- It was discovered in 1960s, but its use started only around 1980s when Vapnik moved to USA
- Vapnik's initial papers were rejected in NeurIPS!



History of SVM

- SVM have long history since 1960
- Use of SVMs have grown since...
- Main paper: "A training algorithm for optimal margin classifiers", by Boser, Guyon, Vapnik
- It was discovered in 1960s, but its use started only around 1980s when Vapnik moved to USA
- Vapnik's initial papers were rejected in NeurIPS!



History of SVM

- SVM have long history since 1960
- Use of SVMs have grown since...
- Main paper: "A training algorithm for optimal margin classifiers", by Boser, Guyon, Vapnik
- It was discovered in 1960s, but its use started only around 1980s when Vapnik moved to USA
- Vapnik's initial papers were rejected in NeurIPS!



Representing the data

- SVMs are used for classifications and regressions

Representing the data

- SVMs are used for classifications and regressions
- How to represent data?

Representing the data

- SVMs are used for classifications and regressions
- How to represent data?
- A person aged 29 has BP of 110 is represented by a vector (110, 29)

Representing the data

- SVMs are used for classifications and regressions
- How to represent data?
- A person aged 29 has BP of 110 is represented by a vector $(110, 29)$
- vectors are tailed at origin $(0, 0)$


Representing the data

- SVMs are used for classifications and regressions
- How to represent data?
- A person aged 29 has BP of 110 is represented by a vector $(110, 29)$
- vectors are tailed at origin $(0, 0)$





Representing the data

- SVMs are used for classifications and regressions
- How to represent data?
- A person aged 29 has BP of 110 is represented by a vector (110, 29)
- vectors are tailed at origin (0, 0)





Patient id	Cholesterol (mg/dl)	Systolic BP (mmHg)	Age (years)	Tail of the vector	Arrow-head of the vector
1	150	110	35	(0,0,0)	(150, 110, 35)
2	250	120	30	(0,0,0)	(250, 120, 30)
3	140	160	65	(0,0,0)	(140, 160, 65)
4	300	180	45	(0,0,0)	(300, 180, 45)

Recall: Equation of Hyperplane





Recall: Equation of Hyperplane

- Let x_0 be any **fixed point** in plane P



Recall: Equation of Hyperplane



- Let x_0 be any **fixed point** in plane P
- Let w be a vector **perpendicular** to P



Recall: Equation of Hyperplane

- Let x_0 be any **fixed point** in plane P
- Let w be a vector **perpendicular** to P
- Let x be **any** point in P , then

$$(x - x_0) \perp w \implies (x - x_0)^T w = 0$$





Recall: Equation of Hyperplane

- Let x_0 be any **fixed point** in plane P
- Let w be a vector **perpendicular** to P
- Let x be **any** point in P , then

$$(x - x_0) \perp w \implies (x - x_0)^T w = 0$$

- Let $b = -w^T x_0$, then **equation of plane** is

$$w^T x + b = 0 \quad \text{or} \quad w \cdot x + b = 0$$



Recall: Equation of Hyperplane



- Let x_0 be any **fixed point** in plane P
- Let w be a vector **perpendicular** to P
- Let x be **any** point in P , then

$$(x - x_0) \perp w \implies (x - x_0)^T w = 0$$

- Let $b = -w^T x$, then **equation of plane** is

$$w^T x + b = 0 \quad \text{or} \quad w \cdot x + b = 0$$

- Changing b , we get parallel planes, see figure on the right



Recall: Equation of Hyperplane



- Let x_0 be any **fixed point** in plane P
- Let w be a vector **perpendicular** to P
- Let x be **any** point in P , then

$$(x - x_0) \perp w \implies (x - x_0)^T w = 0$$

- Let $b = -w^T x$, then **equation of plane** is

$$w^T x + b = 0 \quad \text{or} \quad w \cdot x + b = 0$$

- Changing b , we get parallel planes, see figure on the right
 - increasing b moves in direction of w



Recall: Equation of Hyperplane



- Let x_0 be any **fixed point** in plane P
- Let w be a vector **perpendicular** to P
- Let x be **any** point in P , then

$$(x - x_0) \perp w \implies (x - x_0)^T w = 0$$

- Let $b = -w^T x$, then **equation of plane** is

$$w^T x + b = 0 \quad \text{or} \quad w \cdot x + b = 0$$

- Changing b , we get parallel planes, see figure on the right
 - increasing b moves in direction of w
 - decreasing b moves in direction of $-w$



Recall: Equation of Hyperplane

- Let x_0 be any **fixed point** in plane P
- Let w be a vector **perpendicular** to P
- Let x be **any** point in P , then

$$(x - x_0) \perp w \implies (x - x_0)^T w = 0$$



- Let $b = -w^T x_0$, then **equation of plane** is

$$w^T x + b = 0 \quad \text{or} \quad w \cdot x + b = 0$$

- Changing b , we get parallel planes, see figure on the right

- increasing b moves in direction of w
- decreasing b moves in direction of $-w$

- **Distance between parallel planes** $w^T x + b_1 = 0$ and $w^T x + b_2 = 0$ is



Recall: Equation of Hyperplane

- Let x_0 be any **fixed point** in plane P
- Let w be a vector **perpendicular** to P
- Let x be **any** point in P , then

$$(x - x_0) \perp w \implies (x - x_0)^T w = 0$$

- Let $b = -w^T x$, then **equation of plane** is



$$w^T x + b = 0 \quad \text{or} \quad w \cdot x + b = 0$$

- Changing b , we get parallel planes, see figure on the right

- increasing b moves in direction of w
- decreasing b moves in direction of $-w$

- **Distance between parallel planes** $w^T x + b_1 = 0$ and $w^T x + b_2 = 0$ is

$$D = |b_1 - b_2| / \|w\|_2$$



SVMs for two class classification

- Consider the data:

$$X = \{x_1, x_2, \dots, x_m \in \mathbb{R}^n\}$$

$$Y = \{y_1, y_2, \dots, y_m \in \{+1, -1\}\}$$

SVMs for two class classification

- Consider the data:

$$X = \{x_1, x_2, \dots, x_m \in \mathbb{R}^n\}$$

$$Y = \{y_1, y_2, \dots, y_m \in \{+1, -1\}\}$$

- Here Y is label. It takes $+1$ or -1 values. It is a **two** class problem


SVMs for two class classification

- Consider the data:

$$X = \{x_1, x_2, \dots, x_m \in \mathbb{R}^n\}$$

$$Y = \{y_1, y_2, \dots, y_m \in \{+1, -1\}\}$$

- Here Y is label. It takes $+1$ or -1 values. It is a **two** class problem



SVMs for two class classification


- Consider the data:

$$X = \{x_1, x_2, \dots, x_m \in \mathbb{R}^n\}$$

$$Y = \{y_1, y_2, \dots, y_m \in \{+1, -1\}\}$$

Goal: Find a classifier to separate negative instances from positive ones

- Here Y is label. It takes $+1$ or -1 values. It is a **two** class problem



SVMs for two class classification

- Consider the data:


$$X = \{x_1, x_2, \dots, x_m \in \mathbb{R}^n\}$$

$$Y = \{y_1, y_2, \dots, y_m \in \{+1, -1\}\}$$

- Here Y is label. It takes $+1$ or -1 values. It is a **two** class problem

Goal: Find a classifier to separate negative instances from positive ones

- As shown in figure, there exists infinite such hyperplanes (classifiers)



SVMs for two class classification

- Consider the data:


$$X = \{x_1, x_2, \dots, x_m \in \mathbb{R}^n\}$$

$$Y = \{y_1, y_2, \dots, y_m \in \{+1, -1\}\}$$

- Here Y is label. It takes $+1$ or -1 values. It is a **two** class problem

Goal: Find a classifier to separate negative instances from positive ones

- As shown in figure, there exists infinite such hyperplanes (classifiers)
- Two hyperplanes (shown as dotted) pass through boundary points (support vectors)(yellow)



SVMs for two class classification


- Consider the data:

$$X = \{x_1, x_2, \dots, x_m \in \mathbb{R}^n\}$$

$$Y = \{y_1, y_2, \dots, y_m \in \{+1, -1\}\}$$

- Here Y is label. It takes $+1$ or -1 values. It is a **two** class problem

Goal: Find a classifier to separate negative instances from positive ones



- As shown in figure, there exists infinite such hyperplanes (classifiers)
- Two hyperplanes (shown as dotted) pass through boundary points (support vectors)(yellow)
- SVMs finds hyperplanes that maximizes the gap between data points on boundary

SVMs for two class classification


- Consider the data:

$$X = \{x_1, x_2, \dots, x_m \in \mathbb{R}^n\}$$

$$Y = \{y_1, y_2, \dots, y_m \in \{+1, -1\}\}$$


- Here Y is label. It takes $+1$ or -1 values. It is a **two** class problem

Goal: Find a classifier to separate negative instances from positive ones




- As shown in figure, there exists infinite such hyperplanes (classifiers)
- Two hyperplanes (shown as dotted) pass through boundary points (support vectors)(yellow)
- SVMs finds hyperplanes that maximizes the gap between data points on boundary
- If the support vectors are noisy, SVMs wont work well!

Particular Form of Hyperplanes Passing Through Support Vectors



Particular Form of Hyperplanes Passing Through Support Vectors




- **Quiz:** Why dotted planes through support vectors are

$$w \cdot x + b = -1 \quad (1)$$

$$w \cdot x + b = 1 \quad (2)$$

Particular Form of Hyperplanes Passing Through Support Vectors



- If we move c distance along w , we get to support vector of **red** class, then hyperplane is

$$w \cdot x + b = c \quad (4)$$

- **Quiz:** Why dotted planes through support vectors are


$$w \cdot x + b = -1 \quad (1)$$

$$w \cdot x + b = 1 \quad (2)$$

- Let middle **red** plane be

$$w \cdot x + b = 0 \quad (3)$$

Particular Form of Hyperplanes Passing Through Support Vectors



- **Quiz:** Why dotted planes through support vectors are

$$w \cdot x + b = -1 \quad (1)$$

$$w \cdot x + b = 1 \quad (2)$$

- Let middle **red** plane be

$$w \cdot x + b = 0 \quad (3)$$


- If we move c distance along w , we get to support vector of **red** class, then hyperplane is

$$w \cdot x + b = c \quad (4)$$

- Similarly, if we move c distance along $-w$, we reach support vector of **blue** class

$$w \cdot x + b = -c \quad (5)$$

Particular Form of Hyperplanes Passing Through Support Vectors



- **Quiz:** Why dotted planes through support vectors are

$$w \cdot x + b = -1 \quad (1)$$

$$w \cdot x + b = 1 \quad (2)$$

- Let middle **red** plane be

$$w \cdot x + b = 0 \quad (3)$$

- If we move c distance along w , we get to support vector of **red** class, then hyperplane is

$$w \cdot x + b = c \quad (4)$$

- Similarly, if we move c distance along $-w$, we reach support vector of **blue** class

$$w \cdot x + b = -c \quad (5)$$


- Divide (3), (4), (5) by c to get

$$\tilde{w} \cdot x + \tilde{b} = 0$$

$$\tilde{w} \cdot x + \tilde{b} = -1$$

$$\tilde{w} \cdot x + \tilde{b} = 1$$

Particular Form of Hyperplanes Passing Through Support Vectors



- **Quiz:** Why dotted planes through support vectors are

$$w \cdot x + b = -1 \quad (1)$$

$$w \cdot x + b = 1 \quad (2)$$

- Let middle **red** plane be

$$w \cdot x + b = 0 \quad (3)$$

- If we move c distance along w , we get to support vector of **red** class, then hyperplane is

$$w \cdot x + b = c \quad (4)$$

- Similarly, if we move c distance along $-w$, we reach support vector of **blue** class

$$w \cdot x + b = -c \quad (5)$$

- Divide (3), (4), (5) by c to get


$$\tilde{w} \cdot x + \tilde{b} = 0$$

$$\tilde{w} \cdot x + \tilde{b} = -1$$


$$\tilde{w} \cdot x + \tilde{b} = 1$$

- Rename \tilde{w} by w , and \tilde{b} by b above!

SVM idea: Maximize the Margin




SVM idea: Maximize the Margin



Margin/Gap: Distance between parallel planes passing through support vectors


SVM idea: Maximize the Margin



Margin/Gap: Distance between parallel planes passing through support vectors

Goal: Maximize the margin!

SVM idea: Maximize the Margin



- We have


$$w \cdot x + b = -1$$

$$w \cdot x + b = 1$$

Margin/Gap: Distance between parallel planes passing through support vectors

Goal: Maximize the margin!

SVM idea: Maximize the Margin



Margin/Gap: Distance between parallel planes passing through support vectors

Goal: Maximize the margin!

- We have

$$w \cdot x + b = -1$$


$$w \cdot x + b = 1$$

or equivalently,

$$w \cdot x + b + 1 = 0$$

$$w \cdot x + b - 1 = 0$$

SVM idea: Maximize the Margin



Margin/Gap: Distance between parallel planes passing through support vectors

Goal: Maximize the margin!

- We have

$$w \cdot x + b = -1$$

$$w \cdot x + b = 1$$


or equivalently,

$$w \cdot x + b + 1 = 0$$

$$w \cdot x + b - 1 = 0$$

- Distance/Margin $D = 2/\|w\|$

SVM idea: Maximize the Margin



Margin/Gap: Distance between parallel planes passing through support vectors

Goal: Maximize the margin!

- We have

$$w \cdot x + b = -1$$

$$w \cdot x + b = 1$$

or equivalently,


$$w \cdot x + b + 1 = 0$$

$$w \cdot x + b - 1 = 0$$

- Distance/Margin $D = 2/\|w\|$
- To maximize the gap:

$$\text{minimize } \frac{1}{2} \|w\|^2$$

SVM idea: Maximize the Margin



Margin/Gap: Distance between parallel planes passing through support vectors

Goal: Maximize the margin!

- We have

$$w \cdot x + b = -1$$

$$w \cdot x + b = 1$$

or equivalently,

$$w \cdot x + b + 1 = 0$$


$$w \cdot x + b - 1 = 0$$

- Distance/Margin $D = 2/\|w\|$
- To maximize the gap:


$$\text{minimize } \frac{1}{2} \|w\|^2$$

- Are there constraints?

Impose Constraints, Optimization Model, Prediction




Impose Constraints, Optimization Model, Prediction



Goal: Impose constraints such that all the data points are correctly classified.


Impose Constraints, Optimization Model, Prediction



- For all the **blue data** to be correctly classified, we must have

Goal: Impose constraints such that all the data points are correctly classified.

Impose Constraints, Optimization Model, Prediction




Goal: Impose constraints such that all the data points are correctly classified.

- For all the **blue data** to be correctly classified, we must have

$$w \cdot x + b \leq -1, \text{ if } y_i = -1$$

Impose Constraints, Optimization Model, Prediction




Goal: Impose constraints such that all the data points are correctly classified.

- For all the **blue data** to be correctly classified, we must have

$$w \cdot x + b \leq -1, \text{ if } y_i = -1$$

- Similarly, to classify all **red data** correctly, we must have

Impose Constraints, Optimization Model, Prediction



Goal: Impose constraints such that all the data points are correctly classified.


- For all the **blue data** to be correctly classified, we must have

$$w \cdot x + b \leq -1, \text{ if } y_i = -1$$

- Similarly, to classify all **red data** correctly, we must have

$$w \cdot x + b \geq +1, \text{ if } y_i = +1$$

Impose Constraints, Optimization Model, Prediction



Goal: Impose constraints such that all the data points are correctly classified.

- For all the **blue data** to be correctly classified, we must have

$$w \cdot x + b \leq -1, \text{ if } y_i = -1$$


- Similarly, to classify all **red data** correctly, we must have

$$w \cdot x + b \geq +1, \text{ if } y_i = +1$$

- Equivalently, we have

$$y_i(w \cdot x + b) \geq 1$$

Impose Constraints, Optimization Model, Prediction



Goal: Impose constraints such that all the data points are correctly classified.

Optimization model of SVM (training):

$$\text{minimize } \frac{1}{2} \|\mathbf{w}\|^2$$

$$\text{subject to } y_i(\mathbf{w} \cdot \mathbf{x} + b) \geq 1, \quad i = 1, \dots, m.$$

- For all the **blue data** to be correctly classified, we must have

$$\mathbf{w} \cdot \mathbf{x} + b \leq -1, \text{ if } y_i = -1$$


- Similarly, to classify all **red data** correctly, we must have

$$\mathbf{w} \cdot \mathbf{x} + b \geq +1, \text{ if } y_i = +1$$

- Equivalently, we have

$$y_i(\mathbf{w} \cdot \mathbf{x} + b) \geq 1$$

Impose Constraints, Optimization Model, Prediction



Goal: Impose constraints such that all the data points are correctly classified.

Optimization model of SVM (training):

$$\underset{w}{\text{minimize}} \quad \frac{1}{2} \|w\|^2$$

$$\text{subject to } y_i(w \cdot x + b) \geq 1, \quad i = 1, \dots, m.$$

- For all the **blue data** to be correctly classified, we must have

$$w \cdot x + b \leq -1, \text{ if } y_i = -1$$

- Similarly, to classify all **red data** correctly, we must have

$$w \cdot x + b \geq +1, \text{ if } y_i = +1$$

- Equivalently, we have

$$y_i(w \cdot x + b) \geq 1$$

Prediction:

$$f(x) = \text{sign}(w \cdot x + b)$$

Dual Formulation of SVM Optimization Problem

Standard form:

$$\begin{array}{ll}\text{minimize} & \frac{1}{2} \|w\|^2 \\ \text{subject to} & -y_i(w \cdot x + b) + 1 \leq 0, \quad i = 1, \dots, m.\end{array}$$

Dual Formulation of SVM Optimization Problem

Standard form:

$$\begin{array}{ll}\text{minimize} & \frac{1}{2} \|w\|^2 \\ \text{subject to} & -y_i(w \cdot x + b) + 1 \leq 0, \quad i = 1, \dots, m.\end{array}$$

- ✓ This is primal problem in standard form

Dual Formulation of SVM Optimization Problem

Standard form:

$$\begin{aligned} & \text{minimize} && \frac{1}{2} \|w\|^2 \\ & \text{subject to} && -y_i(w \cdot x + b) + 1 \leq 0, \quad i = 1, \dots, m. \end{aligned}$$

- ✓ This is primal problem in standard form
- ✓ It is convex quadratic programming (QP) because objective is quadratic, and constraints are linear

Dual Formulation of SVM Optimization Problem

Standard form:

$$\begin{aligned} & \text{minimize} && \frac{1}{2} \|w\|^2 \\ & \text{subject to} && -y_i(w \cdot x + b) + 1 \leq 0, \quad i = 1, \dots, m. \end{aligned}$$

- ✓ This is primal problem in standard form
- ✓ It is convex quadratic programming (QP) because objective is quadratic, and constraints are linear
- ✓ The problem can be recast into dual form

Dual Formulation of SVM Optimization Problem

Standard form:

$$\begin{aligned} & \text{minimize} && \frac{1}{2} \|w\|^2 \\ & \text{subject to} && -y_i(w \cdot x + b) + 1 \leq 0, \quad i = 1, \dots, m. \end{aligned}$$

1. Define the Lagrangian

- ✓ This is primal problem in standard form
- ✓ It is convex quadratic programming (QP) because objective is quadratic, and constraints are linear
- ✓ The problem can be recast into dual form

Dual Formulation of SVM Optimization Problem

Standard form:

$$\begin{aligned} & \text{minimize} && \frac{1}{2} \|w\|^2 \\ & \text{subject to} && -y_i(w \cdot x + b) + 1 \leq 0, \quad i = 1, \dots, m. \end{aligned}$$

1. Define the Lagrangian

$$L(w, b, \lambda) = \underbrace{\frac{1}{2} \|w\|_2^2} + \sum_{i=1}^m \lambda_i \underbrace{(-y_i(w \cdot x + b) + 1)}$$

- ✓ This is primal problem in standard form
- ✓ It is convex quadratic programming (QP) because objective is quadratic, and constraints are linear
- ✓ The problem can be recast into dual form

Dual Formulation of SVM Optimization Problem

Standard form:

$$\begin{aligned} & \text{minimize} && \frac{1}{2} \|w\|^2 \\ & \text{subject to} && -y_i(w \cdot x + b) + 1 \leq 0, \quad i = 1, \dots, m. \end{aligned}$$

1. Define the **Lagrangian**

$$L(w, b, \lambda) = \frac{1}{2} \|w\|_2^2 + \sum_{i=1}^m \lambda_i (-y_i(w \cdot x + b) + 1)$$

2. The **dual function** is

- ✓ This is primal problem in standard form
- ✓ It is convex quadratic programming (QP) because objective is quadratic, and constraints are linear
- ✓ The problem can be recast into dual form

Dual Formulation of SVM Optimization Problem

Standard form:

$$\begin{aligned} & \text{minimize} && \frac{1}{2} \|w\|^2 \\ & \text{subject to} && -y_i(w \cdot x + b) + 1 \leq 0, \quad i = 1, \dots, m. \end{aligned}$$

1. Define the **Lagrangian**

$$L(w, b, \lambda) = \frac{1}{2} \|w\|_2^2 + \sum_{i=1}^m \lambda_i (-y_i(w \cdot x + b) + 1)$$

2. The **dual function** is

$$g(\lambda) = \inf_{w, b} L(w, b, \lambda)$$

- ✓ This is primal problem in standard form
- ✓ It is convex quadratic programming (QP) because objective is quadratic, and constraints are linear
- ✓ The problem can be recast into dual form

Dual Formulation of SVM Optimization Problem

Standard form:

$$\begin{aligned} & \underset{w, b}{\text{minimize}} && \frac{1}{2} \|w\|^2 \\ & \text{subject to} && -y_i(w \cdot x + b) + 1 \leq 0, \quad i = 1, \dots, m. \end{aligned}$$

1. Define the **Lagrangian**

$$L(w, b, \lambda) = \frac{1}{2} \|w\|^2 + \sum_{i=1}^m \lambda_i (-y_i(w \cdot x + b) + 1)$$

2. The **dual function** is

$$g(\lambda) = \inf_{w, b} L(w, b, \lambda)$$

3. $L(w, b, \lambda)$ **convex** in w, b , minima given by

$$\nabla_{w, b} L(w, b, \lambda) = 0$$

- ✓ This is primal problem in standard form
- ✓ It is convex quadratic programming (QP) because objective is quadratic, and constraints are linear
- ✓ The problem can be recast into dual form

SVM-Dual Formulation

1. Setting the derivative of L w.r.t b to zero,

SVM-Dual Formulation

1. Setting the derivative of L w.r.t b to **zero**,

$$\frac{\partial L(w, b, \lambda)}{\partial b} = 0 \implies \sum_{i=1}^m \lambda_i y_i = 0$$

SVM-Dual Formulation

1. Setting the derivative of L w.r.t b to **zero**,

$$\frac{\partial L(w, b, \lambda)}{\partial b} = 0 \implies \sum_{i=1}^m \lambda_i y_i = 0$$

2. Setting the derivative of L w.r.t. w to **zero**,

SVM-Dual Formulation

1. Setting the derivative of L w.r.t b to **zero**,

$$\frac{\partial L(w, b, \lambda)}{\partial b} = 0 \implies \sum_{i=1}^m \lambda_i y_i = 0$$

2. Setting the derivative of L w.r.t. w to **zero**,

$$\frac{\partial L(w, b, \lambda)}{\partial w} = 0 \implies w = \sum_{j=1}^m \lambda_j y_j x_j$$

SVM-Dual Formulation

1. Setting the derivative of L w.r.t b to **zero**,

$$\frac{\partial L(w, b, \lambda)}{\partial b} = 0 \implies \sum_{i=1}^m \lambda_i y_i = 0$$

2. Setting the derivative of L w.r.t. w to **zero**,

$$\frac{\partial L(w, b, \lambda)}{\partial w} = 0 \implies w = \sum_{j=1}^m \lambda_j y_j x_j$$

3. Substituting in $\underline{L(w, b, \lambda)} = \frac{1}{2} \|w\|_2^2 + \sum_{i=1}^m \lambda_i (-y_i (w \cdot x_i + b) + 1),$

SVM-Dual Formulation

1. Setting the derivative of L w.r.t b to **zero**,

$$\frac{\partial L(w, b, \lambda)}{\partial b} = 0 \implies \sum_{i=1}^m \lambda_i y_i = 0$$

2. Setting the derivative of L w.r.t. w to **zero**,

$$\frac{\partial L(w, b, \lambda)}{\partial w} = 0 \implies w = \sum_{j=1}^m \lambda_j y_j x_j$$

3. Substituting in $L(w, b, \lambda) = \frac{1}{2} \|w\|_2^2 + \sum_{i=1}^m \lambda_i (-y_i (w \cdot x_i + b) + 1)$,

$$g(\lambda) = \sum_{i=1}^m \lambda_i - \frac{1}{2} \sum_{i=1}^m \lambda_i \lambda_j y_i y_j (x_i \cdot x_j)$$

$$\frac{1}{2} \|w\|_2^2 = \frac{1}{2} w^T w = \frac{1}{2} \left(\sum_{i=1}^m \lambda_i y_i x_i \right)^T \left(\sum_{j=1}^m \lambda_j y_j x_j \right)$$

$$= \frac{1}{2} \sum_{i=1}^m \sum_{j=1}^m \lambda_i \lambda_j y_i y_j (x_i \cdot x_j)$$

$$= \begin{aligned} & - \sum \lambda_i \lambda_j y_i y_j (x_i \cdot x_j) \\ & + \sum \lambda_i + \sum_{i=1}^m \lambda_i b y_i \end{aligned} \xrightarrow{0}$$

because $\sum \lambda_i y_i = 0$

SVM-Dual Formulation

1. Setting the derivative of L w.r.t b to **zero**,

$$\frac{\partial L(w, b, \lambda)}{\partial b} = 0 \implies \sum_{i=1}^m \lambda_i y_i = 0 \quad \}$$

2. Setting the derivative of L w.r.t. w to **zero**,

$$\frac{\partial L(w, b, \lambda)}{\partial w} = 0 \implies w = \sum_{j=1}^m \lambda_j y_j x_j$$

3. Substituting in $L(w, b, \lambda) = \frac{1}{2} \|w\|_2^2 + \sum_{i=1}^m \lambda_i (-y_i (w \cdot x_i + b) + 1)$,

$$g(\lambda) = \sum_{i=1}^m \lambda_i - \frac{1}{2} \sum_{i=1}^m \lambda_i \lambda_j y_i y_j (x_i \cdot x_j)$$

SVM Dual Problem:

$$\text{maximize} \quad \sum_{i=1}^m \lambda_i - \frac{1}{2} \sum_{i=1}^m \lambda_i \lambda_j y_i y_j (x_i \cdot x_j)$$

$$\text{subject to} \quad \lambda_i \geq 0, \quad \sum_{i=1}^m \lambda_i y_i = 0, \quad i = 1, \dots, m$$

SVM Primal Versus Dual Problem

SVM Primal Problem:

$$\text{minimize } \frac{1}{2} \|w\|^2, \quad w \in \mathbb{R}^n$$

$$\text{subject to } -y_i(w \cdot x + b) + 1 \leq 0, \quad i = 1, \dots, m.$$

SVM Primal Versus Dual Problem

SVM Primal Problem:

$$\text{minimize } \frac{1}{2} \|w\|^2, \quad w \in \mathbb{R}^n$$

$$\text{subject to } -y_i(w \cdot x + b) + 1 \leq 0, \quad i = 1, \dots, m.$$

SVM Dual Problem:

$$\text{maximize } \sum_{i=1}^m \lambda_i - \frac{1}{2} \sum_{i,j=1}^m \lambda_i \lambda_j y_i y_j (x_i \cdot x_j)$$

$$\text{subject to } \lambda_i \geq 0, \quad \sum_{i=1}^m \lambda_i y_i = 0, \quad i = 1, \dots, m.$$

SVM Primal Versus Dual Problem

SVM Primal Problem:

$$\begin{aligned} &\text{minimize } \frac{1}{2} \|w\|^2, \quad w \in \mathbb{R}^n \\ &\text{subject to } -y_i(w \cdot x + b) + 1 \leq 0, \quad i = 1, \dots, m. \end{aligned}$$

Primal:

- Variables are $\{\underline{w_1}, \underline{w_2}, \dots, \underline{w_n}\}$

SVM Dual Problem:

$$\begin{aligned} &\text{maximize } \sum_{i=1}^m \lambda_i - \frac{1}{2} \sum_{i,j=1}^m \lambda_i \lambda_j y_i y_j (x_i \cdot x_j) \\ &\text{subject to } \lambda_i \geq 0, \quad \sum_{i=1}^m \lambda_i y_i = 0, \quad i = 1, \dots, m. \end{aligned}$$

SVM Primal Versus Dual Problem

SVM Primal Problem:

$$\begin{aligned} &\text{minimize } \frac{1}{2} \|w\|^2, \quad w \in \mathbb{R}^n \\ &\text{subject to } -y_i(w \cdot x + b) + 1 \leq 0, \quad i = 1, \dots, m. \end{aligned}$$

SVM Dual Problem:

$$\begin{aligned} &\text{maximize } \sum_{i=1}^m \lambda_i - \frac{1}{2} \sum_{i,j=1}^m \lambda_i \lambda_j y_i y_j (x_i \cdot x_j) \\ &\text{subject to } \lambda_i \geq 0, \quad \sum_{i=1}^m \lambda_i y_i = 0, \quad i = 1, \dots, m. \end{aligned}$$

Primal:

- Variables are $\{w_1, w_2, \dots, w_n\}$
- n is number of features

SVM Primal Versus Dual Problem

SVM Primal Problem:

$$\text{minimize } \frac{1}{2} \|w\|^2, \quad w \in \mathbb{R}^n$$

$$\text{subject to } -y_i(w \cdot x + b) + 1 \leq 0, \quad i = 1, \dots, m.$$

Primal:

- Variables are $\{w_1, w_2, \dots, w_n\}$
- n is number of features
- Primal is convex

SVM Dual Problem:

$$\text{maximize } \sum_{i=1}^m \lambda_i - \frac{1}{2} \sum_{i,j=1}^m \lambda_i \lambda_j y_i y_j (x_i \cdot x_j)$$

$$\text{subject to } \lambda_i \geq 0, \quad \sum_{i=1}^m \lambda_i y_i = 0, \quad i = 1, \dots, m.$$

$$\frac{1}{2} w^T I w$$

$P \geq 0$

constraint + objective
for all convex

SVM Primal Versus Dual Problem

SVM Primal Problem:

$$\begin{aligned} &\text{minimize } \frac{1}{2} \|w\|^2, \quad w \in \mathbb{R}^n \\ &\text{subject to } -y_i(w \cdot x + b) + 1 \leq 0, \quad i = 1, \dots, m. \end{aligned}$$

SVM Dual Problem:

$$\begin{aligned} &\text{maximize } \sum_{i=1}^m \lambda_i - \frac{1}{2} \sum_{i,j=1}^m \lambda_i \lambda_j y_i y_j (x_i \cdot x_j) \\ &\text{subject to } \lambda_i \geq 0, \quad \sum_{i=1}^m \lambda_i y_i = 0, \quad i = 1, \dots, m. \end{aligned}$$

Primal:

- Variables are $\{w_1, w_2, \dots, w_n\}$
- n is number of features
- Primal is convex

Dual:

- Variables are $\{\lambda_1, \lambda_2, \dots, \lambda_m\}$

SVM Primal Versus Dual Problem

SVM Primal Problem:

$$\begin{aligned} & \text{minimize } \frac{1}{2} \|w\|^2, \quad w \in \mathbb{R}^n \\ & \text{subject to } -y_i(w \cdot x + b) + 1 \leq 0, \quad i = 1, \dots, m. \end{aligned}$$

SVM Dual Problem:

$$\begin{aligned} & \text{maximize } \sum_{i=1}^m \lambda_i - \frac{1}{2} \sum_{i,j=1}^m \lambda_i \lambda_j y_i y_j (x_i \cdot x_j) \\ & \text{subject to } \lambda_i \geq 0, \quad \sum_{i=1}^m \lambda_i y_i = 0, \quad i = 1, \dots, m. \end{aligned}$$

Primal:

- Variables are $\{w_1, w_2, \dots, w_n\}$
- n is number of features
- Primal is convex

Dual:

- Variables are $\{\lambda_1, \lambda_2, \dots, \lambda_m\}$
- m is number of data samples

SVM Primal Versus Dual Problem

SVM Primal Problem:

$$\begin{aligned} & \text{minimize } \frac{1}{2} \|w\|^2, \quad w \in \mathbb{R}^n \\ & \text{subject to } -y_i(w \cdot x + b) + 1 \leq 0, \quad i = 1, \dots, m. \end{aligned}$$

SVM Dual Problem:

$$\begin{aligned} & \text{maximize } \sum_{i=1}^m \lambda_i - \frac{1}{2} \sum_{i,j=1}^m \lambda_i \lambda_j y_i y_j (x_i \cdot x_j) \\ & \text{subject to } \lambda_i \geq 0, \quad \sum_{i=1}^m \lambda_i y_i = 0, \quad i = 1, \dots, m. \end{aligned}$$

Handwritten notes:
An arrow points from the first term of the objective function to the primal variable w .
An arrow points from the second term of the objective function to the primal constraint.
The term $\lambda_i y_i$ in the constraint is circled.
Below the constraint, there is a handwritten note: $-x_i \cdot x_j$ and $(\partial \lambda_i)$.

Primal:

- Variables are $\{w_1, w_2, \dots, w_n\}$
- n is number of features
- Primal is convex

Dual:

- Variables are $\{\lambda_1, \lambda_2, \dots, \lambda_m\}$
- m is number of data samples
- Dual is convex

SVM Primal Versus Dual Problem

SVM Primal Problem:

$$\text{minimize } \frac{1}{2} \|w\|^2, \quad w \in \mathbb{R}^n$$

$$\text{subject to } -y_i(w \cdot x + b) + 1 \leq 0, \quad i = 1, \dots, m.$$

SVM Dual Problem:

$$\text{maximize } \sum_{i=1}^m \lambda_i - \frac{1}{2} \sum_{i,j=1}^m \lambda_i \lambda_j y_i y_j (x_i \cdot x_j)$$

$$\text{subject to } \lambda_i \geq 0, \quad \sum_{i=1}^m \lambda_i y_i = 0, \quad i = 1, \dots, m.$$



Primal:

- Variables are $\{w_1, w_2, \dots, w_n\}$
- n is number of features
- Primal is convex

Dual:

- Variables are $\{\lambda_1, \lambda_2, \dots, \lambda_m\}$
- m is number of data samples
- Dual is convex

Recommendation: Use dual when the number of samples m are significantly less relative to the number of features n , otherwise use primal.

Strong Duality and KKT Conditions for SVM

SVM Primal Problem:

$$\begin{aligned} &\text{minimize} && \frac{1}{2} \|w\|^2 \\ &\text{subject to} && -y_i(w \cdot x + b) + 1 \leq 0, \quad i = 1, \dots, m. \end{aligned}$$

Strong Duality and KKT Conditions for SVM

SVM Primal Problem:

$$\begin{aligned} & \text{minimize} && \frac{1}{2} \|w\|^2 \\ & \text{subject to} && -y_i(w \cdot x + b) + 1 \leq 0, \quad i = 1, \dots, m. \end{aligned}$$

Recall Slater's condition from slide 12

Slater's condition when some f_i are affine. There exists $x \in \text{int } \mathcal{D}$ with


$$f_i(x) \leq 0, \quad i = 1, \dots, k, \quad f_i(x) < 0, \quad i = k + 1, \dots, m, \quad Ax = b.$$

affine

non-affine

Strong Duality and KKT Conditions for SVM

SVM Primal Problem:

$$\begin{array}{ll}\text{minimize} & \frac{1}{2} \|w\|^2 \\ \text{subject to} & -y_i(w \cdot x + b) + 1 \leq 0, \quad i = 1, \dots, m.\end{array}$$


Recall Slater's condition from slide 12

Slater's condition when some f_i are affine. There exists $x \in \text{int } \mathcal{D}$ with

$$f_i(x) \leq 0, \quad i = 1, \dots, k, \quad f_i(x) < 0, \quad i = k + 1, \dots, m, \quad Ax = b.$$


12

For SVM primal problem

- objective is quadratic and convex, and all inequality constraints are affine
- Slater's condition holds trivially, hence, strong duality holds

SVM Dual Problem:

$$\begin{aligned} \text{maximize} \quad & \sum_{i=1}^m \lambda_i - \frac{1}{2} \sum_{i,j=1}^m \lambda_i \lambda_j y_i y_j (x_i \cdot x_j) \\ \text{subject to} \quad & \lambda_i \geq 0, \quad i = 1, \dots, m, \\ & \sum_{i=1}^m \lambda_i y_i = 0. \end{aligned}$$



SVM Dual Problem:

$$\begin{aligned} &\text{maximize} && \sum_{i=1}^m \lambda_i - \frac{1}{2} \sum_{i,j=1}^m \lambda_i \lambda_j y_i y_j (x_i \cdot x_j) \\ &\text{subject to} && \lambda_i \geq 0, \quad i = 1, \dots, m, \\ &&& \sum_{i=1}^m \lambda_i y_i = 0. \end{aligned}$$

Solving primal from dual:

- Assume that the dual problem is solved to obtain the dual optimal λ^*
- **Recall:** Setting the derivative of L w.r.t. w to **zero**, we got

SVM Dual Problem:

$$\begin{aligned} & \text{maximize} \quad \sum_{i=1}^m \lambda_i - \frac{1}{2} \sum_{i,j=1}^m \lambda_i \lambda_j y_i y_j (x_i \cdot x_j) \\ & \text{subject to} \quad \lambda_i \geq 0, \quad i = 1, \dots, m, \\ & \quad \quad \quad \sum_{i=1}^m \lambda_i y_i = 0. \end{aligned}$$

Solving primal from dual:

- Assume that the dual problem is solved to obtain the dual optimal λ^*
- **Recall:** Setting the derivative of L w.r.t. w to **zero**, we got

$$\frac{\partial L(w, b, \lambda)}{\partial w} = 0 \implies w = \sum_{i=1}^m \lambda_i y_i x_i$$

SVM Dual Problem:

$$\begin{aligned} & \text{maximize} && \sum_{i=1}^m \lambda_i - \frac{1}{2} \sum_{i,j=1}^m \lambda_i \lambda_j y_i y_j (x_i \cdot x_j) \\ & \text{subject to} && \lambda_i \geq 0, \quad i = 1, \dots, m, \\ & && \sum_{i=1}^m \lambda_i y_i = 0. \end{aligned}$$

Solving primal from dual:

- Assume that the dual problem is solved to obtain the dual optimal λ^*
- **Recall:** Setting the derivative of L w.r.t. w to **zero**, we got

$$\frac{\partial L(w, b, \lambda)}{\partial w} = 0 \implies w = \sum_{i=1}^m \lambda_i y_i x_i$$


- Hence we get primal optimal w^* as

$$w^* = \sum_{i=1}^m \lambda_i^* y_i x_i$$


- The primal intercept b^* is

$$b^* = - \frac{\max_{i:y^{(i)}=-1} w^{*T} x^{(i)} + \min_{i:y^{(i)}=+1} w^{*T} x^{(i)}}{2}$$

Computing the optimal intercept b^*




Computing the optimal intercept b^*




- For all x in red plane: $w^* \cdot x = -b$

Computing the optimal intercept b^*



- For all x in red plane: $w^* \cdot x = -b$
- Optimal b^* is such that two support vectors are equal distance


Computing the optimal intercept b^*



- We recall that b^* is such that $w^*x + b^* = 0$ is the middle plane

- For all x in red plane: $w^* \cdot x = -b$
- Optimal b^* is such that two support vectors are equal distance


Computing the optimal intercept b^*



- We recall that b^* is such that $w^*x + b^* = 0$ is the middle plane
- Assume that the normal w^* points in positive class direction

- For all x in red plane: $w^* \cdot x = -b$
- Optimal b^* is such that two support vectors are equal distance


Computing the optimal intercept b^*



- We recall that b^* is such that $w^* \cdot x + b^* = 0$ is the middle plane
- Assume that the normal w^* points in positive class direction
- Take projections of $x^{(i)}$ of positive class on w^* , and find the least farthest

- For all x in red plane: $w^* \cdot x = -b$
- Optimal b^* is such that two support vectors are equal distance

Computing the optimal intercept b^*




- We recall that b^* is such that $w^* \cdot x + b^* = 0$ is the middle plane
- Assume that the normal w^* points in positive class direction
- Take projections of $x^{(i)}$ of positive class on w^* , and find the least farthest

$$\min_{i: y^{(i)} = +1} w^{*T} x^{(i)}$$

- For all x in red plane: $w^* \cdot x = -b$
- Optimal b^* is such that two support vectors are equal distance


Computing the optimal intercept b^*



- For all x in red plane: $w^* \cdot x = -b$
- Optimal b^* is such that two support vectors are equal distance

- We recall that b^* is such that $w^* x + b^* = 0$ is the middle plane
- Assume that the normal w^* points in positive class direction
- Take projections of $x^{(i)}$ of positive class on w^* , and find the least farthest
$$\min_{i: y^{(i)} = +1} w^{*T} x^{(i)}$$
- Similarly, take projections of $x^{(i)}$ of negative class along $-w^*$, and find the farthest point

Computing the optimal intercept b^*




- For all x in red plane: $w^* \cdot x = -b$
- Optimal b^* is such that two support vectors are equal distance

- We recall that b^* is such that $w^* x + b^* = 0$ is the middle plane
- Assume that the normal w^* points in positive class direction
- Take projections of $x^{(i)}$ of positive class on w^* , and find the least farthest
$$\min_{i: y^{(i)} = +1} w^{*T} x^{(i)}$$
- Similarly, take projections of $x^{(i)}$ of negative class along $-w^*$, and find the farthest point

$$\min_{i: y^{(i)} = -1} -w^{*T} x^{(i)} = \max_{i: y^{(i)} = -1} w^{*T} x^{(i)}$$

Computing the optimal intercept b^*





- We recall that b^* is such that $w^* \cdot x + b^* = 0$ is the middle plane
- Assume that the normal w^* points in positive class direction
- Take projections of $x^{(i)}$ of positive class on w^* , and find the least farthest $\min_{i:y^{(i)}=+1} w^{*T} x^{(i)}$
- Similarly, take projections of $x^{(i)}$ of negative class along $-w^*$, and find the farthest point

- For all x in red plane: $w^* \cdot x = -b$
- Optimal b^* is such that two support vectors are equal distance


$$\min_{i:y^{(i)}=-1} -w^{*T} x^{(i)} = \max_{i:y^{(i)}=+1} w^{*T} x^{(i)}$$

Hence, optimal intercept $b^* = -\frac{\max_{i:y^{(i)}=-1} w^{*T} x^{(i)} + \min_{i:y^{(i)}=+1} w^{*T} x^{(i)}}{2}$

Linearly Separable with Noisy Data: Soft-Margin SVM *noisy?*




Linearly Separable with Noisy Data: Soft-Margin SVM



Two red and one blue samples noisy.


Linearly Separable with Noisy Data: Soft-Margin SVM



Two red and one blue samples noisy.

Goal: The data is noisy and it is **not linearly separable**. Modify SVM optimization model to allow for noisy data into account. Formulate optimization problem so that some **noisy data is allowed** between gap/margin (region between dotted lines).

Linearly Separable with Noisy Data: Soft-Margin SVM




- Want to allow for **noisy data**, i.e., allow some sample x_i to be between dotted planes

Two **red** and one **blue** samples noisy.

Goal: The data is noisy and it is **not linearly separable**. Modify SVM optimization model to allow for noisy data into account. Formulate optimization problem so that some **noisy data is allowed** between gap/margin (region between dotted lines).

Linearly Separable with Noisy Data: Soft-Margin SVM



$$y_i (w \cdot x_i + b) \geq 1$$


- Want to allow for noisy data, i.e., allow some sample x_i to be between dotted planes

$$y_i (w \cdot x_i + b) \geq 1 - \xi_i$$

Two **red** and one **blue** samples noisy.

Goal: The data is noisy and it is **not linearly separable**. Modify SVM optimization model to allow for noisy data into account. Formulate optimization problem so that some **noisy data is allowed** between gap/margin (region between dotted lines).

Linearly Separable with Noisy Data: Soft-Margin SVM



Two red and one blue samples noisy.


Goal: The data is noisy and it is **not linearly separable**. Modify SVM optimization model to allow for noisy data into account. Formulate optimization problem so that some **noisy data is allowed** between gap/margin (region between dotted lines).

- Want to allow for **noisy data**, i.e., allow some sample x_i to be between dotted planes

$$y_i(w \cdot x_i + b) \geq 1 - \xi_i$$

- Our margin is no longer strict or hard, it is now **softened**

Linearly Separable with Noisy Data: Soft-Margin SVM



Two **red** and one **blue** samples noisy.


Goal: The data is noisy and it is **not linearly separable**. Modify SVM optimization model to allow for noisy data into account. Formulate optimization problem so that some **noisy data is allowed** between gap/margin (region between dotted lines).

- Want to allow for **noisy data**, i.e., allow some sample x_i to be between dotted planes

$$y_i(w \cdot x_i + b) \geq 1 - \xi_i$$

- Our margin is no longer strict or hard, it is now **softened**
- Also, don't want too many x_i to be between dotted planes, introduce **penalty** in objective

Linearly Separable with Noisy Data: Soft-Margin SVM



Two **red** and one **blue** samples noisy.

Goal: The data is noisy and it is **not linearly separable**. Modify SVM optimization model to allow for noisy data into account. Formulate optimization problem so that some **noisy data is allowed** between gap/margin (region between dotted lines).

- Want to allow for **noisy data**, i.e., allow some sample x_i to be between dotted planes

$$y_i(w \cdot x_i + b) \geq 1 - \xi_i$$


- Our margin is no longer strict or hard, it is now **softened**
- Also, don't want too many x_i to be between dotted planes, introduce **penalty** in objective

min

$$\frac{1}{2} \|w\|^2 + C \sum_{i=1}^m \xi_i$$

(Handwritten orange annotations: a wavy line under the first term, an arrow pointing to the summation term, and a bracket under the summation term.)

Linearly Separable with Noisy Data: Soft-Margin SVM



Two **red** and one **blue** samples noisy.

Goal: The data is noisy and it is **not linearly separable**. Modify SVM optimization model to allow for noisy data into account. Formulate optimization problem so that some **noisy data is allowed** between gap/margin (region between dotted lines).

- Want to allow for **noisy data**, i.e., allow some sample x_i to be between dotted planes

$$y_i(w \cdot x_i + b) \geq 1 - \xi_i$$

- Our margin is no longer strict or hard, it is now **softened**
- Also, don't want too many x_i to be between dotted planes, introduce **penalty** in objective

$$\frac{1}{2} \|w\|^2 + C \sum_{i=1}^m \xi_i$$

- Since it is a **minimization**, large values of ξ_i will be **discouraged**

Primal and Dual Formulation of Soft-Margin SVM

Primal Soft-Margin SVM:

$$\text{minimize} \quad \frac{1}{2} \|w\|^2 + C \sum_{i=1}^m \xi_i$$


$$\text{subject to} \quad y_i(w \cdot x_i + b) \geq 1 - \xi_i, \quad i = 1, \dots, m.$$

Primal and Dual Formulation of Soft-Margin SVM

Primal Soft-Margin SVM:

$$\begin{aligned} &\text{minimize} \quad \frac{1}{2} \|w\|^2 + C \sum_{i=1}^m \xi_i \\ &\text{subject to} \quad y_i(w \cdot x_i + b) \geq 1 - \xi_i, \quad i = 1, \dots, m. \end{aligned}$$


Dual Soft-Margin SVM:

$$\begin{aligned} &\text{Maximize} \quad \sum_{i=1}^m \lambda_i - \frac{1}{2} \sum_{i,j=1}^m \lambda_i \lambda_j y_i y_j x_i \cdot x_j \\ &\text{subject to} \quad 0 \leq \lambda_i \leq C, \quad i = 1, \dots, m, \\ &\quad \quad \quad \sum_{i=1}^m \lambda_i y_i = 0. \end{aligned}$$



Quiz: How did we get the dual form? Try.

Effect of Parameter on soft-margin SVM

$$\text{Minimize } \frac{1}{2} \|w\|^2 + C \sum_{i=1}^m \xi_i \text{ subject to } y_i(w \cdot x_i + b) \geq 1 - \xi_i \text{ for } i = 1, \dots, m.$$



C=100



C=1



C=0.15



C=0.1

- For C very large, soft margin is equivalent to hard margin.
- When C is small, we allow misclassification.
- Here C is a hyperparameter.
- In practice, cross-validations can be used.

Primal and Dual Formulation of Soft-Margin SVM

Primal Soft-Margin SVM:

$$\begin{aligned} &\text{minimize} && \frac{1}{2} \|w\|^2 + C \sum_{i=1}^m \xi_i \\ &\text{subject to} && y_i(w \cdot x_i + b) \geq 1 - \xi_i, \\ &&& i = 1, \dots, m. \end{aligned}$$

Primal and Dual Formulation of Soft-Margin SVM

Primal Soft-Margin SVM:

$$\begin{aligned} &\text{minimize} && \frac{1}{2} \|w\|^2 + C \sum_{i=1}^m \xi_i \\ &\text{subject to} && y_i(w \cdot x_i + b) \geq 1 - \xi_i, \\ &&& i = 1, \dots, m. \end{aligned}$$

Additional advantages of primal over dual

- We see $x_i \cdot x_j$ in dual.

Dual Soft-Margin SVM:

$$\begin{aligned} &\text{Maximize} && \sum_{i=1}^m \lambda_i - \frac{1}{2} \sum_{i,j=1}^m \lambda_i \lambda_j y_i y_j x_i \cdot x_j \\ &\text{subject to} && 0 \leq \lambda_i \leq C, \quad i = 1, \dots, m, \\ &&& \sum_{i=1}^m \lambda_i y_i = 0. \end{aligned}$$

Primal and Dual Formulation of Soft-Margin SVM

Primal Soft-Margin SVM:

$$\begin{aligned} &\text{minimize} && \frac{1}{2} \|w\|^2 + C \sum_{i=1}^m \xi_i \\ &\text{subject to} && y_i(w \cdot x_i + b) \geq 1 - \xi_i, \\ &&& i = 1, \dots, m. \end{aligned}$$

Dual Soft-Margin SVM:

$$\begin{aligned} &\text{Maximize} && \sum_{i=1}^m \lambda_i - \frac{1}{2} \sum_{i,j=1}^m \lambda_i \lambda_j y_i y_j \underbrace{x_i \cdot x_j} \\ &\text{subject to} && 0 \leq \lambda_i \leq C, \quad i = 1, \dots, m, \\ &&& \sum_{i=1}^m \lambda_i y_i = 0. \end{aligned}$$

Additional advantages of ~~primal~~ ^{dual} over dual ~~dual~~ ^{primal}

- We see $x_i \cdot x_j$ in dual.
- Can we make use of this term?

Primal and Dual Formulation of Soft-Margin SVM

Primal Soft-Margin SVM:

$$\begin{aligned} &\text{minimize} && \frac{1}{2} \|w\|^2 + C \sum_{i=1}^m \xi_i \\ &\text{subject to} && y_i(w \cdot x_i + b) \geq 1 - \xi_i, \\ &&& i = 1, \dots, m. \end{aligned}$$

Dual Soft-Margin SVM:

$$\begin{aligned} &\text{Maximize} && \sum_{i=1}^m \lambda_i - \frac{1}{2} \sum_{i,j=1}^m \lambda_i \lambda_j y_i y_j x_i \cdot x_j \\ &\text{subject to} && 0 \leq \lambda_i \leq C, \quad i = 1, \dots, m, \\ &&& \sum_{i=1}^m \lambda_i y_i = 0. \end{aligned}$$

Additional advantages of primal over dual

- We see $x_i \cdot x_j$ in dual.
- Can we make use of this term?
- Sample x_i also appears in constraint in primal. Is that useful?

Primal and Dual Formulation of Soft-Margin SVM

Primal Soft-Margin SVM:

$$\begin{aligned} &\text{minimize} \quad \frac{1}{2} \|w\|^2 + C \sum_{i=1}^m \xi_i \\ &\text{subject to} \quad y_i(w \cdot x_i + b) \geq 1 - \xi_i, \\ &\quad \quad \quad i = 1, \dots, m. \end{aligned}$$

Dual Soft-Margin SVM:

$$\begin{aligned} &\text{Maximize} \quad \sum_{i=1}^m \lambda_i - \frac{1}{2} \sum_{i,j=1}^m \lambda_i \lambda_j y_i y_j x_i \cdot x_j \\ &\text{subject to} \quad 0 \leq \lambda_i \leq C, \quad i = 1, \dots, m, \\ &\quad \quad \quad \sum_{i=1}^m \lambda_i y_i = 0. \end{aligned}$$

Additional advantages of primal over dual

- We see $x_i \cdot x_j$ in dual.
- Can we make use of this term?
- Sample x_i also appears in constraint in primal. Is that useful?
- Fact that $x_i \cdot x_j$ appears only in dual objective and that too as dot product is useful.

Primal and Dual Formulation of Soft-Margin SVM

Primal Soft-Margin SVM:

$$\begin{aligned} &\text{minimize} && \frac{1}{2} \|w\|^2 + C \sum_{i=1}^m \xi_i \\ &\text{subject to} && y_i(w \cdot x_i + b) \geq 1 - \xi_i, \\ &&& i = 1, \dots, m. \end{aligned}$$


Dual Soft-Margin SVM:

$$\begin{aligned} &\text{Maximize} && \sum_{i=1}^m \lambda_i - \frac{1}{2} \sum_{i,j=1}^m \lambda_i \lambda_j y_i y_j x_i \cdot x_j \\ &\text{subject to} && 0 \leq \lambda_i \leq C, \quad i = 1, \dots, m, \\ &&& \sum_{i=1}^m \lambda_i y_i = 0. \end{aligned}$$


Additional advantages of primal over dual

- We see $x_i \cdot x_j$ in dual.
- Can we make use of this term?
- Sample x_i also appears in constraint in primal. Is that useful?
- Fact that $x_i \cdot x_j$ appears only in dual objective and that too as dot product is useful.
- This allows mapping samples to a space where they may be linearly separated.

Non-Linearly Separable Data, Kernel Trick




Non-Linearly Separable Data, Kernel Trick



Here data is **not linearly separable** in the input space


Non-Linearly Separable Data, Kernel Trick



Here data is **not linearly separable** in the input space

After mapping to a new feature space, **data is now linearly separable!**

Non-Linearly Separable Data, Kernel Trick



Here data is **not linearly separable** in the input space

After mapping to a new feature space, **data is now linearly separable!**

Quiz: How to find such map $\Phi: \mathbb{R}^N \rightarrow H$

Kernel Trick

For data in input space

$$f(x) = \text{sign}(w \cdot x + b)$$

$$w = \sum_{i=1}^m \alpha_i y_i x_i$$

Kernel Trick



For data in input space

$$f(x) = \text{sign}(w \cdot x + b)$$

$$w = \sum_{i=1}^m \alpha_i y_i x_i$$

Data in feature space $\Phi(x)$

$$f(x) = \text{sign}(w \cdot \Phi(x) + b)$$

$$w = \sum_{i=1}^m \alpha_i y_i \Phi(x_i)$$

- Solve the dual problem for α_i

Kernel Trick

For data in input space

$$f(x) = \text{sign}(w \cdot x + b)$$

$$w = \sum_{i=1}^m \alpha_i y_i x_i$$

Data in feature space $\Phi(x)$

$$f(x) = \text{sign}(w \cdot \Phi(x) + b)$$

$$w = \sum_{i=1}^m \alpha_i y_i \Phi(x_i)$$

- Solve the dual problem for α_i
- Compute w, b from above

Kernel Trick

For data in input space

$$f(x) = \text{sign}(w \cdot x + b)$$

$$w = \sum_{i=1}^m \alpha_i y_i x_i$$

Data in feature space $\Phi(x)$

$$f(x) = \text{sign}(w \cdot \Phi(x) + b)$$

$$w = \sum_{i=1}^m \alpha_i y_i \Phi(x_i)$$

- Solve the dual problem for α_i
- Compute w, b from above
- Predict using $f(x)$ above

Kernel Trick

For data in input space

$$f(x) = \text{sign}(w \cdot x + b)$$

$$w = \sum_{i=1}^m \alpha_i y_i x_i$$

- Solve the dual problem for α_i
- Compute w, b from above
- Predict using $f(x)$ above

Data in feature space $\Phi(x)$

$$f(x) = \text{sign}(w \cdot \Phi(x) + b)$$

$$w = \sum_{i=1}^m \alpha_i y_i \Phi(x_i)$$

- Solve for α_i using dual

Kernel Trick

For data in input space

$$f(x) = \text{sign}(w \cdot x + b)$$

$$w = \sum_{i=1}^m \alpha_i y_i x_i$$

- Solve the dual problem for α_i
- Compute w, b from above
- Predict using $f(x)$ above

Data in feature space $\Phi(x)$

$$f(x) = \text{sign}(w \cdot \Phi(x) + b)$$

$$w = \sum_{i=1}^m \alpha_i y_i \Phi(x_i)$$

- Solve for α_i using dual
- Compute w, b (x_i by $\Phi(x_i)$)

Kernel Trick

For data in input space

$$f(x) = \text{sign}(w \cdot x + b)$$

$$w = \sum_{i=1}^m \alpha_i y_i x_i$$

- Solve the dual problem for α_i
- Compute w, b from above
- Predict using $f(x)$ above

Data in feature space $\Phi(x)$

$$f(x) = \text{sign}(w \cdot \Phi(x) + b)$$

$$w = \sum_{i=1}^m \alpha_i y_i \Phi(x_i)$$

- Solve for α_i using dual
- Compute w, b (x_i by $\Phi(x_i)$)
- Predict with $f(x)$. **So what?** Trick: substitute w in $f(x)$ above!

Kernel Trick:

$$f(x) = \text{sign} \left(\sum_{i=1}^m \alpha_i y_i \Phi(x_i) \cdot \Phi(x) + b \right) = \text{sign} \left(\sum_{i=1}^m \alpha_i y_i K(x_i, x) + b \right)$$

Kernel Trick



For data in input space

$$f(x) = \text{sign}(w \cdot x + b)$$

$$w = \sum_{i=1}^m \alpha_i y_i x_i$$

- Solve the dual problem for α_i
- Compute w, b from above
- Predict using $f(x)$ above

Data in feature space $\Phi(x)$

$$f(x) = \text{sign}(w \cdot \Phi(x) + b)$$

$$w = \sum_{i=1}^m \alpha_i y_i \Phi(x_i)$$


- Solve for α_i using dual
- Compute w, b (x_i by $\Phi(x_i)$)
- Predict with $f(x)$. **So what?** Trick: substitute w in $f(x)$ above!

Kernel Trick:

$$f(x) = \text{sign} \left(\sum_{i=1}^m \alpha_i y_i \Phi(x_i) \cdot \Phi(x) + b \right) = \text{sign} \left(\sum_{i=1}^m \alpha_i y_i K(x_i, x) + b \right)$$

Dont need to know Φ ; Only a function $K(\cdot, \cdot) : \mathbb{R}^m \times \mathbb{R}^m \rightarrow \mathbb{R}$ will do!

Kernel Trick



For data in input space

$$f(x) = \text{sign}(w \cdot x + b)$$

$$w = \sum_{i=1}^m \alpha_i y_i x_i$$

- Solve the dual problem for α_i
- Compute w, b from above
- Predict using $f(x)$ above

Data in feature space $\Phi(x)$

$$f(x) = \text{sign}(w \cdot \Phi(x) + b)$$

$$w = \sum_{i=1}^m \alpha_i y_i \Phi(x_i)$$

- Solve for α_i using dual
- Compute w, b (x_i by $\Phi(x_i)$)
- Predict with $f(x)$. **So what?** Trick: substitute w in $f(x)$ above!

Kernel Trick:

$$f(x) = \text{sign} \left(\sum_{i=1}^m \alpha_i y_i \underbrace{\Phi(x_i) \cdot \Phi(x)}_{K(x_i, x)} + b \right) = \text{sign} \left(\sum_{i=1}^m \alpha_i y_i \underbrace{K(x_i, x)}_{\text{kernel}} + b \right)$$

Dont need to know Φ ; Only a function $K(\cdot, \cdot) : \mathbb{R}^m \times \mathbb{R}^m \rightarrow \mathbb{R}$ will do!

→ Avoid explicit computation of features or infinite length vectors

Popular Kernels

Kernel: A kernel is a **dot product** in some feature space

$$K(x_i, x_j) = \Phi(x_i) \cdot \Phi(x_j)$$

Popular Kernels

Kernel: A kernel is a **dot product** in some feature space

$$K(x_i, x_j) = \Phi(x_i) \cdot \Phi(x_j)$$

$$-\gamma (x_i - x_j)^T (x_i - x_j)$$

- ✓ $K(x_i, x_j) = \underline{x_i} \cdot \underline{x_j}$
- ✓ $K(x_i, x_j) = \exp(-\gamma \|\underline{x_i} - \underline{x_j}\|^2)$
- ✓ $K(x_i, x_j) = \exp(-\gamma \|x_i - x_j\|)$
- ✓ $K(x_i, x_j) = (p + \underline{x_i} \cdot \underline{x_j})^q$
- ✓ $K(\underline{x_j}, \underline{x_i}) = (p + \underline{x_i} \cdot \underline{x_j})^q \exp(-\gamma \|\underline{x_i} - \underline{x_j}\|^2)$
- $K(x_i, x_j) = \tanh(kx_i \cdot x_j - \delta)$

• Linear kernel ←

• Gaussian kernel ←

• Exponential kernel ←


• Polynomial kernel ←

• Hybrid kernel ←


• Sigmoidal ←

Polynomial Kernel

Polynomial Kernel




Polynomial Kernel




- Data is **not** linearly separable

Polynomial Kernel




- Data is **not** linearly separable
- Apply Kernel $K(x, z) = (x \cdot z)^2$ to map data to higher dimension, to make it linearly separable

Polynomial Kernel



- Data is **not** linearly separable
- Apply Kernel $K(x, z) = (x \cdot z)^2$ to map data to higher dimension, to make it linearly separable
- $x_1 = [x_{(1)}, x_{(2)}]^T$, $z_1 = [z_{(1)}, z_{(2)}]^T$

Polynomial Kernel



- Data is **not** linearly separable
- Apply Kernel $K(x, z) = (x \cdot z)^2$ to map data to higher dimension, to make it linearly separable
- $x_1 = [x_{(1)}, x_{(2)}]^T$, $z_1 = [z_{(1)}, z_{(2)}]^T$

We have

$$K(x \cdot z) = (x \cdot z)^2 = \left(\begin{bmatrix} x_{(1)} \\ x_{(2)} \end{bmatrix} \cdot \begin{bmatrix} z_{(1)} \\ z_{(2)} \end{bmatrix} \right)^2 = (x_{(1)}z_{(1)} + x_{(2)}z_{(2)})^2$$


$$= x_{(1)}^2 z_{(1)}^2 + 2x_{(1)}z_{(1)}x_{(2)}z_{(2)} + x_{(2)}^2 z_{(2)}^2 = \begin{bmatrix} x_{(1)}^2 \\ \sqrt{2}x_{(1)}x_{(2)} \\ x_{(2)}^2 \end{bmatrix} \cdot \begin{bmatrix} z_{(1)}^2 \\ \sqrt{2}z_{(1)}z_{(2)} \\ z_{(2)}^2 \end{bmatrix} = \Phi(x) \cdot \Phi(z)$$

much cheaper than


Cost = 4

Cost = 11

Separation Happens After Using Polynomial Kernel!




Separation Happens After Using Polynomial Kernel!



- In last slide: $\Phi(x) = \begin{bmatrix} x_{(1)}^2 \\ \sqrt{2}x_{(1)}x_{(2)} \\ x_{(2)}^2 \end{bmatrix}$

Separation Happens After Using Polynomial Kernel!




- In last slide: $\Phi(x) = \begin{bmatrix} x_{(1)}^2 \\ \sqrt{2}x_{(1)}x_{(2)} \\ x_{(2)}^2 \end{bmatrix}$
- The points in figure above are

$$x_1 = (0, 1), \quad x_2 = (0, -1)$$

$$x_3 = (1, 0), \quad x_4 = (-1, 0)$$

Separation Happens After Using Polynomial Kernel!



- In last slide: $\Phi(x) = \begin{bmatrix} x_{(1)}^2 \\ \sqrt{2}x_{(1)}x_{(2)} \\ x_{(2)}^2 \end{bmatrix}$

- The points in figure above are

$$x_1 = (0, 1), \quad x_2 = (0, -1)$$

$$x_3 = (1, 0), \quad x_4 = (-1, 0)$$

- We have


$$\Phi(x_1) = [0, 0, 1]^T$$

$$\Phi(x_2) = [0, 0, 1]^T$$

$$\Phi(x_3) = [1, 0, 0]^T$$

$$\Phi(x_4) = [1, 0, 0]^T$$

Separation Happens After Using Polynomial Kernel!



- In last slide: $\Phi(x) = \begin{bmatrix} x_{(1)}^2 \\ \sqrt{2}x_{(1)}x_{(2)} \\ x_{(2)}^2 \end{bmatrix}$

- The points in figure above are

$$x_1 = (0, 1), \quad x_2 = (0, -1)$$

$$x_3 = (1, 0), \quad x_4 = (-1, 0)$$

- We have

$$\Phi(x_1) = [0, 0, 1]^T$$

$$\Phi(x_2) = [0, 0, 1]^T$$

$$\Phi(x_3) = [1, 0, 0]^T$$

$$\Phi(x_4) = [1, 0, 0]^T$$

- After Φ transformation, points are **now linearly separable!**

Same Kernel Function but Different Features

Note: Feature space might not be unique for the same kernel function.

Same Kernel Function but Different Features

Note: Feature space might not be unique for the same kernel function.

Example-1: Consider $\Phi : \mathbb{R}^2 \rightarrow \mathbb{R}^3$

$$\begin{aligned}\Phi(x_1, x_2) &= (x_1^2, x_2^2, \sqrt{2}x_1x_2) \\ \Phi(x) \cdot \Phi(z) &= (x_1^2, x_2^2, \sqrt{2}x_1x_2) \cdot (z_1^2, z_2^2, \sqrt{2}z_1z_2) \\ &= (x_1z_1 + x_2z_2)^2 = (x \cdot z)^2 = \underbrace{K(x, z)}.\end{aligned}$$

Same Kernel Function but Different Features

Note: Feature space might not be unique for the same kernel function.


Example-1: Consider $\Phi : \mathbb{R}^2 \rightarrow \mathbb{R}^3$

$$\begin{aligned}\Phi(x_1, x_2) &= (x_1^2, x_2^2, \sqrt{2}x_1x_2) \\ \Phi(x) \cdot \Phi(z) &= (x_1^2, x_2^2, \sqrt{2}x_1x_2) \cdot (z_1^2, z_2^2, \sqrt{2}z_1z_2) \\ &= (x_1z_1 + x_2z_2)^2 = (x \cdot z)^2 = K(x, z).\end{aligned}$$



Example-2: Consider $\Phi : \mathbb{R}^2 \rightarrow \mathbb{R}^4$

$$\begin{aligned}\Phi(x_1, x_2) &= (x_1^2, x_2^2, x_1x_2, x_2x_1) \\ \Phi(x) \cdot \Phi(z) &= (x_1^2, x_2^2, x_1x_2, x_2x_1) \cdot (z_1^2, z_2^2, z_1z_2, z_2z_1) \\ &= (x \cdot z)^2 = \underline{K(x, z)}.\end{aligned}$$




RBF or Gaussian Kernel






RBF or Gaussian Kernel



RBF or Gaussian Kernel






RBF or Gaussian Kernel



After applying the kernel, we observe:

RBF or Gaussian Kernel



After applying the kernel, we observe:

- **red** data points are bumped up
- **yellow** data points are pushed to cavity

Model Selection: Which Kernels with which parameter?

Parameter C	Polynomial degree <i>d</i>				
	(0.1, 1)	(1, 1)	(10, 1)	(100, 1)	(1000, 1)
	(0.1, 2)	(1, 2)	(10, 2)	(100, 2)	(1000, 2)
	(0.1, 3)	(1, 3)	(10, 3)	(100, 3)	(1000, 3)
	(0.1, 4)	(1, 4)	(10, 4)	(100, 4)	(1000, 4)
	(0.1, 5)	(1, 5)	(10, 5)	(100, 5)	(1000, 5)

Model Selection: Which Kernels with which parameter?

Parameter C	Polynomial degree d				
	(0.1, 1)	(1, 1)	(10, 1)	(100, 1)	(1000, 1)
	(0.1, 2)	(1, 2)	(10, 2)	(100, 2)	(1000, 2)
	(0.1, 3)	(1, 3)	(10, 3)	(100, 3)	(1000, 3)
	(0.1, 4)	(1, 4)	(10, 4)	(100, 4)	(1000, 4)
	(0.1, 5)	(1, 5)	(10, 5)	(100, 5)	(1000, 5)

- Consider polynomial kernel.

Model Selection: Which Kernels with which parameter?

Parameter C	Polynomial degree d				
	(0.1, 1)	(1, 1)	(10, 1)	(100, 1)	(1000, 1)
	(0.1, 2)	(1, 2)	(10, 2)	(100, 2)	(1000, 2)
	(0.1, 3)	(1, 3)	(10, 3)	(100, 3)	(1000, 3)
	(0.1, 4)	(1, 4)	(10, 4)	(100, 4)	(1000, 4)
	(0.1, 5)	(1, 5)	(10, 5)	(100, 5)	(1000, 5)

- Consider polynomial kernel.
- There are parameters: C, d .

Model Selection: Which Kernels with which parameter?

Parameter C	Polynomial degree d				
	(0.1, 1)	(1, 1)	(10, 1)	(100, 1)	(1000, 1)
	(0.1, 2)	(1, 2)	(10, 2)	(100, 2)	(1000, 2)
	(0.1, 3)	(1, 3)	(10, 3)	(100, 3)	(1000, 3)
	(0.1, 4)	(1, 4)	(10, 4)	(100, 4)	(1000, 4)
	(0.1, 5)	(1, 5)	(10, 5)	(100, 5)	(1000, 5)

- Consider polynomial kernel.
- There are parameters: C, d .
- We can consider possible values.

Model Selection: Which Kernels with which parameter?

Parameter C	Polynomial degree d				
	(0.1, 1)	(1, 1)	(10, 1)	(100, 1)	(1000, 1)
	(0.1, 2)	(1, 2)	(10, 2)	(100, 2)	(1000, 2)
	(0.1, 3)	(1, 3)	(10, 3)	(100, 3)	(1000, 3)
	(0.1, 4)	(1, 4)	(10, 4)	(100, 4)	(1000, 4)
	(0.1, 5)	(1, 5)	(10, 5)	(100, 5)	(1000, 5)


- Consider polynomial kernel.
- There are parameters: C, d .
- We can consider possible values.
- Check our choice by doing cross-validations.

Model Selection: Which Kernels with which parameter?

Parameter C	Polynomial degree d				
	(0.1, 1)	(1, 1)	(10, 1)	(100, 1)	(1000, 1)
	(0.1, 2)	(1, 2)	(10, 2)	(100, 2)	(1000, 2)
	(0.1, 3)	(1, 3)	(10, 3)	(100, 3)	(1000, 3)
	(0.1, 4)	(1, 4)	(10, 4)	(100, 4)	(1000, 4)
	(0.1, 5)	(1, 5)	(10, 5)	(100, 5)	(1000, 5)

- Consider polynomial kernel.
- There are parameters: C, d .
- We can consider possible values.
- Check our choice by doing cross-validations.

Recall the main idea of cross-validation:



SVM in Unconstrained Form: loss + penalty form

Primal Soft-Margin SVM:

$$\text{minimize} \quad \frac{1}{2} \|w\|^2 + C \sum_{i=1}^m \xi_i$$

$$\text{subject to} \quad \left. \begin{aligned} y_i(w \cdot x_i + b) &\geq 1 - \xi_i, \\ i &= 1, \dots, m. \end{aligned} \right\}$$

SVM in Unconstrained Form: loss + penalty form

Primal Soft-Margin SVM:

$$\text{minimize} \quad \frac{1}{2} \|w\|^2 + C \sum_{i=1}^m \xi_i$$

$$\text{subject to} \quad y_i(w \cdot x_i + b) \geq 1 - \xi_i, \\ i = 1, \dots, m.$$

1. Write constraint as

$$y_i(w \cdot x_i + b) \geq 1 - \xi_i$$

$$\xi_i \geq \underbrace{-(w \cdot x_i + b)y_i + 1}$$

SVM in Unconstrained Form: loss + penalty form

Primal Soft-Margin SVM:

$$\begin{aligned} &\text{minimize} && \frac{1}{2} \|w\|^2 + C \sum_{i=1}^m \xi_i \\ &\text{subject to} && y_i(w \cdot x_i + b) \geq 1 - \xi_i, \\ &&& i = 1, \dots, m. \end{aligned}$$

2. Write constraints on ξ as function?

$$\xi_i = \max(0, 1 - y_i f(x_i))$$

Handwritten note: max is not diff.

1. Write constraint as

$$y_i(w \cdot x_i + b) \geq 1 - \xi_i$$

$$\xi_i \geq -(\underbrace{w \cdot x_i + b}_{f(x_i)}) y_i + 1$$

and $\xi_i \geq 0$.

SVM in Unconstrained Form: loss + penalty form

Primal Soft-Margin SVM:

$$\begin{aligned} & \text{minimize} && \frac{1}{2} \|w\|^2 + C \sum_{i=1}^m \xi_i \\ & \text{subject to} && y_i(w \cdot x_i + b) \geq 1 - \xi_i, \\ & && i = 1, \dots, m. \end{aligned}$$

1. Write constraint as

$$\begin{aligned} y_i(w \cdot x_i + b) &\geq 1 - \xi_i \\ \xi_i &\geq -(w \cdot x_i + b)y_i + 1 \end{aligned}$$

and $\xi_i \geq 0$.

2. Write constraints on ξ as function?

$$\xi_i = \max(0, 1 - y_i f(x_i))$$

3. Substituting ξ_i in objective

$$\begin{aligned} &= \frac{1}{2} \|w\|^2 + C \sum_{i=1}^m \max(0, 1 - y_i f(x_i)) \\ &= \frac{1}{2C} \|w\|^2 + \sum_{i=1}^m \max(0, 1 - y_i f(x_i)) \\ &= \lambda \|w\|^2 + \sum_{i=1}^m \max(0, 1 - y_i f(x_i)) \\ &= \lambda \|w\|^2 + \sum_{i=1}^m [0, 1 - y_i f(x_i)]_+ \end{aligned}$$

Unconstrained SVM: Find w, b s.t. minimize $\sum_{i=1}^m [1 - y_i f(x_i)]_+ + \lambda \|w\|_2^2$

SVM in Unconstrained Form: loss + penalty form

Primal Soft-Margin SVM:

$$\begin{aligned} &\text{minimize} && \frac{1}{2} \|w\|^2 + C \sum_{i=1}^m \xi_i \\ &\text{subject to} && y_i(w \cdot x_i + b) \geq 1 - \xi_i, \\ &&& i = 1, \dots, m. \end{aligned}$$

1. Write constraint as

$$\begin{aligned} y_i(w \cdot x_i + b) &\geq 1 - \xi_i \\ \xi_i &\geq -(w \cdot x_i + b)y_i + 1 \end{aligned}$$

and $\xi_i \geq 0$.

2. Write constraints on ξ as function?

$$\xi_i = \max(0, 1 - y_i f(x_i))$$

3. Substituting ξ_i in objective

$$\begin{aligned} &= \frac{1}{2} \|w\|^2 + C \sum_{i=1}^m \max(0, 1 - y_i f(x_i)) \\ &= \frac{1}{2C} \|w\|^2 + \sum_{i=1}^m \max(0, 1 - y_i f(x_i)) \\ &= \lambda \|w\|^2 + \sum_{i=1}^m \max(0, 1 - y_i f(x_i)) \\ &= \lambda \|w\|^2 + \sum_{i=1}^m [0, 1 - y_i f(x_i)]_+ \end{aligned}$$

Unconstrained SVM: Find w, b s.t. minimize $\sum_{i=1}^m [1 - y_i f(x_i)]_+ + \lambda \|w\|_2^2$

Penalty: $\lambda \|w\|_2^2$.

Loss (Hinge Loss): $\sum_{i=1}^m [1 - y_i f(x_i)]_+$.



Variety of Loss+Penalty Formulations for SVM


Loss	Penalty function	Resulting algorithm
Hinge Loss: $\sum_{i=1}^m [1 - y_i f(x_i)]_+$	$\lambda \ w\ _2^2$	SVM
Mean Squared Error $\sum_{i=1}^m (y_i - f(x_i))^2$	$\lambda \ w\ _2^2$	Ridge Regression
Mean Squared Error $\sum_{i=1}^m (y_i - f(x_i))^2$	$\lambda \ w\ _1$	Lasso
Mean Squared Error $\sum_{i=1}^m (y_i - f(x_i))^2$	$\lambda_1 \ w\ _1 + \lambda_2 \ w\ _2^2$	Elastic Net
Hinge Loss: $\sum_{i=1}^m [1 - y_i f(x_i)]_+$	$\lambda \ w\ _1$	1-Norm SVM

Variety of Loss+Penalty Formulations for SVM


Loss	Penalty function	Resulting algorithm
Hinge Loss: $\sum_{i=1}^m [1 - y_i f(x_i)]_+$	$\lambda \ w\ ^2$	SVM
Mean Squared Error $\sum_{i=1}^m (y_i - f(x_i))^2$	$\lambda \ w\ _2^2$	Ridge Regression
Mean Squared Error $\sum_{i=1}^m (y_i - f(x_i))^2$	$\lambda \ w\ _1$	Lasso
Mean Squared Error $\sum_{i=1}^m (y_i - f(x_i))^2$	$\lambda_1 \ w\ _1 + \lambda_2 \ w\ _2^2$	Elastic Net
Hinge Loss: $\sum_{i=1}^m [1 - y_i f(x_i)]_+$	$\lambda \ w\ _1$	1-Norm SVM

Algorithm	Loss	Penalty
SVM	convex, non-differentiable	convex, differentiable
Ridge Regression	convex, differentiable	convex, differentiable
Lasso	convex, differentiable	convex, non-differentiable
Elastic Net	convex, differentiable	convex, non-differentiable
Hinge Loss	convex, non-differentiable	convex, non-differentiable

Hard Margin SVM Regression: ϵ -SVR




Hard Margin SVM Regression: ϵ -SVR



Goal: Find a linear function that fits the red data points.

Hard Margin SVM Regression: ϵ -SVR




Goal: Find a **linear function** that fits the **red** data points.

Optimization Model:

$$\begin{aligned} & \text{minimize} && \frac{1}{2} \|w\|^2 \\ & \text{subject to:} && y_i(w \cdot x_i + b) \leq \epsilon, \\ & && y_i(w \cdot x_i + b) \geq -\epsilon, \\ & && i = 1, \dots, m. \end{aligned}$$

Hard Margin SVM Regression: ϵ -SVR



Goal: Find a **linear function** that fits the **red** data points.

Optimization Model:

$$\begin{aligned} &\text{minimize} && \frac{1}{2} \|w\|^2 \\ &\text{subject to:} && y_i(w \cdot x_i + b) \leq \epsilon, \\ &&& y_i(w \cdot x_i + b) \geq -\epsilon, \\ &&& i = 1, \dots, m. \end{aligned}$$

Remark: Hence, the model suggests that the difference between y_i and fitted function should be smaller than ϵ (hyperparameter) and larger than $-\epsilon$. That is

$$|y_i - (w \cdot x_i + b)| \leq \epsilon.$$

Here y_i is the **height** of the samples x_i , and **not** +1 or -1 labels!

Formulate Dual Margin Kernel SVM Problem for QP

Dual Soft-Margin Kernel SVM:

$$\text{Maximize} \quad \sum_{i=1}^m \lambda_i - \frac{1}{2} \sum_{i,j=1}^m \lambda_i \lambda_j y_i y_j \Phi(x_i) \cdot \Phi(x_j)$$

$$\text{subject to} \quad 0 \leq \lambda_i \leq C, \quad i = 1, \dots, m,$$

$$\sum_{i=1}^m \lambda_i y_i = 0.$$

Formulate Dual Margin Kernel SVM Problem for QP

Dual Soft-Margin Kernel SVM:

$$\begin{aligned} &\text{Maximize} && \sum_{i=1}^m \lambda_i - \frac{1}{2} \sum_{i,j=1}^m \lambda_i \lambda_j y_i y_j \underbrace{\Phi(x_i) \cdot \Phi(x_j)} \\ &\text{subject to} && \underbrace{0 \leq \lambda_i \leq C, \quad i = 1, \dots, m,} \\ &&& \underbrace{\sum_{i=1}^m \lambda_i y_i = 0.} \end{aligned}$$

Recall QP:

$$\begin{aligned} &\text{minimize} && \frac{1}{2} \underbrace{x^T P x + q^T x,} \\ &\text{subject to} && \underbrace{Gx \leq h,} \\ &&& \underbrace{Ax = b} \end{aligned}$$

How to set: x, P, q, G, h, A, b ?

Formulate Dual Margin Kernel SVM Problem for QP

Dual Soft-Margin Kernel SVM:

$$\begin{aligned} &\text{Maximize} && \sum_{i=1}^m \lambda_i - \frac{1}{2} \sum_{i,j=1}^m \lambda_i \lambda_j y_i y_j \Phi(x_i) \cdot \Phi(x_j) \\ &\text{subject to} && 0 \leq \lambda_i \leq C, \quad i = 1, \dots, m, \\ &&& \sum_{i=1}^m \lambda_i y_i = 0. \end{aligned}$$

Recall QP:

$$\begin{aligned} &\text{minimize} && \frac{1}{2} \underline{x}^T P x + q^T x, \\ &\text{subject to} && Gx \leq h, \\ &&& Ax = b \end{aligned}$$

How to set: x, P, q, G, h, A, b ?

1. Set $\underline{x} = [\lambda_1, \lambda_2, \dots, \lambda_m]^T$

Formulate Dual Margin Kernel SVM Problem for QP

Dual Soft-Margin Kernel SVM:

$$\begin{aligned} &\text{Maximize} && \sum_{i=1}^m \lambda_i - \frac{1}{2} \sum_{i,j=1}^m \lambda_i \lambda_j y_i y_j \underbrace{\Phi(x_i) \cdot \Phi(x_j)} \\ &\text{subject to} && 0 \leq \lambda_i \leq C, \quad i = 1, \dots, m, \\ &&& \sum_{i=1}^m \lambda_i y_i = 0. \end{aligned}$$

Recall QP:

$$\begin{aligned} &\text{minimize} && \frac{1}{2} x^T P x + q^T x, \\ &\text{subject to} && Gx \leq h, \\ &&& Ax = b \end{aligned}$$

How to set: x, P, q, G, h, A, b ?

1. Set $x = [\lambda_1, \lambda_2, \dots, \lambda_m]^T$
2. Let $\underbrace{K(x_i, x_j)} = \underbrace{\Phi(x_i) \cdot \Phi(x_j)}$

Formulate Dual Margin Kernel SVM Problem for QP

Dual Soft-Margin Kernel SVM:

$$\begin{aligned} &\text{Maximize} && \sum_{i=1}^m \lambda_i - \frac{1}{2} \sum_{i,j=1}^m \lambda_i \lambda_j \underbrace{y_i y_j}_{\text{K}} \underbrace{\Phi(x_i) \cdot \Phi(x_j)}_{\text{K}} \\ &\text{subject to} && 0 \leq \lambda_i \leq C, \quad i = 1, \dots, m, \\ &&& \sum_{i=1}^m \lambda_i y_i = 0. \end{aligned}$$

Recall QP:

$$\begin{aligned} &\text{minimize} && \frac{1}{2} x^T P x + q^T x, \\ &\text{subject to} && Gx \leq h, \\ &&& Ax = b \end{aligned}$$

How to set: x, P, q, G, h, A, b ?

1. Set $x = [\lambda_1, \lambda_2, \dots, \lambda_m]^T$
2. Let $K(x_i, x_j) = \Phi(x_i) \cdot \Phi(x_j)$
3. Set matrix $\underbrace{P}_{ij} = \underbrace{y_i y_j}_{\text{K}} \underbrace{K(x_i, x_j)}_{\text{K}}$

$$[\lambda_1 \dots \lambda_m]^T \leftarrow P \begin{bmatrix} 1 \\ \vdots \\ 1 \end{bmatrix}$$

Formulate Dual Margin Kernel SVM Problem for QP

Dual Soft-Margin Kernel SVM:

$$\begin{aligned} &\text{Maximize} \quad \sum_{i=1}^m \lambda_i - \frac{1}{2} \sum_{i,j=1}^m \lambda_i \lambda_j y_i y_j \Phi(x_i) \cdot \Phi(x_j) \\ &\text{subject to} \quad 0 \leq \lambda_i \leq C, \quad i = 1, \dots, m, \\ &\quad \quad \quad \sum_{i=1}^m \lambda_i y_i = 0. \end{aligned}$$

Handwritten notes:

$$\min \quad -\sum \lambda_i + \frac{1}{2} \sum \lambda_i \lambda_j y_i y_j \Phi(x_i) \cdot \Phi(x_j)$$

(An arrow points from the handwritten expression to the objective function in the Dual Soft-Margin Kernel SVM formulation.)

Recall QP:

$$\begin{aligned} &\text{minimize} \quad \frac{1}{2} x^T P x + \underline{q}^T x, \\ &\text{subject to} \quad Gx \leq h, \\ &\quad \quad \quad Ax = b \end{aligned}$$

How to set: x, P, q, G, h, A, b ?

1. Set $x = [\lambda_1, \lambda_2, \dots, \lambda_m]^T$
2. Let $K(x_i, x_j) = \Phi(x_i) \cdot \Phi(x_j)$
3. Set matrix $P_{ij} = y_i y_j K(x_i, x_j)$
4. q is column vector containing -1.

Formulate Dual Margin Kernel SVM Problem for QP

Dual Soft-Margin Kernel SVM:

$$\text{Maximize} \quad \sum_{i=1}^m \lambda_i - \frac{1}{2} \sum_{i,j=1}^m \lambda_i \lambda_j y_i y_j \Phi(x_i) \cdot \Phi(x_j)$$

$$\text{subject to} \quad 0 \leq \lambda_i \leq C, \quad i = 1, \dots, m,$$

$$\sum_{i=1}^m \lambda_i y_i = 0.$$

Recall QP:

$$\text{minimize} \quad \frac{1}{2} x^T P x + q^T x,$$

$$\text{subject to} \quad Gx \leq h,$$

$$\underline{Ax = b}$$

How to set: x, P, q, G, h, A, b ?

1. Set $x = [\lambda_1, \lambda_2, \dots, \lambda_m]^T$
2. Let $K(x_i, x_j) = \Phi(x_i) \cdot \Phi(x_j)$
3. Set matrix $P_{ij} = y_i y_j K(x_i, x_j)$
4. q is column vector containing -1.
5. For equality constraint, set

$$A = [y_1, y_2, \dots, y_m]^T, \quad b = 0$$

Formulate Dual Margin Kernel SVM Problem for QP

Dual Soft-Margin Kernel SVM:

$$\text{Maximize} \quad \sum_{i=1}^m \lambda_i - \frac{1}{2} \sum_{i,j=1}^m \lambda_i \lambda_j y_i y_j \Phi(x_i) \cdot \Phi(x_j)$$

$$\text{subject to} \quad \underbrace{0 \leq \lambda_i \leq C}_{\sum_{i=1}^m \lambda_i y_i = 0}, \quad i = 1, \dots, m,$$

$$\sum_{i=1}^m \lambda_i y_i = 0.$$

Handwritten notes:

- $-\lambda_i \leq 0$
- $\lambda_i \geq 0$
- $\lambda_i \leq C$

Recall QP:

$$\text{minimize} \quad \frac{1}{2} x^T P x + q^T x,$$

$$\text{subject to} \quad Gx \leq h,$$

$$Ax = b$$

How to set: x, P, q, G, h, A, b ?

1. Set $x = [\lambda_1, \lambda_2, \dots, \lambda_m]^T$
2. Let $K(x_i, x_j) = \Phi(x_i) \cdot \Phi(x_j)$
3. Set matrix $P_{ij} = y_i y_j K(x_i, x_j)$
4. q is column vector containing -1.
5. For equality constraint, set
 $A = [y_1, y_2, \dots, y_m]^T, b = 0$
6. Inequalities: $\underbrace{-\lambda_i \leq 0}, \underbrace{\lambda_i \leq C}$

Formulate Dual Margin Kernel SVM Problem for QP

Dual Soft-Margin Kernel SVM:

$$\begin{aligned} &\text{Maximize} && \sum_{i=1}^m \lambda_i - \frac{1}{2} \sum_{i,j=1}^m \lambda_i \lambda_j y_i y_j \Phi(x_i) \cdot \Phi(x_j) \\ &\text{subject to} && 0 \leq \lambda_i \leq C, \quad i = 1, \dots, m, \\ &&& \sum_{i=1}^m \lambda_i y_i = 0. \end{aligned}$$

Recall QP:

$$\begin{aligned} &\text{minimize} && \frac{1}{2} x^T P x + q^T x, \\ &\text{subject to} && Gx \leq h, \\ &&& Ax = b \end{aligned}$$

How to set: x, P, q, G, h, A, b ?

1. Set $x = [\lambda_1, \lambda_2, \dots, \lambda_m]^T$
2. Let $K(x_i, x_j) = \Phi(x_i) \cdot \Phi(x_j)$
3. Set matrix $P_{ij} = y_i y_j K(x_i, x_j)$
4. q is column vector containing -1.
5. For equality constraint, set
 $A = [y_1, y_2, \dots, y_m]^T, b = 0$
6. Inequalities: $-\lambda_i \leq 0, \lambda_i \leq C$

7. Set G as

Formulate Dual Margin Kernel SVM Problem for QP

Dual Soft-Margin Kernel SVM:

$$\begin{aligned} & \text{Maximize} \quad \sum_{i=1}^m \lambda_i - \frac{1}{2} \sum_{i,j=1}^m \lambda_i \lambda_j y_i y_j \Phi(x_i) \cdot \Phi(x_j) \\ & \text{subject to} \quad 0 \leq \lambda_i \leq C, \quad i = 1, \dots, m, \\ & \quad \quad \quad \sum_{i=1}^m \lambda_i y_i = 0. \end{aligned}$$

Recall QP:

$$\begin{aligned} & \text{minimize} \quad \frac{1}{2} x^T P x + q^T x, \\ & \text{subject to} \quad Gx \leq h, \\ & \quad \quad \quad Ax = b \end{aligned}$$

How to set: x, P, q, G, h, A, b ?



1. Set $x = [\lambda_1, \lambda_2, \dots, \lambda_m]^T$
2. Let $K(x_i, x_j) = \Phi(x_i) \cdot \Phi(x_j)$
3. Set matrix $P_{ij} = y_i y_j K(x_i, x_j)$
4. q is column vector containing -1.
5. For equality constraint, set $A = [y_1, y_2, \dots, y_m]^T$, $b = 0$
6. Inequalities: $-\lambda_i \leq 0, \lambda_i \leq C$

$h =$

$$\begin{bmatrix} C \\ C \\ C \\ \vdots \\ C \\ 0 \\ 0 \\ \vdots \\ 0 \end{bmatrix}$$

7. Set G as

$$G = \begin{bmatrix} 1 & 0 & \dots & 0 \\ 0 & 1 & \dots & 0 \\ \vdots & \ddots & \dots & \vdots \\ 0 & 0 & \dots & 1 \\ -1 & 0 & \dots & 0 \\ \vdots & \ddots & \dots & \vdots \\ 0 & 0 & \dots & -1 \end{bmatrix}$$

$\begin{bmatrix} \lambda_1 \\ \lambda_2 \\ \vdots \\ \lambda_m \end{bmatrix}$
 x

$-\lambda_i \leq 0$

Formulate Dual Margin Kernel SVM Problem for QP

Dual Soft-Margin Kernel SVM:

$$\begin{aligned} &\text{Maximize} \quad \sum_{i=1}^m \lambda_i - \frac{1}{2} \sum_{i,j=1}^m \lambda_i \lambda_j y_i y_j \Phi(x_i) \cdot \Phi(x_j) \\ &\text{subject to} \quad 0 \leq \lambda_i \leq C, \quad i = 1, \dots, m, \\ &\quad \quad \quad \sum_{i=1}^m \lambda_i y_i = 0. \end{aligned}$$

Recall QP:

$$\begin{aligned} &\text{minimize} \quad \frac{1}{2} x^T P x + q^T x, \\ &\text{subject to} \quad Gx \leq h, \\ &\quad \quad \quad Ax = b \end{aligned}$$

How to set: x, P, q, G, h, A, b ?

1. Set $x = [\lambda_1, \lambda_2, \dots, \lambda_m]^T$
2. Let $K(x_i, x_j) = \Phi(x_i) \cdot \Phi(x_j)$
3. Set matrix $P_{ij} = y_i y_j K(x_i, x_j)$
4. q is column vector containing -1.
5. For equality constraint, set $A = [y_1, y_2, \dots, y_m]^T, b = 0$
6. Inequalities: $-\lambda_i \leq 0, \lambda_i \leq C$

7. Set G as

$$G = \begin{bmatrix} 1 & 0 & \dots & 0 \\ 0 & 1 & \dots & 0 \\ \vdots & \ddots & \dots & \vdots \\ 0 & 0 & \dots & 1 \\ -1 & 0 & \dots & 0 \\ \vdots & \ddots & \dots & \vdots \\ 0 & 0 & \dots & -1 \end{bmatrix}$$

8. $h = [C, C, \dots, C | 0, \dots, 0]^T$

Handwritten notes in orange:

- $T^1 \leq C$
- $T^2 \leq C$
- $T^m \leq C$
- $I \begin{bmatrix} T^1 \\ T^2 \\ \vdots \\ T^m \end{bmatrix} \leq \begin{bmatrix} C \\ C \\ \vdots \\ C \end{bmatrix}$

Algorithms for Dual Soft Margin Kernel SVM

Algorithm 1 Algorithm for solving Dual Kernel SVM using QP

1: Initialization:

- Compute $K = XX^T$, if possible using input space. ✓
- For linear kernel, return K , for polynomial of degree d , return $\frac{1}{d}K^d$. ✓
- For RBF kernel, compute $K = \exp(-(x - x')^2/2\sigma^2)$. ✓

2: Training: Assemble matrices and vectors to solve QP for dual

$$\min_x (x^T P x + q^T x), \quad \text{subject to } Gx \leq h, Ax = b.$$

- Define x, P, q, G, h, b as described in previous slide.

pass to solver.