

Bayesian inference of causal effects with proxy measurements of the interference network

Bar Weinstein
Tel Aviv University


ISDSA Conference, May 2024

Joint work with Daniel Nevo

Outline

- Causal inference 101.
- Networked interference.
- Available proxies of the interference network.
- Inferential framework.

Causal Inference 101




Interference

- Interference occurs when treatments of one unit affect the outcomes of other units.
- Units interact, resulting in the spread of treatment effects.
- Examples include
 - 1 Protection against infectious diseases (Halloran et al. 1995; Hayek et al. 2022)
 - 2 Spread of addictive drugs (Buchanan et al. 2023)
 - 3 A/B testing in online platforms (Aral 2016)
 - 4 Social influence between friends (Bearman et al. 2004; Christakis et al. 2007)
 - 5 Diffusion of financial information (Banerjee et al. 2013)


Interference

- Interference occurs when treatments of one unit affect the outcomes of other units.
- Units interact, resulting in the spread of treatment effects.
- Examples include
 - 1 Protection against infectious diseases (Halloran et al. 1995; Hayek et al. 2022)
 - 2 Spread of addictive drugs (Buchanan et al. 2023)
 - 3 A/B testing in online platforms (Aral 2016)
 - 4 Social influence between friends (Bearman et al. 2004; Christakis et al. 2007)
 - 5 Diffusion of financial information (Banerjee et al. 2013)
- Encompass various terms such as spillovers, peer effects, contamination, etc.
- Interference structure between units can be represented within a network.


Networked interference



Networked interference



Networked interference



Formal settings

- Finite population of units $i \in \{1, \dots, N\}$.
- \mathbf{Z} : population-level treatment vector with treatments' space \mathcal{Z} .
 - For example, for binary treatments $\mathcal{Z} \subseteq \{0, 1\}^N$.
- $Y_i(\mathbf{z})$: *potential* or *counterfactual* outcomes for each $\mathbf{z} \in \mathcal{Z}$.
 - Outcomes under hypothetical intervention that sets $\mathbf{Z} = \mathbf{z}$.

Formal settings

- Finite population of units $i \in \{1, \dots, N\}$.
- \mathbf{Z} : population-level treatment vector with treatments' space \mathcal{Z} .
 - For example, for binary treatments $\mathcal{Z} \subseteq \{0, 1\}^N$.
- $Y_i(\mathbf{z})$: *potential* or *counterfactual* outcomes for each $\mathbf{z} \in \mathcal{Z}$.
 - Outcomes under hypothetical intervention that sets $\mathbf{Z} = \mathbf{z}$.
- $\mathbf{X} \subseteq \mathbb{R}^{N \times q}$: covariates matrix.
- \mathbf{A}^* : adjacency matrix of the true interference network.
 - For simplicity, undirected and unweighted.

Causal estimands

- Comparisons of population-level interventions:

1 **Static.** $\mu(\mathbf{z}) = N^{-1} \sum_{i=1}^N \mathbb{E}[Y_i(\mathbf{z}) | \mathbf{A}^*]$

2 **Dynamic.** $\mu(h) = N^{-1} \sum_{i=1}^N \mathbb{E}[Y_i(h) | \mathbf{A}^*]$, for function $h \equiv h(\mathbf{X}, \mathbf{A}^*)$.

3 **Stochastic.** $\mu(\alpha_0) = N^{-1} \sum_{i=1}^N \sum_{\mathbf{z} \in \mathcal{Z}} \pi_{\alpha_0}(\mathbf{z}) \mathbb{E}[Y_i(\mathbf{z}) | \mathbf{A}^*]$, for stochastic policy $\pi_{\alpha_0}(\mathbf{z})$.

Causal estimands

- Comparisons of population-level interventions:

1 **Static.** $\mu(\mathbf{z}) = N^{-1} \sum_{i=1}^N \mathbb{E}[Y_i(\mathbf{z}) | \mathbf{A}^*]$

2 **Dynamic.** $\mu(h) = N^{-1} \sum_{i=1}^N \mathbb{E}[Y_i(h) | \mathbf{A}^*]$, for function $h \equiv h(\mathbf{X}, \mathbf{A}^*)$.

3 **Stochastic.** $\mu(\alpha_0) = N^{-1} \sum_{i=1}^N \sum_{\mathbf{z} \in \mathcal{Z}} \pi_{\alpha_0}(\mathbf{z}) \mathbb{E}[Y_i(\mathbf{z}) | \mathbf{A}^*]$, for stochastic policy $\pi_{\alpha_0}(\mathbf{z})$.

Assumptions and estimands are expressed via \mathbf{A}^* . Can we accurately measure it?

Measuring A^*

- Accurately measuring networks of social interactions is a formidable task.
 - Social networks from self-reported elicitation of social relations.
 - Measured relations might not reflect those relevant to the studied mechanism.

Measuring A^*

- Accurately measuring networks of social interactions is a formidable task.
 - Social networks from self-reported elicitation of social relations.
 - Measured relations might not reflect those relevant to the studied mechanism.
- However, it is commonly assumed that the measured network accurately represents the interference structure (Forastiere et al. 2020; Ogburn et al. 2022; Tchetgen Tchetgen et al. 2020).
- Misspecified network result in biased estimation (Weinstein et al. 2023).

Measuring \mathbf{A}^*

- Accurately measuring networks of social interactions is a formidable task.
 - Social networks from self-reported elicitation of social relations.
 - Measured relations might not reflect those relevant to the studied mechanism.
- However, it is commonly assumed that the measured network accurately represents the interference structure (Forastiere et al. 2020; Ogburn et al. 2022; Tchetgen Tchetgen et al. 2020).
- Misspecified network result in biased estimation (Weinstein et al. 2023).
- Assume that researchers observe proxy measurements $\mathbf{A} = (\mathbf{A}_1, \dots, \mathbf{A}_B)$ of \mathbf{A}^* .
 - Single or multiple imprecisely measured networks.
 - Multiple networks from varying sources.
 - Multi-layer networks.

Measuring \mathbf{A}^*


- Accurately measuring networks of social interactions is a formidable task.
 - Social networks from self-reported elicitation of social relations.
 - Measured relations might not reflect those relevant to the studied mechanism.
- However, it is commonly assumed that the measured network accurately represents the interference structure (Forastiere et al. 2020; Ogburn et al. 2022; Tchetgen Tchetgen et al. 2020).
- Misspecified network result in biased estimation (Weinstein et al. 2023).
- Assume that researchers observe proxy measurements $\mathbf{A} = (\mathbf{A}_1, \dots, \mathbf{A}_B)$ of \mathbf{A}^* .
 - Single or multiple imprecisely measured networks.
 - Multiple networks from varying sources.
 - Multi-layer networks.
- **Our goal: estimate causal effects using the proxy measurements \mathbf{A} .**

Structural model

- \mathbf{Z}_{-i} affects Y_i only through values of fixed, possibly unknown, summarizing function of neighbors' treatments (Aronow et al. 2017).
- **Assumption 1.** *For any two treatments $\mathbf{z}, \mathbf{z}' \in \mathcal{Z}$, if $z_i = z'_i$ and $f(\mathbf{z}_{-i}, \mathbf{A}_i^*) = f(\mathbf{z}'_{-i}, \mathbf{A}_i^*)$, then $Y_i(\mathbf{z}) = Y_i(\mathbf{z}')$.*

Structural model

- \mathbf{Z}_{-i} affects Y_i only through values of fixed, possibly unknown, summarizing function of neighbors' treatments (Aronow et al. 2017).
- **Assumption 1.** For any two treatments $\mathbf{z}, \mathbf{z}' \in \mathcal{Z}$, if $z_i = z'_i$ and $f(\mathbf{z}_{-i}, \mathbf{A}_i^*) = f(\mathbf{z}'_{-i}, \mathbf{A}_i^*)$, then $Y_i(\mathbf{z}) = Y_i(\mathbf{z}')$.
- Data are generated by sequentially evaluating a set of structural equations.
- Represented by a population-level directed acyclic graph (DAG):



Structural model

- \mathbf{A}^* and \mathbf{X} block all back-door paths between \mathbf{Z} and \mathbf{Y} , i.e., $Y_i(\mathbf{z}) \perp\!\!\!\perp \mathbf{Z} | \mathbf{A}^*, \mathbf{X}$.
 - Accounting for latent network confounding $\mathbf{A}^* \leftarrow \mathbf{U} \rightarrow \mathbf{Y}$ is possible.

Structural model

- \mathbf{A}^* and \mathbf{X} block all back-door paths between \mathbf{Z} and \mathbf{Y} , i.e., $Y_i(\mathbf{z}) \perp\!\!\!\perp \mathbf{Z} | \mathbf{A}^*, \mathbf{X}$.
 - Accounting for latent network confounding $\mathbf{A}^* \leftarrow \mathbf{U} \rightarrow \mathbf{Y}$ is possible.
- Require the specification of the following probabilistic models:
 - 1 *True network.* $p(\mathbf{A}^* | \mathbf{X})$.
 - Random graphs, stochastic block model, ERGM, latent space models, etc.

Structural model

- \mathbf{A}^* and \mathbf{X} block all back-door paths between \mathbf{Z} and \mathbf{Y} , i.e., $Y_i(\mathbf{z}) \perp\!\!\!\perp \mathbf{Z} | \mathbf{A}^*, \mathbf{X}$.
 - Accounting for latent network confounding $\mathbf{A}^* \leftarrow \mathbf{U} \rightarrow \mathbf{Y}$ is possible.
- Require the specification of the following probabilistic models:
 - 1 *True network.* $p(\mathbf{A}^* | \mathbf{X})$.
 - Random graphs, stochastic block model, ERGM, latent space models, etc.
 - 2 *Observed proxies.* $p(\mathbf{A} | \mathbf{A}^*, \mathbf{X})$.
 - Measurement error models (differential or not) (e.g., Young et al. 2021).
 - Multi-layer network models with \mathbf{A}^* being the “consensus” network (e.g., Sosa et al. 2022).

Structural model

- \mathbf{A}^* and \mathbf{X} block all back-door paths between \mathbf{Z} and \mathbf{Y} , i.e., $Y_i(\mathbf{z}) \perp\!\!\!\perp \mathbf{Z} | \mathbf{A}^*, \mathbf{X}$.
 - Accounting for latent network confounding $\mathbf{A}^* \leftarrow \mathbf{U} \rightarrow \mathbf{Y}$ is possible.
- Require the specification of the following probabilistic models:
 - 1 *True network.* $p(\mathbf{A}^* | \mathbf{X})$.
 - Random graphs, stochastic block model, ERGM, latent space models, etc.
 - 2 *Observed proxies.* $p(\mathbf{A} | \mathbf{A}^*, \mathbf{X})$.
 - Measurement error models (differential or not) (e.g., Young et al. 2021).
 - Multi-layer network models with \mathbf{A}^* being the “consensus” network (e.g., Sosa et al. 2022).
 - 3 *Outcome model.* $p(\mathbf{Y} | \mathbf{Z}, \mathbf{A}^*, \mathbf{X})$.

Structural model

- \mathbf{A}^* and \mathbf{X} block all back-door paths between \mathbf{Z} and \mathbf{Y} , i.e., $Y_i(\mathbf{z}) \perp\!\!\!\perp \mathbf{Z} | \mathbf{A}^*, \mathbf{X}$.
 - Accounting for latent network confounding $\mathbf{A}^* \leftarrow \mathbf{U} \rightarrow \mathbf{Y}$ is possible.
- Require the specification of the following probabilistic models:
 - 1 *True network.* $p(\mathbf{A}^* | \mathbf{X})$.
 - Random graphs, stochastic block model, ERGM, latent space models, etc.
 - 2 *Observed proxies.* $p(\mathbf{A} | \mathbf{A}^*, \mathbf{X})$.
 - Measurement error models (differential or not) (e.g., Young et al. 2021).
 - Multi-layer network models with \mathbf{A}^* being the “consensus” network (e.g., Sosa et al. 2022).
 - 3 *Outcome model.* $p(\mathbf{Y} | \mathbf{Z}, \mathbf{A}^*, \mathbf{X})$.
- Few limitations (inc. dependence between units). The primary constraint is the computational burden.

Nonparameteric identification

- Observed outcomes Y_i are related to the potential outcomes $Y_i(\mathbf{z})$ as follows.
- **Assumption 2** (Consistency). *If $\mathbf{Z} = \mathbf{z}$, then $Y_i = Y_i(\mathbf{z})$, $\forall \mathbf{z} \in \mathcal{Z}$*

Nonparameteric identification

- Observed outcomes Y_i are related to the potential outcomes $Y_i(\mathbf{z})$ as follows.
- **Assumption 2** (Consistency). *If $\mathbf{Z} = \mathbf{z}$, then $Y_i = Y_i(\mathbf{z})$, $\forall \mathbf{z} \in \mathcal{Z}$*
- Causal estimands are identified via: $\mathbb{E}[Y_i(\mathbf{z})|\mathbf{A}^*] = \mathbb{E}_{\mathbf{X}}\mathbb{E}[Y_i|\mathbf{Z} = \mathbf{z}, \mathbf{A}^*, \mathbf{X}]$.

Nonparameteric identification

- Observed outcomes Y_i are related to the potential outcomes $Y_i(\mathbf{z})$ as follows.
- **Assumption 2** (Consistency). *If $\mathbf{Z} = \mathbf{z}$, then $Y_i = Y_i(\mathbf{z})$, $\forall \mathbf{z} \in \mathcal{Z}$*
- Causal estimands are identified via: $\mathbb{E}[Y_i(\mathbf{z})|\mathbf{A}^*] = \mathbb{E}_{\mathbf{X}}\mathbb{E}[Y_i|\mathbf{Z} = \mathbf{z}, \mathbf{A}^*, \mathbf{X}]$.
- Further assuming that Y_i depends on \mathbf{A}^* through summary statistics:
- **Assumption 3.** *For some fixed functions ϕ_1, ϕ_2, ϕ_3 ,*

$$p(Y_i|\mathbf{Z}, \mathbf{A}^*, \mathbf{X}) = p(Y_i|Z_i, f(\phi_1(\mathbf{Z}_{-i}, \mathbf{A}_i^*)), \mathbf{X}_i, \phi_2(\mathbf{X}_{-i}, \mathbf{A}_i^*), \phi_{3,i}(\mathbf{A}^*)), \forall i$$

- yields

$$\mathbb{E}[Y_i(\mathbf{z})|\mathbf{A}^*] = \mathbb{E}_{\mathbf{X}}\mathbb{E}[Y_i|Z_i, f(\phi_1), \mathbf{X}_i, \phi_2, \phi_{3,i}]$$

Bayesian inference

- We propose a Bayesian framework for inference.
- Denote the observed data by $\mathbf{O} = (\mathbf{Y}, \mathbf{Z}, \mathbf{X}, \mathbf{A})$.

Bayesian inference

- We propose a Bayesian framework for inference.
- Denote the observed data by $\mathbf{O} = (\mathbf{Y}, \mathbf{Z}, \mathbf{X}, \mathbf{A})$.
- Let η, θ, γ parameterize the outcome, network, and proxies distributions, respectively. They can be in a finite or infinite space. Assume prior independence.

Bayesian inference

- We propose a Bayesian framework for inference.
- Denote the observed data by $\mathbf{O} = (\mathbf{Y}, \mathbf{Z}, \mathbf{X}, \mathbf{A})$.
- Let η, θ, γ parameterize the outcome, network, and proxies distributions, respectively. They can be in a finite or infinite space. Assume prior independence.
- Posterior can be written as

$$\begin{aligned}
 p(\eta, \theta, \gamma, \mathbf{A}^* | \mathbf{O}) &\propto p(\mathbf{Y} | \mathbf{Z}, \mathbf{A}^*, \mathbf{X}, \eta) p(\eta) \\
 &\quad \cdot p(\mathbf{A} | \mathbf{A}^*, \mathbf{X}, \gamma) p(\gamma) \\
 &\quad \cdot p(\mathbf{A}^* | \mathbf{X}, \theta) p(\theta),
 \end{aligned}$$

- Mixed space of continuous and discrete latent variables. Discrete space consists of $\mathcal{O}(2^{N^2})$ terms!

Sampling from the posterior

- Metropolis-Hasting (and Gibbs) have slow convergence and poor mixing.
- Modern methods, e.g., Hamiltonian Monte Carlo, require continuous posterior. Extension for mixed spaces doesn't scale well (e.g., Nishimura et al. 2020; Zhou 2020).

Sampling from the posterior

- Metropolis-Hasting (and Gibbs) have slow convergence and poor mixing.
- Modern methods, e.g., Hamiltonian Monte Carlo, require continuous posterior. Extension for mixed spaces doesn't scale well (e.g., Nishimura et al. 2020; Zhou 2020).
- We propose a Bayesian modularization approach (Bayarri et al. 2009; Jacob et al. 2017). Posterior can be written as:

$$p(\eta, \theta, \gamma, \mathbf{A}^* | \mathbf{O}) \propto \underbrace{p(\eta | \mathbf{Y}, \mathbf{Z}, \mathbf{A}^*, \mathbf{X})}_{\text{Outcome module}} \underbrace{p(\theta, \gamma, \mathbf{A}^* | \mathbf{X}, \mathbf{A})}_{\text{Network module}} \underbrace{p(\mathbf{Y} | \mathbf{Z}, \mathbf{A}^*, \mathbf{X})}_{\text{Feedback term}}$$

Sampling from the posterior

- Metropolis-Hasting (and Gibbs) have slow convergence and poor mixing.
- Modern methods, e.g., Hamiltonian Monte Carlo, require continuous posterior. Extension for mixed spaces doesn't scale well (e.g., Nishimura et al. 2020; Zhou 2020).
- We propose a Bayesian modularization approach (Bayarri et al. 2009; Jacob et al. 2017). Posterior can be written as:

$$p(\eta, \theta, \gamma, \mathbf{A}^* | \mathbf{O}) \propto \underbrace{p(\eta | \mathbf{Y}, \mathbf{Z}, \mathbf{A}^*, \mathbf{X})}_{\text{Outcome module}} \underbrace{p(\theta, \gamma, \mathbf{A}^* | \mathbf{X}, \mathbf{A})}_{\text{Network module}} \underbrace{p(\mathbf{Y} | \mathbf{Z}, \mathbf{A}^*, \mathbf{X})}_{\text{Feedback term}}$$

- Use the cut-posterior that removes the feedback between modules:

$$\tilde{p}(\eta, \theta, \gamma, \mathbf{A}^* | \mathbf{O}) \propto p(\eta | \mathbf{Y}, \mathbf{Z}, \mathbf{A}^*, \mathbf{X}) p(\theta, \gamma, \mathbf{A}^* | \mathbf{X}, \mathbf{A})$$

Sampling from the posterior

- Sampling from the posterior is performed sequentially by sampling from each module:

- 1 **Network.** $p(\theta, \gamma, \mathbf{A}^* | \mathbf{X}, \mathbf{A}) = p(\mathbf{A}^* | \mathbf{X}, \mathbf{A}, \theta, \gamma) \sum_{\mathbf{A}^*} p(\theta, \gamma, \mathbf{A}^* | \mathbf{X}, \mathbf{A}) \Rightarrow$ can be simplified!
- 2 **Outcome.** $p(\eta | \mathbf{Y}, \mathbf{Z}, \mathbf{A}^*, \mathbf{X}) \propto p(\eta) p(\mathbf{Y} | \mathbf{Z}, \mathbf{A}^*, \mathbf{X}, \eta) \Rightarrow$ Bayesian regression.

Sampling from the posterior

- Sampling from the posterior is performed sequentially by sampling from each module:
 - 1 Network.** $p(\theta, \gamma, \mathbf{A}^* | \mathbf{X}, \mathbf{A}) = p(\mathbf{A}^* | \mathbf{X}, \mathbf{A}, \theta, \gamma) \sum_{\mathbf{A}^*} p(\theta, \gamma, \mathbf{A}^* | \mathbf{X}, \mathbf{A}) \Rightarrow$ can be simplified!
 - 2 Outcome.** $p(\eta | \mathbf{Y}, \mathbf{Z}, \mathbf{A}^*, \mathbf{X}) \propto p(\eta) p(\mathbf{Y} | \mathbf{Z}, \mathbf{A}^*, \mathbf{X}, \eta) \Rightarrow$ Bayesian regression.
- In practice, performed via either:
 - Sample (θ_m, γ_m) . Then, sample \mathbf{A}_m^* . For each \mathbf{A}_m^* , sample η from the outcome module. Networks can be sampled via either
 - For each (θ_m, γ_m) , sample one network ("*Three-stage*").
 - Sample networks using the posterior mean $\mathbb{E}[\theta, \gamma | \cdot]$ ("*Two-stage*").

Sampling from the posterior


- Sampling from the posterior is performed sequentially by sampling from each module:
 - Network.** $p(\theta, \gamma, \mathbf{A}^* | \mathbf{X}, \mathbf{A}) = p(\mathbf{A}^* | \mathbf{X}, \mathbf{A}, \theta, \gamma) \sum_{\mathbf{A}^*} p(\theta, \gamma, \mathbf{A}^* | \mathbf{X}, \mathbf{A}) \Rightarrow$ can be simplified!
 - Outcome.** $p(\eta | \mathbf{Y}, \mathbf{Z}, \mathbf{A}^*, \mathbf{X}) \propto p(\eta) p(\mathbf{Y} | \mathbf{Z}, \mathbf{A}^*, \mathbf{X}, \eta) \Rightarrow$ Bayesian regression.
- In practice, performed via either:
 - Sample (θ_m, γ_m) . Then, sample \mathbf{A}_m^* . For each \mathbf{A}_m^* , sample η from the outcome module. Networks can be sampled via either
 - For each (θ_m, γ_m) , sample one network ("*Three-stage*").
 - Sample networks using the posterior mean $\mathbb{E}[\theta, \gamma | \cdot]$ ("*Two-stage*").
 - Sample multiple \mathbf{A}^* . Estimate summary statistics $\hat{\phi} = M^{-1} \sum_m \phi(\mathbf{Z}, \mathbf{X}, \mathbf{A}_m^*)$. Sample η by replacing ϕ with the plug-in estimator $\hat{\phi}$ ("*Plug-in*").

Numerical illustration



- Implemented in the probabilistic programming language *NumPyro*.
- Observed proxy network from a random noise measurement error model. $N = 300$.

Numerical illustration

- Implemented in the probabilistic programming language *NumPyro*.
- Observed proxy network from a random noise measurement error model. $N = 300$.



Numerical illustration



THANK YOU!

 GitHub:// barwein

 barwein@mail.tau.ac.il

Part I

Appendix

Positivity assumption

Assumption (positivity). $p(\mathbf{Z}_i = z, f(\mathbf{Z}_{-i}, \mathbf{A}_i^*) = c \mid \mathbf{X} = \mathbf{x}) > 0, \forall i, z \in \{0, 1\}, c \in \mathcal{C}, \mathbf{x}.$

Nonparameteric identification

$$\begin{aligned}\mathbb{E}[Y_i(\mathbf{z})|\mathbf{A}^*] &= \mathbb{E}_{\mathbf{X}}\mathbb{E}[Y_i(\mathbf{z})|\mathbf{A}^*, \mathbf{X}] \\ &= \mathbb{E}_{\mathbf{X}}\mathbb{E}[Y_i(\mathbf{z})|\mathbf{Z} = \mathbf{z}, \mathbf{A}^*, \mathbf{X}] \\ &= \mathbb{E}_{\mathbf{X}}\mathbb{E}[Y_i|\mathbf{Z} = \mathbf{z}, \mathbf{A}^*, \mathbf{X}] \\ &= \mathbb{E}_{\mathbf{X}}\mathbb{E}[Y_i|z_i, f(\phi_1(\mathbf{z}_{-i}, \mathbf{A}_i^*)), \mathbf{X}_i, \phi_2(\mathbf{X}_{-i}, \mathbf{A}_i^*), \phi_{3,i}(\mathbf{A}^*)]\end{aligned}$$

Nonparameteric identification – stochastic estimand

$$\begin{aligned} N^{-1} \sum_{i=1}^N \sum_{\mathbf{z} \in \mathcal{Z}} \pi_{\alpha_0}(\mathbf{z}) \mathbb{E}[Y_i(\mathbf{z}) | \mathbf{A}^*] &= N^{-1} \sum_{\mathbf{z} \in \mathcal{Z}} \pi_{\alpha_0}(\mathbf{z}) \sum_{i=1}^N \mathbb{E}[Y_i(\mathbf{z}) | \mathbf{A}^*] \\ &= N^{-1} \sum_{\mathbf{z} \in \mathcal{Z}} \pi_{\alpha_0}(\mathbf{z}) \sum_{i=1}^N \mathbb{E}_{\mathbf{X}} \mathbb{E}[Y_i | \mathbf{Z} = \mathbf{z}, \mathbf{A}^*, \mathbf{X}], \end{aligned}$$

DAG factorization

$$p(\mathbf{Y}, \mathbf{Z}, \mathbf{A}, \mathbf{A}^*, \mathbf{X}) = p(\mathbf{Y}|\mathbf{Z}, \mathbf{A}^*, \mathbf{X})p(\mathbf{Z}|\mathbf{X}, \mathbf{A}, \mathbf{A}^*)p(\mathbf{A}|\mathbf{A}^*, \mathbf{X})p(\mathbf{A}^*|\mathbf{X})p(\mathbf{X}),$$

Example – random noise

We observe single \mathbf{A} generated by measuring \mathbf{A}^* with random error independently across edges with false positive and negative rates γ_0, γ_1 , respectively. Namely,

$\Pr(A_{ij} = 1 - k | A_{ij}^* = k) = \gamma_k$, $k = 0, 1$, and we can write

$$p(\mathbf{A} | \mathbf{A}^*, \mathbf{X}, \gamma) = \prod_{i>j} \sum_{k=0,1} \mathbb{I}\{A_{ij}^* = k\} \gamma_k^{A_{ij}} (1 - \gamma_k)^{1-A_{ij}} = \prod_{i>j} \xi_{ij}(A_{ij}^*, \gamma),$$

where $\xi_{ij}(k, \gamma) = \gamma_k^{A_{ij}} (1 - \gamma_k)^{1-A_{ij}}$, $k = 0, 1$. Extending to heterogeneous random noise that depends on covariates is possible as well.

Example – Repeated noisy measurements

Researchers observe $\mathbf{A} = (\mathbf{A}_1, \dots, \mathbf{A}_B)$ networks each pertain to different noisy measurement of the true network (De Bacco et al. 2023; Redhead et al. 2023). A model can be assumed for each of the networks:

$$p(\mathbf{A}_b | \mathbf{A}^*, \mathbf{X}, \gamma_b) = \prod_{i>j} \xi_{b,ij}(\mathbf{A}_{ij}^*, \gamma_b),$$

for some function $\xi_{b,ij}$. The joint distribution of $\mathbf{A}_1, \dots, \mathbf{A}_B$ can be assumed to be independent or dependent (e.g., via a hierarchical model).

Example – multi-layer networks

Researchers observe multi-layer collection of networks $\mathbf{A} = (\mathbf{A}_1, \dots, \mathbf{A}_B)$. Each \mathbf{A}_b measures different social relationships. It is possible to model the observed multi-layer networks using LSM with shared latent positions:

$$p(\mathbf{A}_b | \mathbf{X}, \gamma_b, \mathbf{w}) = \prod_{i>j} \psi(\gamma_b - \|\mathbf{w}_i - \mathbf{w}_j\|)^{A_{b,ij}} (1 - \psi(\gamma_b - \|\mathbf{w}_i - \mathbf{w}_j\|))^{1-A_{b,ij}},$$

for $\psi : \mathbb{R} \rightarrow [0, 1]$ and latent positions $\mathbf{w}_i \in \mathbb{R}^d$. Covariates can also be included. Dependence between the networks can be modeled with multinomial (Salter-Townshend et al. 2017) or hierarchical (Sosa et al. 2022) models. The true network \mathbf{A}^* can be viewed as a “consensus” or aggregate version of the observed networks. That can be achieved with a simple union (Salter-Townshend et al. 2017), or in a model-based approach (Sosa et al. 2022).

Feedback between modules

The posterior can be written as: $p(\eta, \theta, \gamma, \mathbf{A}^* | \mathbf{O}) = p(\eta | \mathbf{O}, \theta, \gamma, \mathbf{A}^*)p(\theta, \gamma, \mathbf{A}^* | \mathbf{O})$. Thus

$$\begin{aligned} p(\theta, \gamma, \mathbf{A}^* | \mathbf{O}) &= \int_{\eta} p(\eta, \theta, \gamma, \mathbf{A}^* | \mathbf{O}) d\eta \\ &\propto p(\theta)p(\gamma)p(\mathbf{A} | \mathbf{A}^*, \mathbf{X}, \gamma)p(\mathbf{A}^* | \mathbf{X}, \theta)p(\mathbf{Y} | \mathbf{Z}, \mathbf{A}^*, \mathbf{X}) \\ &\equiv p(\theta, \gamma, \mathbf{A}^* | \mathbf{X}, \mathbf{A})p(\mathbf{Y} | \mathbf{Z}, \mathbf{A}^*, \mathbf{X}) \end{aligned}$$

where $p(\mathbf{Y} | \mathbf{Z}, \mathbf{A}^*, \mathbf{X}) = \int_{\eta} p(\mathbf{Y} | \mathbf{Z}, \mathbf{A}^*, \mathbf{X}, \eta)p(\eta) d\eta$. In addition,

$$p(\eta | \mathbf{O}, \theta, \gamma, \mathbf{A}^*) = \frac{p(\eta, \theta, \gamma, \mathbf{A}^* | \mathbf{O})}{p(\theta, \gamma, \mathbf{A}^* | \mathbf{O})} \propto p(\mathbf{Y} | \mathbf{Z}, \mathbf{A}^*, \mathbf{X}, \eta)p(\eta)$$

Sampling from network module

Assume that both the network generation model $p(\mathbf{A}^*|\mathbf{X}, \theta)$ and the observed network model $p(\mathbf{A}|\mathbf{A}^*, \mathbf{X}, \gamma)$ are dyad independent, i.e., can be written as

$$p(\mathbf{A}^*|\mathbf{X}, \theta) = \prod_{i>j} \nu_{ij}(\mathbf{A}_{ij}^*, \theta), \quad p(\mathbf{A}|\mathbf{A}^*, \mathbf{X}, \gamma) = \prod_{i>j} \xi_{ij}(\mathbf{A}_{ij}^*, \gamma),$$

for some functions ν_{ij}, ξ_{ij} . Thus,

$$\begin{aligned} p(\theta, \gamma|\mathbf{X}, \mathbf{A}) &\propto p(\theta)p(\gamma) \sum_{\mathbf{A}^*} \prod_{i>j} \xi_{ij}(\mathbf{A}_{ij}^*, \gamma) \nu_{ij}(\mathbf{A}_{ij}^*, \theta) \\ &= p(\theta)p(\gamma) \prod_{i>j} \sum_{k=0,1} \xi_{ij}(k, \gamma) \nu_{ij}(k, \theta) \end{aligned}$$

Sampling networks

Generating samples of \mathbf{A}^* is reduced to sampling edges independently from

$$\begin{aligned} p(\mathbf{A}^* | \mathbf{X}, \mathbf{A}, \theta, \gamma) &= \frac{p(\theta, \gamma, \mathbf{A}^* | \mathbf{X}, \mathbf{A})}{\sum_{\mathbf{A}^*} p(\theta, \gamma, \mathbf{A}^* | \mathbf{X}, \mathbf{A})} \\ &= \prod_{i>j} \frac{\xi_{ij}(\mathbf{A}_{ij}^*, \gamma) \nu_{ij}(\mathbf{A}_{ij}^*, \theta)}{\sum_{k=0,1} \xi_{ij}(k, \gamma) \nu_{ij}(k, \theta)}, \end{aligned}$$

Three-stage sampling

Algorithm 1: Three-stage sampling

```
foreach  $m = 1, \dots, M$  do  
    Sample  $\theta_m, \gamma_m \sim p(\theta, \gamma | \mathbf{X}, \mathbf{A})$  ;  
    Sample  $\mathbf{A}_m^* \sim p(\mathbf{A}^* | \mathbf{X}, \mathbf{A}, \theta_m, \gamma_m)$  ;  
    foreach  $\ell = 1, \dots, L$  do  
        Sample  $\eta_\ell \sim p(\eta | \mathbf{Y}, \mathbf{Z}, \mathbf{A}_m^*, \mathbf{X})$  ;  
    end  
end
```

Two-stage sampling

Algorithm 2: Two-stage sampling

foreach $m = 1, \dots, M$ **do**

 | Sample $\theta_m, \gamma_m \sim p(\theta, \gamma | \mathbf{X}, \mathbf{A})$;

end

Compute posterior means $\bar{\theta} = M^{-1} \sum_m \theta_m$ and similarly for $\bar{\gamma}$;

foreach $m = 1, \dots, M$ **do**

 | Sample $\mathbf{A}_m^* \sim p(\mathbf{A}^* | \mathbf{X}, \mathbf{A}, \bar{\theta}, \bar{\gamma})$;

foreach $\ell = 1, \dots, L$ **do**

 | Sample $\eta_\ell \sim p(\eta | \mathbf{Y}, \mathbf{Z}, \mathbf{A}_m^*, \mathbf{X})$;

end

end

Plugin sampling

Algorithm 3: Plug-in sampling

foreach $m = 1, \dots, M$ **do**

 | Sample $\theta_m, \gamma_m \sim p(\theta, \gamma | \mathbf{X}, \mathbf{A})$;

 | Sample $\mathbf{A}_m^* \sim p(\mathbf{A}^* | \mathbf{X}, \mathbf{A}, \theta_m, \gamma_m)$;

end

Estimate the outcome models' summary statistics

$$\mathbf{A}^* \equiv \phi(\mathbf{Z}, \mathbf{A}^*, \mathbf{X}) = M^{-1} \sum_m \phi(\mathbf{Z}, \mathbf{A}_m^*, \mathbf{X}) ;$$

foreach $\ell = 1, \dots, L$ **do**

 | Sample $\eta_\ell \sim p(\eta | \mathbf{Y}, \mathbf{Z}, \mathbf{A}^*, \mathbf{X})$;

end

Simulations DGP

- Outcome model $Y_i = \eta_0 + \eta_1 Z_i + \eta_2 \sum_{j \neq i} Z_j A_{ij}^* + \eta_3 X_i + \varepsilon_i$
- Network generation $\text{logit}(\Pr(A_{ij}^* = 1)) = \theta_0 + \theta_1 |X_i - X_j|$.
- Observed network from a random noise measurement error model.
- $N = 300$ for 300 iterations.

References I

Aral, S. (2016). "Networked experiments". In: *The Oxford handbook of the economics of networks*, pp. 376–411.

Aronow, P. M. and C. Samii (Dec. 2017). "Estimating average causal effects under general interference, with application to a social network experiment". en. In: *The Annals of Applied Statistics* 11.4. ISSN: 1932-6157. DOI: 10.1214/16-AOAS1005. URL: <https://projecteuclid.org/journals/annals-of-applied-statistics/volume-11/issue-4/Estimating-average-causal-effects-under-general-interference-with-application-to/10.1214/16-AOAS1005.full> (visited on 08/01/2022).

Banerjee, A. et al. (July 2013). "The Diffusion of Microfinance". In: *Science* 341.6144. DOI: 10.1126/science.1236498.

Bayarri, M. J., J. O. Berger, and F. Liu (Mar. 2009). "Modularization in Bayesian analysis, with emphasis on analysis of computer models". In: *Bayesian Analysis* 4.1. Publisher: International Society for Bayesian Analysis, pp. 119–150. ISSN: 1936-0975, 1931-6690. DOI: 10.1214/09-BA404. URL: <https://projecteuclid.org/journals/bayesian-analysis/volume-4/issue-1/Modularization-in-Bayesian-analysis-with-emphasis-on-analysis-of-computer/10.1214/09-BA404.full> (visited on 03/18/2024).

Bearman, P. S. and J. Moody (2004). "Suicide and Friendships Among American Adolescents". In: *American Journal of Public Health* 94.1. PMID: 14713704, pp. 89–95. DOI: 10.2105/AJPH.94.1.89. eprint: <https://doi.org/10.2105/AJPH.94.1.89>. URL: <https://doi.org/10.2105/AJPH.94.1.89>.

Buchanan, A. L. et al. (Feb. 2023). "Methods for Assessing Spillover in Network-Based Studies of HIV/AIDS Prevention among People Who Use Drugs". In: *Pathogens* 12.2, p. 326. DOI: 10.3390/pathogens12020326.

Christakis, N. A. and J. H. Fowler (July 2007). "The Spread of Obesity in a Large Social Network over 32 Years". In: *New England Journal of Medicine* 357.4, pp. 370–379. DOI: 10.1056/nejmsa066082.

References II

- De Bacco, C. et al. (2023). "Latent network models to account for noisy, multiply reported social network data". In: *Journal of the Royal Statistical Society Series A: Statistics in Society* 186.3, pp. 355–375.
- Forastiere, L., E. M. Airoidi, and F. Mealli (June 2020). "Identification and Estimation of Treatment and Interference Effects in Observational Studies on Networks". In: *Journal of the American Statistical Association* 116.534, pp. 901–918. DOI: [10.1080/01621459.2020.1768100](https://doi.org/10.1080/01621459.2020.1768100).
- Halloran, M. E. and C. J. Struchiner (Mar. 1995). "Causal Inference in Infectious Diseases". In: *Epidemiology* 6.2, pp. 142–151. DOI: [10.1097/00001648-199503000-00010](https://doi.org/10.1097/00001648-199503000-00010).
- Hayek, S. et al. (2022). "Indirect protection of children from SARS-CoV-2 infection through parental vaccination". In: *Science* 375.6585, pp. 1155–1159.
- Jacob, P. E. et al. (Aug. 2017). "Better together? Statistical learning in models made of modules". In: arXiv:1708.08719 [stat]. DOI: [10.48550/arXiv.1708.08719](https://doi.org/10.48550/arXiv.1708.08719). URL: <http://arxiv.org/abs/1708.08719> (visited on 03/18/2024).
- Nishimura, A., D. B. Dunson, and J. Lu (June 2020). "Discontinuous Hamiltonian Monte Carlo for discrete parameters and discontinuous likelihoods". In: *Biometrika* 107.2, pp. 365–380. ISSN: 0006-3444. DOI: [10.1093/biomet/asz083](https://doi.org/10.1093/biomet/asz083). URL: <https://doi.org/10.1093/biomet/asz083> (visited on 03/14/2024).
- Ogburn, E. L. et al. (2022). "Causal Inference for Social Network Data". In: *Journal of the American Statistical Association*, pp. 1–46. DOI: [10.1080/01621459.2022.2131557](https://doi.org/10.1080/01621459.2022.2131557). eprint: <https://doi.org/10.1080/01621459.2022.2131557>. URL: <https://doi.org/10.1080/01621459.2022.2131557>.

References III

- Redhead, D., R. McElreath, and C. T. Ross (2023). "Reliable network inference from unreliable data: A tutorial on latent network modeling using STRAND.". In: *Psychological methods*.
- Salter-Townshend, M. and T. H. McCormick (2017). "Latent space models for multiview network data". In: *The annals of applied statistics* 11.3, p. 1217.
- Sosa, J. and B. Betancourt (May 2022). "A latent space model for multilayer network data". In: *Computational Statistics & Data Analysis* 169, p. 107432. ISSN: 0167-9473. DOI: 10.1016/j.csda.2022.107432.
- Tchetgen Tchetgen, E. J., I. R. Fulcher, and I. Shpitser (Oct. 2020). "Auto-G-Computation of Causal Effects on a Network". In: *Journal of the American Statistical Association* 116.534, pp. 833–844. DOI: 10.1080/01621459.2020.1811098.
- Weinstein, B. and D. Nevo (2023). "Causal inference with misspecified network interference structure". In: *arXiv preprint arXiv:2302.11322*.
- Young, J.-G., G. T. Cantwell, and M. E. J. Newman (Mar. 2021). "Bayesian inference of network structure from unreliable data". en. In: *Journal of Complex Networks* 8.6. arXiv:2008.03334 [physics, stat], cnaa046. ISSN: 2051-1310, 2051-1329. DOI: 10.1093/comnet/cnaa046. URL: <http://arxiv.org/abs/2008.03334> (visited on 08/01/2022).
- Zhou, G. (2020). "Mixed Hamiltonian Monte Carlo for Mixed Discrete and Continuous Variables". In: *Advances in Neural Information Processing Systems*. Vol. 33. Curran Associates, Inc., pp. 17094–17104. URL: <https://proceedings.neurips.cc/paper/2020/hash/c6a01432c8138d46ba39957a8250e027-Abstract.html> (visited on 03/05/2024).