

# BLG202E HOMEWORK № 4

Başar Demir, 150180080

31/05/2020

## 1 Problem 1

### 1.1 Newton's Divided Difference Method

$$p_n(x) = \sum_{j=0}^n \left( f[x_0, x_1, \dots, x_j] \prod_{i=0}^{j-1} (x - x_i) \right)$$

The equation that is given above is generalized Newton's Divided Difference Interpolation formula. We need a quadratic interpolation, so  $n = 2$ . Our data points are,

- $x_0 = 0.00, y_0 = 1.20$
- $x_1 = -0.60, y_1 = 1.04$
- $x_2 = -1.20, y_2 = 0.00$

If we apply Newton's Divided Difference Method to our data points:

For  $j=0$ :

$$c_0 = f(x_0) \rightarrow c_0 = \mathbf{1.20}$$

For  $j=1$ :

$$c_1 = \frac{f(x_1) - f(x_0)}{x_1 - x_0} \rightarrow c_1 = \frac{f(-0.60) - f(0.00)}{-0.60 - 0.00} = \frac{4}{15} \approx \mathbf{0.26}$$

$$c_1(x - x_0) = 0.26(x)$$

For  $j=2$ :

$$c_2 = \frac{\frac{f(x_2) - f(x_1)}{x_2 - x_1} - \frac{f(x_1) - f(x_0)}{x_1 - x_0}}{x_2 - x_0} \rightarrow c_2 = \frac{\frac{f(-1.20) - f(-0.60)}{-1.20 + 0.60} - \frac{f(-0.60) - f(0)}{-0.60 - 0}}{-1.20 - 0}$$

$$c_2 = \frac{\frac{1.04}{0.60} - \frac{0.16}{0.60}}{-1.20} \approx \mathbf{-1.22}$$

$$c_2(x - x_0)(x - x_1) = -1.76(x)(x + 0.60)$$

If we sum all results, we get our interpolation equation:

$$p(x) = 1.20 + 0.26(x) - 1.22(x)(x + 0.60)$$

Interpolated value of point  $x=-1.04$ :

$$p(1.04) = 1.20 + 0.26(-1.04) - 1.22(1.04)(-1.04 + 0.60) \approx \mathbf{0.371}$$

In dataset  $f(-1.04)=0.60$ , therefore absolute value of the relative error is

$$\frac{|0.60 - 0.371|}{|0.60|} \approx \mathbf{0.38166}$$

## 2 Deriving Formulas Using Taylor Series

With the help of Taylor Series, we can determine derivation any point of the function without differentiation with a small truncation error.

### 2.0.1 Second Order Centered Difference Formula

We are trying to find derivative of  $x = x_0$  by using  $x = x_0 + h$  and  $x = x_0 - h$ . By Taylor Formula:

$$\begin{aligned} f(x_0 + h) &= f(x_0) + hf'(x_0) + \frac{h^2}{2}f''(x_0) + \frac{h^3}{6}f'''(x_0) \\ f(x_0 - h) &= f(x_0) - hf'(x_0) + \frac{h^2}{2}f''(x_0) - \frac{h^3}{6}f'''(x_0) \end{aligned}$$

If we subtract second from the first and take the derivative to left:

$$f'(x_0) = \frac{f(x_0 + h) - f(x_0 - h)}{2h} - \frac{h^2}{6}f'''(x_0)$$

Truncation error which is second order:

$$\varepsilon(h) = \frac{h^2}{6}f'''(x_0)$$

### 2.0.2 Fourth Order Centered Difference Formula

We are trying to find derivative of  $x = x_0$  by using  $x = x_0 + h, x = x_0 - h, x = x_0 + 2h, x = x_0 - 2h$ .

By Taylor Formula:

For  $x = x_0 + h, x = x_0 - h$ :

$$\begin{aligned} f(x_0 + h) &= f(x_0) + hf'(x_0) + \frac{h^2}{2}f''(x_0) + \frac{h^3}{6}f'''(x_0) + \frac{h^4}{24}f^{iv}(x_0) + \frac{h^5}{120}f^v(x_0) + \frac{h^6}{720}f^{vi}(x_0) \\ f(x_0 - h) &= f(x_0) - hf'(x_0) + \frac{h^2}{2}f''(x_0) - \frac{h^3}{6}f'''(x_0) + \frac{h^4}{24}f^{iv}(x_0) - \frac{h^5}{120}f^v(x_0) + \frac{h^6}{720}f^{vi}(x_0) \end{aligned}$$

If we subtract second from the first:

$$\begin{aligned} f(x_0 + h) - f(x_0 - h) &= 2hf'(x_0) + \frac{2h^3}{6}f'''(x_0) + \frac{2h^5}{120}f^v(x_0) \\ 1)f'(x_0) &= \frac{f(x_0 + h) - f(x_0 - h)}{2h} - \frac{h^2}{6}f'''(x_0) - \frac{h^4}{120}f^v(x_0) \end{aligned}$$

For  $x = x_0 + 2h, x = x_0 - 2h$ :

$$f(x_0 + 2h) = f(x_0) + 2hf'(x_0) + \frac{4h^2}{2}f''(x_0) + \frac{8h^3}{6}f'''(x_0) + \frac{16h^4}{24}f^{iv}(x_0) + \frac{32h^5}{120}f^v(x_0) + \frac{64h^6}{720}f^{vi}(x_0)$$

$$f(x_0-2h) = f(x_0) - 2hf'(x_0) + \frac{4h^2}{2}f''(x_0) - \frac{8h^3}{6}f'''(x_0) + \frac{16h^4}{24}f^{iv}(x_0) - \frac{32h^5}{120}f^v(x_0) + \frac{64h^6}{720}f^{vi}(x_0)$$

If we subtract second from the first:

$$f(x_0 + 2h) - f(x_0 - 2h) = 4hf'(x_0) + \frac{16h^3}{6}f'''(x_0) + \frac{64h^5}{120}f^v(x_0)$$

$$2)f'(x_0) = \frac{f(x_0 + 2h) - f(x_0 - 2h)}{4h} - \frac{4h^2}{6}f'''(x_0) - \frac{16h^4}{120}f^v(x_0)$$

If we subtract 4 times of first equation from the second one to cancel out third order derivative.

$$-3f'(x_0) = \frac{f(x_0 + 2h) - f(x_0 - 2h)}{4h} - \frac{4f(x_0 + h) - 4f(x_0 - h)}{2h} - \frac{12h^4}{120}f^v(x_0)$$

$$f'(x_0) = \frac{1}{12h} \left( f(x_0 - 2h) - 8f(x_0 - h) + 8f(x_0 + h) - f(x_0 + 2h) \right) + \frac{h^4}{30}f^v(x_0)$$

Truncation error which is fourth order:

$$\varepsilon(h) = \frac{h^4}{30}f^v(x_0)$$

### 2.0.3 Sixth Order Centered Difference Formula

We are trying to find derivative of  $x = x_0$  by using  $x = x_0 + h, x = x_0 - h, x = x_0 + 2h, x = x_0 + 3h, x = x_0 - 3h$ . By Taylor Formula:

For  $x = x_0 + h, x = x_0 - h$ :

$$\begin{aligned} f(x_0 + h) &= f(x_0) + hf'(x_0) + \frac{h^2}{2}f''(x_0) + \frac{h^3}{6}f'''(x_0) + \frac{h^4}{24}f^{iv}(x_0) + \frac{h^5}{120}f^v(x_0) \\ &\quad + \frac{h^6}{720}f^{vi}(x_0) + \frac{h^7}{5040}f^{vii}(x_0) + \frac{h^8}{40320}f^{viii}(x_0) \\ f(x_0 - h) &= f(x_0) - hf'(x_0) + \frac{h^2}{2}f''(x_0) - \frac{h^3}{6}f'''(x_0) + \frac{h^4}{24}f^{iv}(x_0) - \frac{h^5}{120}f^v(x_0) \\ &\quad + \frac{h^6}{720}f^{vi}(x_0) - \frac{h^7}{5040}f^{vii}(x_0) + \frac{h^8}{40320}f^{viii}(x_0) \end{aligned}$$

If we subtract second from the first:

$$f(x_0 + h) - f(x_0 - h) = 2hf'(x_0) + \frac{h^3}{3}f'''(x_0) + \frac{h^5}{60}f^v(x_0) + \frac{h^7}{2520}f^{vii}(x_0)$$

$$1)f'(x_0) = \frac{f(x_0 + h) - f(x_0 - h)}{2h} - \frac{h^2}{6}f'''(x_0) - \frac{h^4}{120}f^v(x_0) - \frac{h^6}{5040}f^{vii}(x_0)$$

For  $x = x_0 + 2h, x = x_0 - 2h$ :

$$\begin{aligned} f(x_0 + 2h) &= f(x_0) + 2hf'(x_0) + \frac{4h^2}{2}f''(x_0) + \frac{8h^3}{6}f'''(x_0) + \frac{16h^4}{24}f^{iv}(x_0) + \frac{32h^5}{120}f^v(x_0) \\ &\quad + \frac{64h^6}{720}f^{vi}(x_0) + \frac{128h^7}{5040}f^{vii}(x_0) + \frac{256h^8}{40320}f^{viii}(x_0) \\ f(x_0 - 2h) &= f(x_0) - 2hf'(x_0) + \frac{4h^2}{2}f''(x_0) - \frac{8h^3}{6}f'''(x_0) + \frac{16h^4}{24}f^{iv}(x_0) - \frac{32h^5}{120}f^v(x_0) \\ &\quad + \frac{64h^6}{720}f^{vi}(x_0) - \frac{128h^7}{5040}f^{vii}(x_0) + \frac{256h^8}{40320}f^{viii}(x_0) \end{aligned}$$

If we subtract second from the first:

$$f(x_0 + 2h) - f(x_0 - 2h) = 4hf'(x_0) + \frac{8h^3}{3}f'''(x_0) + \frac{32h^5}{60}f^v(x_0) + \frac{128h^7}{2520}f^{vii}(x_0)$$

$$2)f'(x_0) = \frac{f(x_0 + 2h) - f(x_0 - 2h)}{4h} - \frac{2h^2}{3}f'''(x_0) - \frac{8h^4}{120}f^v(x_0) - \frac{32h^6}{2520}f^{vii}(x_0)$$

For  $x = x_0 + 3h$ ,  $x = x_0 - 3h$ : For  $x = x_0 + h$ ,  $x = x_0 - h$ :

$$f(x_0 + 3h) = f(x_0) + 3hf'(x_0) + \frac{9h^2}{2}f''(x_0) + \frac{27h^3}{6}f'''(x_0) + \frac{81h^4}{24}f^{iv}(x_0) + \frac{243h^5}{120}f^v(x_0) + \frac{729h^6}{720}f^{vi}(x_0) + \frac{2187h^7}{5040}f^{vii}(x_0) + \frac{6561h^8}{40320}f^{viii}(x_0)$$

$$f(x_0 - 3h) = f(x_0) - 3hf'(x_0) + \frac{9h^2}{2}f''(x_0) - \frac{27h^3}{6}f'''(x_0) + \frac{81h^4}{24}f^{iv}(x_0) - \frac{243h^5}{120}f^v(x_0) + \frac{729h^6}{720}f^{vi}(x_0) - \frac{2187h^7}{5040}f^{vii}(x_0) + \frac{6561h^8}{40320}f^{viii}(x_0)$$

If we subtract second from the first:

$$f(x_0 + 3h) - f(x_0 - 3h) = 6hf'(x_0) + \frac{27h^3}{3}f'''(x_0) + \frac{243h^5}{60}f^v(x_0) + \frac{2187h^7}{2520}f^{vii}(x_0)$$

$$3)f'(x_0) = \frac{f(x_0 + 3h) - f(x_0 - 3h)}{6h} - \frac{27h^2}{18}f'''(x_0) - \frac{243h^4}{360}f^v(x_0) - \frac{2187h^6}{15120}f^{vii}(x_0)$$

We have found three centered second order approximations. These are given below. I will increase the order by doing algebraic operations.

$$1)f'(x_0) = \frac{f(x_0 + h) - f(x_0 - h)}{2h} - \frac{h^2}{6}f'''(x_0) - \frac{h^4}{120}f^v(x_0) - \frac{h^6}{5040}f^{vii}(x_0)$$

$$2)f'(x_0) = \frac{f(x_0 + 2h) - f(x_0 - 2h)}{4h} - \frac{2h^2}{3}f'''(x_0) - \frac{8h^4}{120}f^v(x_0) - \frac{32h^6}{2520}f^{vii}(x_0)$$

$$3)f'(x_0) = \frac{f(x_0 + 3h) - f(x_0 - 3h)}{6h} - \frac{27h^2}{18}f'''(x_0) - \frac{243h^4}{360}f^v(x_0) - \frac{2187h^6}{15120}f^{vii}(x_0)$$

If I subtract 4 times equation 1 from equation 2:

$$a)-3f'(x_0) = \frac{3f(x_0 + 2h) - 3f(x_0 - 2h) + 24f(x_0 + h) - 24f(x_0 - h)}{12h} - \frac{36h^4}{360}f^v(x_0) - \frac{180h^6}{15120}f^{vii}(x_0)$$

If I subtract 9 times equation 1 from equation 3:

$$b)-8f'(x_0) = \frac{2f(x_0 + 3h) - 2f(x_0 - 3h) - 54f(x_0 + h) + 54f(x_0 - h)}{12h} - \frac{216h^4}{360}f^v(x_0) - \frac{2160h^6}{15120}f^{vii}(x_0)$$

We have found two centered fourth order approximations. Then, we will cancel fourth order and we will get sixth order approximation. If I subtract 6 times equation a from equation b:

$$10f'(x_0) = \frac{2(f(x_0+3h)-f(x_0-3h)) + 18(f(x_0-2h)-f(x_0+2h)) + 90(f(x_0+h)-f(x_0-h))}{12h} - \frac{1080h^6}{15120}f^{vii}(x_0)$$

$$f'(x_0) = \frac{2(f(x_0+3h)-f(x_0-3h)) + 18(f(x_0-2h)-f(x_0+2h)) + 90(f(x_0+h)-f(x_0-h))}{120h} - \frac{108h^6}{15120}f^{vii}(x_0)$$

Truncation error which is sixth order:

$$\varepsilon(h) = \frac{108h^6}{15120}f^{vii}(x_0)$$

## 2.1 Python Implementation

Listing 1: Python Implementation of Derivation Algorithm

```
1 import math
2 import matplotlib.pyplot as plt
3 import numpy as np
4 import sympy as sym
5 from sympy import Symbol, Derivative
6 #Function which returns theoretical and real absolute error
7 def errorComparasion(f,diff,x0,k,order,h):
8     #Finds approximate value of f' using centered difference formula
9     res= derivative(f,x0,k,order,h)
10    #Calculates h value
11    h = math.pow(h,-k)
12    #Calculates absolute error
13    absolute_error=abs(diff(1,x0)-res)
14    #Finds corresponding theoretical error values
15    if order==2:
16        theoretical = (h**2/6)*diff(3,x0)
17    elif order==4:
18        theoretical = (h**4/30)*diff(5,x0)
19    elif order==6:
20        theoretical = (h**6/15120)*diff(7,x0)
21    #Returns absolute and theoretical errors
22    return absolute_error, theoretical
23 #Function which calculates derivatives with using centered difference
24 def derivative(f,x0,k,order,h):
25     #Calculates h value
26     h = math.pow(h,-k)
27     #Returns corresponding centered difference formulas for the first ↵
28     derivative
29     if order==2:#Returns corresponding
30         res= (f(x0+h)-f(x0-h))/(2*h)
31         return res
32     elif order==4:
33         res = (1/(12*h))*( f(x0-2*h)-8*f(x0-h)+8*f(x0+h)-f(x0+2*h))
34         return res
35     elif order==6:
36         res =(2*( f(x0+3*h)-f(x0-3*h))+18*(f(x0-2*h)-f(x0+2*h))+90*(f(↵
37             x0+h)-f(x0-h)))/(120*h)
38         return res
39     else:
40         #Prints invalid input error
41         print("Order can be only 2, 4 or 6")
42         return
43 def f(x):
```

```

42     return math.exp(x)*math.cos(2*x)
43 #Function which returns higher order derivatives of f function7
44 def diff(order,x0):
45     x = sym.Symbol('x')
46     function= sym.exp(x)*sym.cos(2*x)
47     for i in range(order):
48         function = Derivative(function, x)
49         function.doit()
50     return sym.N(function.subs(x, x0))
51 #Function which prints theoretical and real absolute errors of ↵
    calculations
52 def Compare(f,diff,h,k_0,k_1, x_0):
53     for order in [2,4,6]:
54         print("Order of "+str(order)+":")
55         for k in range (k_0,k_1+1):
56             print("Step size: "+str(h)+"^"+str(k))
57             absolute, theoretical=errorComparasion(f,diff,x_0,k,order,↵
                h)
58             print("Absolute error:" +str(absolute)+" | "+"Theoretical ↵
                error:" +str(theoretical))
59 def Graph(f,diff,h,k_0,k_1, x_0):
60     second=[] #list for second order approximations
61     fourth=[] #list for fourth order approximations
62     sixth=[] #list for sixth order approximations
63     h_array=[] #list for keeping h values
64     #Fills h array with negative powers of 10 up to -10th power
65     for k in range(k_0,k_1+1):
66         h1 = math.pow(h,-k)
67         h_array.append(h1)
68     #Fills lists with approximations for every power of 10
69     for order in [2,4,6]:
70         for k in range (k_0,k_1+1):
71             absolute_error, theoretical =errorComparasion(f,diff,x_0,k↵
                ,order,h)
72             if(order==2):
73                 second.append(absolute_error)
74             if(order==4):
75                 fourth.append(absolute_error)
76             if(order==6):
77                 sixth.append(absolute_error)
78     plt.figure()
79     plt.xlabel("h")
80     plt.ylabel("Absolute Error")
81     plt.loglog( h_array,second,"--",label="second order")
82     plt.loglog(h_array, fourth,"--",label="fourth order")
83     plt.loglog( h_array,sixth,"--",label="sixth order")
84     plt.legend()

```

85

86 `Graph(f,diff,10,1,10, 1)`

87 `Compare(f,diff,10,1,10, 1)`

---




Figure 1: Errors-h & Order Relations of  $e^x \cos(2x)$

Observations show that higher order formulas are more reliable and they gives more closer results. In the graphic, we can observe that sixth order approximations are in below. But for lower h values, absolute error is getting bigger. Because of round-off errors, computer cannot perform operations smoothly.

In the table which is given in next page, for smaller k values our the theoretical error is very close to real absolute error. But increase in k values causes larger mismatches between theory and real results.

In essence, differentiation operations with using Taylor series give us accurate results, if we choose optimal h values. In my opinion, it is simple and reliable technique.

Absolute Error- Theoretical Findings			
Order	k	Absolute Error	Theoretical Error
2	1	0.0290173150437871	0.0289778359422243
4		0.000155637768665784	0.000158487446007179
6		5.09898860734381e-6	-4.76154062133788e-8
2	2	0.000289782321460486	0.000289778359422243
4		1.58458925980653e-8	1.58487446007179e-8
6		5.14965847742133e-12	-4.76154062133788e-14
2	3	2.89778441153743e-6	2.89778359422243e-6
4		1.04449782156735e-12	1.58487446007179e-12
6		6.35047570085590e-13	-4.76154062133788e-20
2	4	2.89773307571295e-8	2.89778359422243e-8
4		1.15463194561016e-12	1.58487446007179e-16
6		8.59756710269721e-13	-4.76154062133788e-26
2	5	2.71405120599866e-10	2.89778359422243e-10
4		2.65050204006911e-11	1.58487446007179e-20
6		3.05755420981768e-11	-4.76154062133788e-32
2	6	1.49279699712679e-10	2.89778359422243e-12
4		1.67784008908711e-10	1.58487446007179e-24
6		2.08491002240407e-10	-4.76154062133788e-38
2	7	1.29401023230002e-9	2.89778359422243e-14
4		2.58927101981499e-9	1.58487446007179e-28
6		3.07036707170028e-9	-4.76154062133788e-44
2	8	6.47755094007607e-9	2.89778359422243e-16
4		1.57294106628569e-8	1.58487446007179e-32
6		1.49892604994761e-8	-4.76154062133788e-50
2	9	1.60055902753697e-7	2.89778359422243e-18
4		2.15567053984955e-7	1.58487446007179e-36
6		2.48873743835532e-7	-4.76154062133788e-56
2	10	2.84033307096365e-7	2.89778359422243e-20
4		6.54107648045965e-7	1.58487446007179e-40
6		8.76152253859175e-7	-4.76154062133788e-62