

# Extensive Study of the Wobbling Properties in $^{163}\text{Lu}$ for the Positive and Negative Parity States

Robert Poenaru <sup>\*1,2</sup> and Apolodor Aristotel Raduta <sup>†2,3</sup>

<sup>1</sup>Doctoral School of Physics, University of Bucharest, Romania

<sup>2</sup>*Horia Hulubei* National Institute for Physics and Nuclear Engineering,  
Măgurele-Bucharest, Romania

<sup>3</sup>Academy of Romanian Scientists, Bucharest, Romania

January 25, 2021

## Abstract

A new interpretation on the wobbling structure in  $^{163}\text{Lu}$  is developed, based on the concept of parity symmetry. It is known that four wobbling bands are experimentally observed in this isotope, where three of them are considered as wobbling phonon excitations (namely  $TSD_2$ ,  $TSD_3$ , and  $TSD_4$ ) and the yrast band for the ground state (that is  $TSD_1$ ). In the present work, the trial function that is used for obtaining the wobbling spectrum is analyzed in terms of its behavior under the rotation operation. Indeed, due to a specific symmetry to rotations with  $\pi$  around the 2-axis of the triaxial system, the parity becomes a good quantum number. As such, the trial function admits solutions with negative parity, which belong to the rotational states in  $TSD_4$ . A unified description of all the triaxial super-deformed bands in  $^{163}\text{Lu}$  is achieved with the new formalism.

## 1 Introduction

Triaxiality in nuclei has become an interesting topic for physicists over the years, mainly due to its great challenge of measure it experimentally, but also for its large number of characteristics that are said to be resulting from these kind of shapes. Moreover, stable triaxial shapes are of rare occurrence across the chart of nuclides [1], since the predominant character of nuclei is either spherical or axially symmetric. Over the last two decades, it has been shown that triaxiality plays a crucial role in measurements of important quantities like proton emission probabilities [2], separation energies of the nucleons [1], and also fission barriers in heavy nuclei [3], however, concrete evidences

---

<sup>\*</sup>E-mail: robert.poenaru@drd.unibuc.ro

<sup>†</sup>E-mail: raduta@nipne.ro

of triaxiality in nuclei were still missing or under investigation. A tremendous work was given in finding a clear signature for non-axially symmetric shapes: effects such as anomalous signature splitting [4], signature inversion [5], and staggering of  $\gamma$  bands [6] were pointed, but only recently two clear fingerprints of nuclear triaxiality have emerged in the literature, based on both experimental and theoretical findings. Indeed, the phenomena of *chiral symmetry breaking* and that of *wobbling motion* (W.M.) are considered as unique characteristics of nuclear triaxiality.

Chirality consists in the existence of a pair of chiral twin bands with an identical structure and almost similar energies. These bands are expected to appear due to the coupling of valence nucleons and the collective mode of rotation that could drive the total spin away from any of the three principal planes, giving rise to both left-handed and right-handed orientation of the angular momentum vectors [7]. A rigorous investigation of all the nuclei with chiral bands is given by Xiong and Wang in [8], where reportedly a total of 59 chiral doublet bands in 47 such nuclei are confirmed. As a matter of fact, 8 of these nuclei have multiple chiral doublets.

On the other hand, the experimental observations regarding wobbling motion have been quite rare, even though this kind of collective motion has been theoretically predicted almost 50 years ago by Bohr and Mottelson [9] when they were investigating the rotational modes of a triaxial nucleus by means of a Triaxial Rotor Model (TRM). Therein, they showed that for a triaxial rotor, the main rotational motion is around the axis with the largest moment of inertia (MOI), as it is energetically the most favorable. This mode is quantum mechanically disturbed by the rotation around the other two axes, since rotation around any of the three principal axes of the system are possible, due to the anisotropy of the three different MOIs (that is  $\mathcal{I}_1 \neq \mathcal{I}_2 \neq \mathcal{I}_3$ ).

W.M. can be viewed as the quantum analogue to the motion of the asymmetric top, whose rotation around the axis with largest MOI is energetically favored and stable. A uniform rotation about this axis will have the lowest energy for a given angular momentum (spin). As the energy increases, this axis will start to precess with a harmonic type of oscillation about the space-fixed angular momentum vector, giving rise to a family of wobbling bands, each characterized by a wobbling phonon number  $n_w$ . The resulting quantal spectrum will be a sequence of rotational  $\Delta I = 2$  bands, with alternating signature number for each wobbling excitation. According to [9], it is possible to obtain the wobbling spectrum of any triaxial rigid rotor, by using the information related to its angular momentum  $I$ , moments of inertia  $\mathcal{I}_{1,2,3}$ , rotational frequency  $\omega_{\text{rot}}$ , wobbling frequency  $\omega_{\text{wob}}$  as follows:

$$E_{\text{rot}} = \sum_i \left( \frac{\hbar^2}{2\mathcal{I}_i} \right) I_i^2 \approx \frac{\hbar^2}{2\mathcal{I}_1} I(I+1) + \hbar\omega_{\text{wob}} \left( n_w + \frac{1}{2} \right), \quad (1)$$

with  $\omega_{\text{wob}}$  given by the following expression:

$$\hbar\omega_{\text{wob}} = \hbar\omega_{\text{rot}} \sqrt{\frac{(\mathcal{I}_1 - \mathcal{I}_2)(\mathcal{I}_1 - \mathcal{I}_3)}{\mathcal{I}_2\mathcal{I}_3}}. \quad (2)$$




Figure 1: Family of wobbling bands for a simple triaxial rotor (left-side). Tilting of the angular momentum vector away from the rotational axis (right-side). This schematic representation was done for an arbitrary set of MOIs  $\mathcal{I}_1 : \mathcal{I}_2 : \mathcal{I}_3 = 25 : 5 : 2$ .

where the rotational frequency of the rigid rotor is given by  $\hbar\omega_{\text{rot}} = \frac{\hbar I^2}{\mathcal{I}_1}$ . In Eq. 1, the approximation of very large MOI along 1-axis is considered (i.e.,  $\mathcal{I}_1 \gg \mathcal{I}_2, \mathcal{I}_3$ ), and  $I(I+1) = I_1^2 + I_2^2 + I_3^2$ . One can see that the wobbling motion is expressed as a 1-dimensional vibration with only one variable, since the energy of the zero-point fluctuation is  $\frac{\hbar\omega_{\text{wob}}}{2}$  [10].

Just for an illustrative purpose, Figure 1 shows a theoretical spectrum for the wobbling bands within a triaxial rigid rotor. The family of wobbling bands are obtained from a set of three moments of inertia (along the three principal axes), a given angular momentum, and different wobbling phonon numbers. Moreover, in Figure 1, the tilting of the angular momentum away from the rotational axis is sketched, where the tilt increases with the increase in the wobbling excitation. In a given sequence of wobbling bands, both the intra-band  $\Delta I = 2$  as well as inter-band  $\Delta I = 1$  transitions have a strong  $E2$  collective character.

It is important to mention that the wobbling spectrum described by Eq. 1 and graphically represented in Figure 1 was firstly predicted for an even-even triaxial nucleus [9]. This predicted wobbling mode has not been experimentally confirmed yet. However, the first experimental evidence for wobbling excitations in nuclei was for an even-odd nucleus, namely  $^{163}\text{Lu}$ , where a single one-phonon wobbling band was measured initially [11], followed by two additional wobbling bands discovered one year later [12, 13].

After the first discovery of wobbling bands in  $^{163}\text{Lu}$  ( $Z = 71$ ), an entire series of even-odd isotopes with  $A \approx 160$  were experimentally confirmed as *wobblers*:  $^{161}\text{Lu}$ ,  $^{165}\text{Lu}$ ,  $^{167}\text{Lu}$ , and  $^{167}\text{Ta}$ . In these nuclei, the wobbling mode appears due to the coupling of a valence nucleon (the so-called  $\pi(i_{13/2})$  intruder) to a triaxial core, driving the entire nuclear system up to large deformation ( $\epsilon \approx 0.4$ ) [14].

With time, several nuclei in which WM occurs were also reported in regions of smaller  $A$ . Indeed, two isotopes with  $A \approx 130$ :  $^{133}\text{La}$  [15] and  $^{135}\text{Pr}$  [16, 17] were identified as having wobbling bands, which emerged from the coupling with a triaxial even-even core of another intruder (the  $\pi(h_{11/2})$  nucleon) for  $^{135}\text{Pr}$ , and an additional pair of positive parity quasi-protons which are making an alignment with the short axis of the triaxial rotor for  $^{105}\text{Pr}$ . The resulting coupling in both cases have a deformation  $\epsilon = 0.16$  [15, 16], which is obviously smaller than the deformation in the heavier nuclei within the  $A \approx 160$  region. A third nucleus that also lies in this mass region was confirmed very recently by Chakraborty et. al. in [18], namely the odd- $A$   $^{127}\text{Xe}$ , where a total of four wobbling bands have been reported by the team (two yrast bands, and two excited phonon bands with  $n_w = 1$  and  $n_w = 2$ ).

Some additional progress towards a more comprehensive wobbling spectroscopy was made in the  $A \approx 100$  mass region, with an experimental evidence for  $^{105}\text{Pd}$  that showed of two such bands that are built on a  $\nu(h_{11/2})$  configuration, the first one so far in which a valence neutron couples to the triaxial core [19]. The resulting configuration drives the nuclear system up to deformation  $\epsilon \approx 0.26$ .

The heaviest nuclei known so far in which WM has been experimentally observed are the isotopes  $Z = 79$  with  $A = 183$  [20] and  $A = 187$  [21], respectively. However, for the case of  $^{187}\text{Au}$ , there is an ongoing debate [22] whether the two wobbling bands ( $n_w = 0$  and  $n_w = 1$ ) are bands with wobbling character, or if they are of magnetic nature (which would exclude the wobbling phonon interpretation).

Regarding the wobbling motion for the even-even nuclei (behavior that was described above through the schematic representation from Figure 1), the experimental results are very fragmentary, with unclear evidence on such collective behavior in nuclei. However, some embryos of even-even wobblers have been reported in the recent years. For example, the  $^{112}\text{Ru}$  ( $Z = 44$ ) nucleus has three wobbling bands [23], with two of them being excited (one- and two-phonon wobbling bands). Another example is the even-even  $^{130}\text{Ba}$  ( $Z = 56$ ) [24–26]. Indeed, for  $^{112}\text{Ru}$ , the ground band together with the odd and even spin members of the  $\gamma$  band with were interpreted as zero-(yrast), one-, and two-phonon wobbling bands. Unfortunately, since there are no data concerning the electromagnetic transitions, its wobbling character is still unclear. On the other hand, for the nucleus  $^{130}\text{Ba}$ , from its recent study regarding the band structure [24], a pair of bands with even and odd spins were proposed as zero- and one-phonon wobbling bands, respectively. What it is worth noting for this case is the fact that these two bands are built on a configuration in which two aligned protons that emerge from the bottom of  $h_{j=11/2}$  shell couple with the triaxial core. One remarks the change in nature of the wobbling motion from a purely collective form, but in the presence of two aligned quasiparticles [25].

Regarding the interpretation of the wobbling motion which occurs in the nuclei that were mentioned above, it is mandatory to discuss some aspects related to its behavior with the increase in total angular momentum (nuclear spin). It is a long lasting debate on whether certain nuclei behave as *longitudinal wobblers* (LW) or *transverse wobblers* (TW). The concepts of LW and TW emerged from an extensive study done by Frauendorf et. al. [27] in which the team discussed the possible coupling schemes that a valence

nucleon can create with the triaxial core, thus giving rise to two possible scenarios. Based on microscopic calculations using the Quasiparticle Triaxial Rotor (QTR) model, they showed that if the odd valance nucleon aligns its angular momentum vector  $\vec{j}$  with the axis of largest MOI, the nuclear system is of longitudinal wobbling character. On the other hand, if the odd nucleon aligns its a.m. vector  $\vec{j}$  with an axis perpendicular to the one with largest MOI, then the nuclear system has a transverse wobbling character. From the microscopic calculations, it was shown that for LW, the wobbling energy  $E_{\text{wob}}$  (see Eq. 3) has an *increasing* behavior with an increase in spin, while for TW the energy  $E_{\text{wob}}$  *decreases* with spin.

Within the nuclei that were mentioned above, most of them are of TW type, with only  $^{133}\text{La}$  [15],  $^{127}\text{Xe}$  [18], and  $^{183,187}\text{Au}$  [20, 21] being nuclei with LW character. The energy that characterizes the type of wobbling in a nuclear system is the energy of the first excited band (the one-phonon  $n_w = 1$  wobbling band) relative to the yrast ground band (zero-phonon  $n_w = 0$  wobbling band):

$$E_{\text{wob}} = E_1(I) - \left( \frac{E_0(I+1) + E_0(I-1)}{2} \right), \quad (3)$$

with 0 and 1 representing the wobbling phonon number  $n_w$ .

The odd nucleons that couple with the rigid triaxial core will influence the appearance of a particular wobbling regime (LW or TW). In all the wobblers, there is a proton from a certain orbital which is coupling with the core, except for the case of  $^{105}\text{Pd}$ , where the valance nucleon is a neutron. The nature of the odd quasiparticle (i.e., particle or hole) and its "position" in the deformed  $j$ -shell (i.e. bottom or top) will determine whether its angular momentum  $\vec{j}$  will align with the *short* ( $s$ ) or *long* ( $l$ ) axes of the triaxial rotor, respectively (with the notations short  $s$ , long  $l$ , and medium  $m$  axes of a triaxial ellipsoid). The reasoning behind this has to do with the minimization of the overall energy of the system: in the first case, a maximal overlap of its density distribution with the triaxial core will determine a minimal energy, while in the second case, a minimal overlap of the density distribution of the particle with the core will result in a minimal energy. Moreover, if the quasiparticle emerges from the middle of the  $j$ -shell, then it tends to align its angular momentum vector  $\vec{j}$  with the *medium* ( $m$ ) axis of the triaxial core. Figure 2 aims at depicting the type of alignment of a quasiparticle with the triaxial core.

As previously mentioned, for a given angular momentum, uniform rotation around the axis with the largest MOI corresponds to a minimum energy. For a triaxial rotor, this is equivalent to rotation around the  $m$  axis. Therefore, Frauendorf [27] classified the LW as the situation when the odd nucleon will align its angular momentum along the  $m$ -axis, while TW being the situation where  $j$  is aligned perpendicular to the  $m$ -axis (with  $s$ - or  $l$ -axis alignment depending on the  $j$ -shell orbital from which the odd nucleon arises). It is worthwhile to mention the fact that the analysis done in Ref. [27] was within a so-called *Frozen Alignment* approximation, where the angular momentum of the odd particle  $\vec{j}$  is rigidly aligned with one of the three principal axes of the triaxial ellipsoid (that is  $s$ -,  $l$ - or  $m$ -axis).




Figure 2: The wobbling regime (LW or TW) based on the type of alignment for an odd quasiparticle with the triaxial core. Figure based on the quantal analysis done in [27].

For a better understanding of the wobbling regimes in terms of angular momentum alignment, the schematic illustration from Figure 3 depicts a simple wobblers (the case firstly developed by Bohr and Mottelson [9]) - inset A.0, a longitudinal wobblers - inset A.1, and lastly a transverse wobblers - inset A.1.

## 2 Theoretical Background

The Hamiltonian of the system is given in terms of a term that corresponds to the core deformation, and a second term which corresponds to the valence nucleon moving in a mean-field with quadrupole character (generated by the triaxial core).

$$\hat{H} = \hat{H}_{\text{rot}} + \hat{H}_{\text{s.p.}} \quad (4)$$

## References

- [1] Peter Möller, Ragnar Bengtsson, B Gillis Carlsson, Peter Olivius, and Takatoshi Ichikawa. Global calculations of ground-state axial shape asymmetry of nuclei. *Physical review letters*, 97(16):162502, 2006.
- [2] DS Delion, RJ Liotta, and Ramon Wyss. Theories of proton emission. *Physics reports*, 424(3):113–174, 2006.




Figure 3: The geometry for the angular momentum for a simple wobbler: A.0, a longitudinal wobbler: A.1, and a transverse wobbler: A.2.

- [3] Peter Möller, Arnold J Sierk, Takatoshi Ichikawa, Akira Iwamoto, Ragnar Bengtsson, Henrik Uhrenholt, and Sven Åberg. Heavy-element fission barriers. *Physical Review C*, 79(6):064304, 2009.
- [4] I Hamamoto and H Sagawa. Triaxial deformation in odd- $z$  light rare-earth nuclei. *Physics Letters B*, 201(4):415–419, 1988.
- [5] R Bengtsson, H Frisk, FR May, and JA Pinston. Signature inversion—a fingerprint of triaxiality. *Nuclear Physics A*, 415(2):189–214, 1984.
- [6] J Stachel, N Kaffrell, E Grosse, H Emling, H Folger, R Kulessa, and D Schwalm. Triaxiality and its dynamics in 104ru investigated by multiple coulomb excitation. *Nuclear Physics A*, 383(3):429–467, 1982.
- [7] Stefan Frauendorf and Jie Meng. Tilted rotation of triaxial nuclei. *Nuclear Physics A*, 617(2):131–147, 1997.
- [8] BW Xiong and YY Wang. Nuclear chiral doublet bands data tables. *Atomic Data and Nuclear Data Tables*, 125:193–225, 2019.
- [9] Aage Bohr and Ben R Mottelson. *Nuclear structure*, volume 1. World Scientific, 1998.
- [10] Gudrun B Hagemann and Ikuko Hamamoto. Quantized wobbling in nuclei. *Nuclear Physics News*, 13(3):20–24, 2003.
- [11] SW Ødegård, GB Hagemann, DR Jensen, M Bergström, B Herskind, G Sletten, S Törmänen, JN Wilson, PO Tjøm, I Hamamoto, et al. Evidence for the wobbling mode in nuclei. *Physical review letters*, 86(26):5866, 2001.
- [12] DR Jensen, GB Hagemann, I Hamamoto, SW Ødegård, B Herskind, G Sletten, JN Wilson, K Spohr, H Hübel, P Bringel, et al. Evidence for second-phonon nuclear wobbling. *Physical review letters*, 89(14):142503, 2002.

- [13] D Ringkøbing Jensen, GB Hagemann, I Hamamoto, SW Ødegård, M Bergström, B Herskind, G Sletten, S Törmänen, JN Wilson, PO Tjøm, et al. Wobbling phonon excitations, coexisting with normal deformed structures in 163lu. *Nuclear Physics A*, 703(1-2):3–44, 2002.
- [14] H Schnack-Petersen, Ragnar Bengtsson, RA Bark, P Bosetti, A Brockstedt, H Carlsson, LP Ekström, GB Hagemann, B Herskind, F Ingebreetsen, et al. Superdeformed triaxial bands in 163,165 lu. *Nuclear Physics A*, 594(2):175–202, 1995.
- [15] S Biswas, R Palit, S Frauendorf, U Garg, W Li, GH Bhat, JA Sheikh, J Sethi, S Saha, Purnima Singh, et al. Longitudinal wobbling in 133 la. *The European Physical Journal A*, 55(9):1–7, 2019.
- [16] James Till Matta. Transverse wobbling in 135 pr. In *Exotic Nuclear Excitations: The Transverse Wobbling Mode in 135 Pr*, pages 77–93. Springer, 2017.
- [17] N Sensharma, U Garg, S Zhu, AD Ayangeakaa, S Frauendorf, W Li, GH Bhat, JA Sheikh, MP Carpenter, QB Chen, et al. Two-phonon wobbling in 135pr. *Physics Letters B*, 792:170–174, 2019.
- [18] S Chakraborty, HP Sharma, SS Tiwary, C Majumder, AK Gupta, P Banerjee, S Ganguly, S Rai, S Kumar, A Kumar, et al. Multiphonon longitudinal wobbling in 127xe. *Physics Letters B*, 811:135854, 2020.
- [19] J Timár, QB Chen, B Kruzsicz, D Sohler, I Kuti, SQ Zhang, J Meng, P Joshi, R Wadsworth, K Starosta, et al. Experimental evidence for transverse wobbling in pd 105. *Physical review letters*, 122(6):062501, 2019.
- [20] S Nandi, G Mukherjee, QB Chen, S Frauendorf, R Banik, Soumik Bhattacharya, Shabir Dar, S Bhattacharyya, C Bhattacharya, S Chatterjee, et al. First observation of multiple transverse wobbling bands of different kinds in au 183. *Physical Review Letters*, 125(13):132501, 2020.
- [21] N Sensharma, U Garg, QB Chen, S Frauendorf, DP Burdette, JL Cozzi, KB Howard, S Zhu, MP Carpenter, P Copp, et al. Longitudinal wobbling motion in au 187. *Physical review letters*, 124(5):052501, 2020.
- [22] S Guo, XH Zhou, CM Petrache, EA Lawrie, S Mthembu, YD Fang, HY Wu, HL Wang, HY Meng, GS Li, et al. Risk of misinterpretation of low-spin non-yrast bands as wobbling bands. *arXiv preprint arXiv:2011.14354*, 2020.
- [23] JH Hamilton, SJ Zhu, YX Luo, AV Ramayya, S Frauendorf, JO Rasmussen, JK Hwang, SH Liu, GM Ter-Akopian, AV Daniel, et al. Super deformation to maximum triaxiality in a= 100–112; superdeformation, chiral bands and wobbling motion. *Nuclear Physics A*, 834(1-4):28c–31c, 2010.



- [24] CM Petrache, PM Walker, S Guo, QB Chen, S Frauendorf, YX Liu, RA Wyss, D Mengoni, YH Qiang, A Astier, et al. Diversity of shapes and rotations in the  $\gamma$ -soft 130ba nucleus: First observation of a t-band in the a= 130 mass region. *Physics Letters B*, 795:241–247, 2019.
- [25] YK Wang, FQ Chen, and PW Zhao. Two quasiparticle wobbling in the even-even nucleus 130ba. *Physics Letters B*, 802:135246, 2020.
- [26] QB Chen, S Frauendorf, and CM Petrache. Transverse wobbling in an even-even nucleus. *Physical Review C*, 100(6):061301, 2019.
- [27] S Frauendorf and F Dönau. Transverse wobbling: A collective mode in odd-a triaxial nuclei. *Physical Review C*, 89(1):014322, 2014.