

A Systematic Description of the Wobbling Motion in Odd-Mass Nuclei Within a Semi-Classical Formalism

PhD Candidate

Robert Poenaru^{1,2}

Scientific Supervisor

Prof. Dr. Em. A. A. Raduta²

¹Doctoral School of Physics, UB

²Department of Theoretical Physics, IFIN-HH

A presentation for the degree of Doctor of Philosophy

May 12, 2023


- 1 Aim and Motivation
- 2 Nuclear Shapes
- 3 Triaxiality and Wobbling Motion

Aim

Research Objectives


- Extend the current interpretation of the **nuclear triaxiality** in the context of its unique fingerprint: **Wobbling Motion**
- Adopt a framework that is as close as possible to **classical physics**.
- Provide new formalisms for the phenomena related to **nuclear deformation**.

Objectives exclusive to the thesis

- Give the reader enough context towards a better understanding of the underlying concepts, methods, and results.
-  create a completely *open-source* project.

Motivation

- **Nuclear Triaxiality** has become a *hot topic* within the scientific community.
- Identifying nuclei with triaxial deformations represents a real **experimental** and **theoretical** challenge.



Nuclear Deformation

Nuclear Radius

The **shape** of the nucleus is most generally described in terms of the *nuclear radius*:

$$R(\theta, \varphi) = R_0 \left(1 + \sum_{\lambda=0}^{\infty} \sum_{\mu=-\lambda}^{\lambda} \alpha_{\lambda\mu} Y_{\lambda}^{\mu}(\theta, \varphi) \right) \quad (1)$$

Quadrupole deformations $\lambda = 2$

- **For us:** Most relevant modes are the **quadrupole vibrations** $\lambda = 2$
 \implies *Play a crucial role in the rotational spectra of nuclei:*
- *Bohr, 1969:* Coordinates $\alpha_{2\mu}$ can be reduced to only two *deformation parameters*: β_2 (**eccentricity**) and γ (**triaxiality**).

Axial shapes

Collective coordinates



- Most of the nuclei are either **spherical** or **axially symmetric** in their ground-state (*Budaca, 2018*).
- Moments of inertia: $\mathcal{I}_{1,2,3}$: two are equal, one is different.



Figure: **spherical:** $\beta_2 = 0$ **prolate:** $\beta_2 > 0$ **oblate:** $\beta_2 < 0$. ($\gamma = 0^\circ$).

Non-axial shapes

- The triaxiality parameter $\gamma \neq 0^\circ$: departure from axial symmetry.
- Moments of inertia: $\mathcal{I}_1 \neq \mathcal{I}_2 \neq \mathcal{I}_3$.



Fingerprints of Triaxiality

Evidence

- Currently, there are **only two** well-established phenomena uniquely attributed to triaxial deformation.
 - Wobbling Motion** - WM (*Bohr and Mottelson, 1970s*)
 - Chiral Motion - χ M (*Frauendorf, 1997*)
- These two can be measured/detected experimentally.

Experimental observations

First experimental evidence for **nuclear wobbling motion** in 2001.


Goal

Describe the elusive character of Wobbling Motion in the context of nuclear triaxiality.

Q Probing triaxiality in nuclei

Triaxial nuclei can be observed/obtained in several experiments:

- Nuclear fission: $A \rightarrow B + C$
- Nuclear fusion: $X + Y \rightarrow Z$
- **Fusion-evaporation reactions:** Long-lived + enhanced deformation
 $Beam(N_1, E) + Target(N_2) \longrightarrow N_3^* \rightarrow \dots \rightarrow triaxial(N_4)$



Q Nuclear facilities



Figure: Gammasphere detector, ANL-ATLAS USA. Source: aps.org






Figure: a) IDS detector, CERN. Source: isolde.web.cern.ch b) JUROGAM II, Finland. Source: twitter.com

Wobbling Motion



Wobbling Effect

- The **total angular momentum** of the nucleus **precesses** and **oscillates** around \mathcal{I}_{\max} .

Rotational Energy

Thank you for your attention ♥