
Edge-significant interactions: is the edge signal carried by M (composition) or by Q (proximity coupling)?
matched baseline M z (clipped at 40)
ratio Q = E/M z
leiden: CXCL12→CXCR4 fibroblast→immune
paper: CXCL12→CXCR4 fibroblast→immune
paper: CXCL12→CXCR4 fibroblast→fibroblast
leiden: CXCL12→CXCR4 immune→immune
leiden: CXCL12→CXCR4 fibroblast→epithelial
extra: PTN→SDC4 epithelial→epithelial
paper: CXCL12→CXCR4 immune→fibroblast
paper: CXCL12→CXCR4 endothelial→immune
paper: CXCL12→CXCR4 fibroblast→endothelial
leiden: CXCL12→CXCR4 immune→epithelial
extra: CD86→CTLA4 fibroblast→immune
leiden: CXCL12→CXCR4 fibroblast→fibroblast
paper: CXCL12→CXCR4 immune→immune
paper: CXCL12→CXCR4 endothelial→fibroblast
paper: CXCL12→CXCR4 fibroblast→epithelial
leiden: CXCL12→CXCR4 immune→fibroblast
leiden: CXCL12→CXCR4 epithelial→immune
extra: CD86→CTLA4 fibroblast→fibroblast
extra: CD86→CTLA4 immune→epithelial
−10
0
10
20
30
40
z vs label-permutation null