

Functions and its properties

Let A, B be two nonempty sets.

The **cartesian product** of A and B is

$$A \times B = \{(a, b) | a \in A, b \in B\}$$

Note: $A \times B \neq B \times A$.

Let $A = \{1, 2, 3\}$, $B = \{a, b\}$. Then

$$A \times B = \{(1, a), (2, a), (3, a), (1, b), (2, b), (3, b)\}$$

Definition (Function)

A function f from A to B is an assignment for each element a of A , a unique element b of B . $a \rightsquigarrow b$

A function f from A to B is a subset of $A \times B$ such that **for each** $a \in A$, there **exists unique** $b \in B$ such that $(a, b) \in f$.

Notation: $f : A \rightarrow B$: a function f from A to B
for each $(a, b) \in f$, we write $f(a) = b$.

Examples of function

- ▶ Let $A = \{a, b\}$ and $\{1, 2, 3\}$. Then

$$f = \{(a, 2), (b, 3)\}$$

is a function; whereas

$$g = \{(a, 1), (a, 2), (b, 1)\}$$

is not a function.

- ▶ Let $A = B = \mathbb{R}$ and $f(x) = x^2$. Then $f : \mathbb{R} \rightarrow \mathbb{R}$ is a function. Define $g : \mathbb{R} \rightarrow \mathbb{R}$ as $g(x) := y$, if $y^2 = x$. Then g is not a function from \mathbb{R} to \mathbb{R} .
- ▶ Let A be a non-empty set. The identity function on A , 1_A is defined as $1_A(a) = a$ for all $a \in A$.

Definition

Suppose

$$f : A \rightarrow B$$

$$g : B \rightarrow C$$

are functions. The **composite** of f and g ,
 $g \circ f : A \rightarrow C$ is a function defined as

$$g \circ f(a) := g(f(a)) \text{ for each } a \in A.$$

Let $f : A \rightarrow B$ be a function.

Domain of $f=A$.

Codomain of $f=B$.

Range of $f = \{b \in B | f(a) = b \text{ for some } a \in A\}$

f is said to be **onto/surjective** if $\text{Range of } f = B$.

f is said to be **injective** if $f(a) = f(a')$ then $a = a'$

Define $f, g : \mathbb{R} \rightarrow \mathbb{R}$ as

$$f(x) = x^2,$$

$$g(x) = +\sqrt{|x|}$$

$$g \circ f(x) = |x|$$

What is $f \circ g$?

Theorem

Let $f, g : \mathbb{R} \rightarrow \mathbb{R}$ be two functions. Suppose $g \circ f$ is injective. Then f is injective

Proof.

Let $f(a) = f(a')$.

We need to prove $a = a'$

Hence $g(f(a)) = g(f(a'))$, i.e., $g \circ f(a) = g \circ f(a')$

Since $g \circ f$ is injective, $a = a'$. □

Let $f : A \rightarrow B$ be a function, i.e.,

$$f = \{(a, f(a)) \mid a \in A\}$$

Consider the set

$$g = \{(f(a), a) \mid a \in A\} \subseteq B \times A$$

If g is a function from B to A , then f is said to be **invertible** with $f^{-1} = g$.

Note: $f \circ f^{-1} = 1_B$ and $f^{-1} \circ f = 1_A$

Example: Let $A = \{a, b, c\}$ and $B = \{1, 2, 3\}$.

$$f = \{(a, 2), (b, 1), (c, 2)\}$$

Is f invertible?

Let $f : A \rightarrow B$ be a function, i.e.,

$$f = \{(a, f(a)) \mid a \in A\}$$

Consider the set

$$g = \{(f(a), a) \mid a \in A\} \subseteq B \times A$$

If g is a function from B to A , then f is said to be **invertible** with $f^{-1} = g$.

Note: $f \circ f^{-1} = 1_B$ and $f^{-1} \circ f = 1_A$

Example: Let $A = \{a, b, c\}$ and $B = \{1, 2, 3\}$.

$$f = \{(a, 2), (b, 1), (c, 2)\}$$

Is f invertible? **No.**

Is $g = \{(a, 2), (b, 1), (c, 3)\}$ invertible?

Let $f : A \rightarrow B$ be a function, i.e.,

$$f = \{(a, f(a)) \mid a \in A\}$$

Consider the set

$$g = \{(f(a), a) \mid a \in A\} \subseteq B \times A$$

If g is a function from B to A , then f is said to be **invertible** with $f^{-1} = g$.

Note: $f \circ f^{-1} = 1_B$ and $f^{-1} \circ f = 1_A$

Example: Let $A = \{a, b, c\}$ and $B = \{1, 2, 3\}$.

$$f = \{(a, 2), (b, 1), (c, 2)\}$$

Is f invertible? **No.**

Is $g = \{(a, 2), (b, 1), (c, 3)\}$ invertible? **Yes.**

Theorem (Criteria for invertible functions)

A function $f : A \rightarrow B$ is invertible if and only if f is injective and onto.

Theorem (Criteria for invertible functions)

A function $f : A \rightarrow B$ is invertible if and only if f is injective and onto.

Proof.

Let f be invertible.

Since $f^{-1} : B \rightarrow A$ is a function, Domain of $f^{-1} = B$.

Hence Range of $f = B$, i.e., f is onto.

Theorem (Criteria for invertible functions)

A function $f : A \rightarrow B$ is invertible if and only if f is injective and onto.

Proof.

Let f be invertible.

Since $f^{-1} : B \rightarrow A$ is a function, Domain of $f^{-1} = B$.

Hence Range of $f = B$, i.e., f is onto.

Let $f(a) = f(a')$

Since f^{-1} is a function,

$$f^{-1}(f(a)) = f^{-1}(f(a'))$$

$$\text{i.e., } a = a'$$



Theorem

Let A and B be two nonempty finite sets of same cardinality and $f : A \rightarrow B$ be a function. If f is injective, then f is invertible.

Theorem

Let A and B be two nonempty finite sets of same cardinality and $f : A \rightarrow B$ be a function. If f is injective, then f is invertible.

Proof.

Let $A = \{a_1, \dots, a_n\}$.

Since f is injective, $f(a_1), f(a_2), \dots, f(a_n)$ are all distinct elements of B .

But $|B| = |A| = n$. Hence

$$B = \{f(a_1), f(a_2), \dots, f(a_n)\}, \text{ i.e., } f \text{ is onto.}$$



Theorem

Let A and B be two nonempty finite sets of same cardinality and $f : A \rightarrow B$ be a function. If f is injective, then f is invertible.

Proof.

Let $A = \{a_1, \dots, a_n\}$.

Since f is injective, $f(a_1), f(a_2), \dots, f(a_n)$ are all distinct elements of B .

But $|B| = |A| = n$. Hence

$$B = \{f(a_1), f(a_2), \dots, f(a_n)\}, \text{ i.e., } f \text{ is onto.}$$



Theorem

Let A and B be two nonempty finite sets of same cardinality and $f : A \rightarrow B$ be a function. If f is surjective, then f is invertible.

Application of invertible functions in Cryptography

Let $A = \{a, b, \dots, z\}$ and $f : A \rightarrow A$ be an invertible function. Take a message. Encode the message by replacing each alphabet of it by its image under f .
(In order to decode this encoded message, f must have inverse.)
The recipient decodes the message by applying f^{-1} to each alphabet.

Graph of a function

Definition (Graph of a function)

Let $f : A \rightarrow B$ be a function. The graph of f is the set of ordered pairs

$$\{(a, f(a)) \mid a \in A\}$$

Graph of a function

Definition (Graph of a function)

Let $f : A \rightarrow B$ be a function. The graph of f is the set of ordered pairs

$$\{(a, f(a)) | a \in A\}$$

Question: Display the graph of a function $f : \mathbb{Z} \rightarrow \mathbb{Z}$ defined as $f(n) = n^2$.

Sequences and summations

Sequence is a **ordered list** of elements.

Definition (Sequence)

A sequence of the set S is a function

$$a : A \rightarrow S \quad \text{where } A \subseteq \mathbb{N}$$

We will write it as $(a_n)_{n=1}^{\infty}$, where $a_n = a(n)$, n^{th} term of the sequence.

Sequences and summations

Sequence is a **ordered list** of elements.

Definition (Sequence)

A sequence of the set S is a function

$$a : A \rightarrow S \quad \text{where } A \subseteq \mathbb{N}$$

We will write it as $(a_n)_{n=1}^{\infty}$, where $a_n = a(n)$, n^{th} term of the sequence.

Examples:

1. $a_n = \frac{1}{n}$ sequence of rational numbers.

Sequences and summations

Sequence is a **ordered list** of elements.

Definition (Sequence)

A sequence of the set S is a function

$$a : A \rightarrow S \quad \text{where } A \subseteq \mathbb{N}$$

We will write it as $(a_n)_{n=1}^{\infty}$, where $a_n = a(n)$, n^{th} term of the sequence.

Examples:

1. $a_n = \frac{1}{n}$ sequence of rational numbers.
2. **(Arithmetic progression)** $\{a, a + d, a + 2d, a + 3d, \dots\}$. n^{th} term is $a + (n - 1)d$

Sequences and summations

Sequence is a **ordered list** of elements.

Definition (Sequence)

A sequence of the set S is a function

$$a : A \rightarrow S \quad \text{where } A \subseteq \mathbb{N}$$

We will write it as $(a_n)_{n=1}^{\infty}$, where $a_n = a(n)$, n^{th} term of the sequence.

Examples:

1. $a_n = \frac{1}{n}$ sequence of rational numbers.
2. (**Arithmetic progression**) $\{a, a + d, a + 2d, a + 3d, \dots\}$. n^{th} term is $a + (n - 1)d$
3. (**Geometric progression**) $\{a, ar, ar^2, ar^3, \dots\}$. n^{th} term is $ar^{(n-1)}$

Task: Given first few terms of the sequence, identify the remaining terms:

$\{1, \frac{1}{2}, \frac{1}{3}, \dots\}$ be a sequence. n^{th} term of the sequence is

Task: Given first few terms of the sequence, identify the remaining terms:

$\{1, \frac{1}{2}, \frac{1}{3}, \dots\}$ be a sequence. n^{th} term of the sequence is $\frac{1}{n}$.

$\{6, 2, \frac{2}{3}, \frac{2}{9}, \frac{2}{27}, \dots\}$ be a sequence. n^{th} term of the sequence is

Task: Given first few terms of the sequence, identify the remaining terms:

$\{1, \frac{1}{2}, \frac{1}{3}, \dots\}$ be a sequence. n^{th} term of the sequence is $\frac{1}{n}$.

$\{6, 2, \frac{2}{3}, \frac{2}{9}, \frac{2}{27}, \dots\}$ be a sequence. n^{th} term of the sequence is $\frac{2}{3^{n-2}}$.

$\{2, 6, 18, 54, 162, \dots\}$ be a sequence. Then n^{th} term of the sequence is

Task: Given first few terms of the sequence, identify the remaining terms:

$\{1, \frac{1}{2}, \frac{1}{3}, \dots\}$ be a sequence. n^{th} term of the sequence is $\frac{1}{n}$.

$\{6, 2, \frac{2}{3}, \frac{2}{9}, \frac{2}{27}, \dots\}$ be a sequence. n^{th} term of the sequence is $\frac{2}{3^{n-2}}$.

$\{2, 6, 18, 54, 162, \dots\}$ be a sequence. Then n^{th} term of the sequence is $2 \cdot (3)^{n-1}$.

Let $\{a_n\}_{n=1}^{\infty}$ be a sequence.

$$\sum_{i=m}^n a_i := a_m + a_{m+1} + \cdots + a_n$$

Let $\{a_n\}_{n=1}^{\infty}$ be a sequence.

$$\sum_{i=m}^n a_i := a_m + a_{m+1} + \cdots + a_n$$

$$\sum_{i=m}^n (a_i + b_i) = \sum_{i=m}^n (a_i) + \sum_{i=m}^n (b_i)$$

Let $\{a_n\}_{n=1}^{\infty}$ be a sequence.

$$\sum_{i=m}^n a_i := a_m + a_{m+1} + \cdots + a_n$$

$$\sum_{i=m}^n (a_i + b_i) = \sum_{i=m}^n (a_i) + \sum_{i=m}^n (b_i)$$

$$\sum_{i=m}^n (ca_i) = c \sum_{i=m}^n (a_i), \text{ where } c \text{ is a constant}$$

Let $\{a_n\}_{n=1}^{\infty}$ be a sequence.

$$\sum_{i=m}^n a_i := a_m + a_{m+1} + \cdots + a_n$$

$$\sum_{i=m}^n (a_i + b_i) = \sum_{i=m}^n (a_i) + \sum_{i=m}^n (b_i)$$

$$\sum_{i=m}^n (ca_i) = c \sum_{i=m}^n (a_i), \text{ where } c \text{ is a constant}$$

$$\sum_{i=1}^4 \sum_{j=2}^3 ij = \sum_{i=1}^4 2i + 3i = (2+3) + (4+6) + (6+9) + (8+12) = 50$$

Let $\{a_n\}_{n=1}^{\infty}$ be a sequence.

$$\sum_{i=m}^n a_i := a_m + a_{m+1} + \cdots + a_n$$

$$\sum_{i=m}^n (a_i + b_i) = \sum_{i=m}^n (a_i) + \sum_{i=m}^n (b_i)$$

$$\sum_{i=m}^n (ca_i) = c \sum_{i=m}^n (a_i), \text{ where } c \text{ is a constant}$$

$$\sum_{i=1}^4 \sum_{j=2}^3 ij = \sum_{i=1}^4 2i + 3i = (2+3) + (4+6) + (6+9) + (8+12) = 50$$

$$\sum_{i=1}^4 \sum_{j=2}^3 ij = \sum_{i=1}^4 i \sum_{j=2}^3 j = \sum_{i=1}^4 i(5) = 5(1+2+3+4) = 50$$

Question: Let a, r be real numbers and $r \neq 0, 1$. Find $\sum_{j=0}^n ar^j$.

Question: Let a, r be real numbers and $r \neq 0, 1$. Find $\sum_{j=0}^n ar^j$.

Let $S = \sum_{j=0}^n ar^j$

$$rS = \sum_{j=0}^n ar^{j+1}$$

$$= \sum_{k=1}^{n+1} ar^k$$

$$= \left(\sum_{k=0}^n ar^k \right) + (ar^{n+1} - a)$$

$$rS = S + (ar^{n+1} - a)$$

$$(r - 1)S = (ar^{n+1} - a)$$

$$S = \frac{(ar^{n+1} - a)}{r - 1} \text{ since } r \neq 1$$

Determine whether each of these functions from \mathbf{Z} to \mathbf{Z} is one-to-one.

- a) $f(n) = n - 1$ b) $f(n) = n^2 + 1$
c) $f(n) = n^3$ d) $f(n) = \lceil n/2 \rceil$

Which functions in Exercise 12 are onto?

Determine whether $f: \mathbf{Z} \times \mathbf{Z} \rightarrow \mathbf{Z}$ is onto if

- a) $f(m, n) = 2m - n$.
b) $f(m, n) = m^2 - n^2$.
c) $f(m, n) = m + n + 1$.
d) $f(m, n) = |m| - |n|$.
e) $f(m, n) = m^2 - 4$.

Infinite sets

We learnt that if A is a finite set then $|A| = \text{no. of distinct elements of } A$.

Infinite sets

We learnt that if A is a finite set then $|A| = \text{no. of distinct elements of } A$.

What about cardinality of infinite sets?

Infinite sets

We learnt that if A is a finite set then $|A| = \text{no. of distinct elements of } A$.

What about cardinality of infinite sets?

We saw

If S is a finite set, then every injective function $f : S \rightarrow S$ is surjective.

Infinite sets

We learnt that if A is a finite set then $|A| = \text{no. of distinct elements of } A$.

What about cardinality of infinite sets?

We saw

If S is a finite set, then every injective function $f : S \rightarrow S$ is surjective.

This statement is same as

If there exists an injective function $f : S \rightarrow S$ which is not surjective, then S is not finite(that is, infinite).

Infinite sets

We learnt that if A is a finite set then $|A| = \text{no. of distinct elements of } A$.

What about cardinality of infinite sets?

We saw

If S is a finite set, then every injective function $f : S \rightarrow S$ is surjective.

This statement is same as

If there exists an injective function $f : S \rightarrow S$ which is not surjective, then S is not finite (that is, infinite).

Definition

A set S is said to be infinite if there exists an injective function $f : S \rightarrow S$ which is not surjective.

Infinite sets

We learnt that if A is a finite set then $|A| = \text{no. of distinct elements of } A$.

What about cardinality of infinite sets?

We saw

If S is a finite set, then every injective function $f : S \rightarrow S$ is surjective.

This statement is same as

If there exists an injective function $f : S \rightarrow S$ which is not surjective, then S is not finite (that is, infinite).

Definition

A set S is said to be infinite if there exists an injective function $f : S \rightarrow S$ which is not surjective.

Example: The sets $\mathbb{N}, \mathbb{Q}, \mathbb{R}$ are infinite sets.

Infinite sets

We learnt that if A is a finite set then $|A| = \text{no. of distinct elements of } A$.

What about cardinality of infinite sets?

We saw

If S is a finite set, then every injective function $f : S \rightarrow S$ is surjective.

This statement is same as

If there exists an injective function $f : S \rightarrow S$ which is not surjective, then S is not finite (that is, infinite).

Definition

A set S is said to be infinite if there exists an injective function $f : S \rightarrow S$ which is not surjective.

Example: The sets $\mathbb{N}, \mathbb{Q}, \mathbb{R}$ are infinite sets.

Definition (countable set)

A set S is said to be countable if either it is finite or if there exists a **bijective** (injective and surjective) function $f : \mathbb{N} \rightarrow S$.

Cardinality of infinite sets

We define the cardinality of $\mathbb{N} := \aleph_0$

Cardinality of infinite sets

We define the cardinality of $\mathbb{N} := \aleph_0$

If there exists an injective function $f : A \rightarrow B$, then we say $|A| \leq |B|$.

Cardinality of infinite sets

We define the cardinality of $\mathbb{N} := \aleph_0$

If there exists an injective function $f : A \rightarrow B$, then we say

$$|A| \leq |B|.$$

$|A| = |B|$ if and only if there exists a bijective function $f : A \rightarrow B$

Cardinality of infinite sets

We define the cardinality of $\mathbb{N} := \aleph_0$

If there exists an injective function $f : A \rightarrow B$, then we say

$$|A| \leq |B|.$$

$|A| = |B|$ if and only if there exists a bijective function $f : A \rightarrow B$

Theorem (Schröder-Bernstein-Cantor theorem)

If $f : A \rightarrow B$ and $g : B \rightarrow A$ are injective functions then there exists a bijective function $h : A \rightarrow B$.

Cardinality of infinite sets

We define the cardinality of $\mathbb{N} := \aleph_0$

If there exists an injective function $f : A \rightarrow B$, then we say

$$|A| \leq |B|.$$

$|A| = |B|$ if and only if there exists a bijective function $f : A \rightarrow B$

Theorem (Schröder-Bernstein-Cantor theorem)

If $f : A \rightarrow B$ and $g : B \rightarrow A$ are injective functions then there exists a bijective function $h : A \rightarrow B$.

Exercise: Show that $|[0, 1]| = |(0, 1)|$.

Cardinality of infinite sets

We define the cardinality of $\mathbb{N} := \aleph_0$

If there exists an injective function $f : A \rightarrow B$, then we say

$$|A| \leq |B|.$$

$|A| = |B|$ if and only if there exists a bijective function $f : A \rightarrow B$

Theorem (Schröder-Bernstein-Cantor theorem)

If $f : A \rightarrow B$ and $g : B \rightarrow A$ are injective functions then there exists a bijective function $h : A \rightarrow B$.

Exercise: Show that $|[0, 1]| = |(0, 1)|$.

If the set A is countably infinite, then

$$|A| = \aleph_0$$

Cardinality of infinite sets

We define the cardinality of $\mathbb{N} := \aleph_0$

If there exists an injective function $f : A \rightarrow B$, then we say

$$|A| \leq |B|.$$

$|A| = |B|$ if and only if there exists a bijective function $f : A \rightarrow B$

Theorem (Schröder-Bernstein-Cantor theorem)

If $f : A \rightarrow B$ and $g : B \rightarrow A$ are injective functions then there exists a bijective function $h : A \rightarrow B$.

Exercise: Show that $|[0, 1]| = |(0, 1)|$.

If the set A is countably infinite, then

$$|A| = \aleph_0$$

An infinite set is said to be **uncountable** if it is not countable.

Cardinality of infinite sets

We define the cardinality of $\mathbb{N} := \aleph_0$

If there exists an injective function $f : A \rightarrow B$, then we say

$$|A| \leq |B|.$$

$|A| = |B|$ if and only if there exists a bijective function $f : A \rightarrow B$

Theorem (Schröder-Bernstein-Cantor theorem)

If $f : A \rightarrow B$ and $g : B \rightarrow A$ are injective functions then there exists a bijective function $h : A \rightarrow B$.

Exercise: Show that $|[0, 1]| = |(0, 1)|$.

If the set A is countably infinite, then

$$|A| = \aleph_0$$

An infinite set is said to be **uncountable** if it is not countable.

$\mathbb{N}, \mathbb{Z}, \mathbb{Q}$ are countable.

Theorem

\mathbb{R} is *uncountable*.

Theorem

\mathbb{R} is *uncountable*.

Proof: Suppose not, i.e., \mathbb{R} is countable.

Theorem

\mathbb{R} is *uncountable*.

Proof: Suppose not, i.e., \mathbb{R} is countable.

Then $[0, 1]$ is also countable.

Theorem

\mathbb{R} is *uncountable*.

Proof: Suppose not, i.e., \mathbb{R} is countable.

Then $[0, 1]$ is also countable.

Let $f : \mathbb{N} \rightarrow [0, 1]$ be a bijection.

r_1, r_2, r_3, \dots be all real numbers of $[0, 1]$, where $f(n) = r_n$

Theorem

\mathbb{R} is uncountable.

Proof: Suppose not, i.e., \mathbb{R} is countable.

Then $[0, 1]$ is also countable.

Let $f : \mathbb{N} \rightarrow [0, 1]$ be a bijection.

r_1, r_2, r_3, \dots be all real numbers of $[0, 1]$, where $f(n) = r_n$

Use Cantor's diagonalization argument as follows:

$$r_1 = 0.d_{11}d_{12}d_{13}\dots$$

$$r_2 = 0.d_{21}d_{22}d_{23}\dots$$

$$r_3 = 0.d_{31}d_{32}d_{33}\dots \text{ and so on}$$

Construct $r = 0.d_1d_2d_3d_4\dots$ as follows:

Choose d_i different from d_{ii} for all i

Theorem

\mathbb{R} is uncountable.

Proof: Suppose not, i.e., \mathbb{R} is countable.

Then $[0, 1]$ is also countable.

Let $f : \mathbb{N} \rightarrow [0, 1]$ be a bijection.

r_1, r_2, r_3, \dots be all real numbers of $[0, 1]$, where $f(n) = r_n$

Use Cantor's diagonalization argument as follows:

$$r_1 = 0.d_{11}d_{12}d_{13}\dots$$

$$r_2 = 0.d_{21}d_{22}d_{23}\dots$$

$$r_3 = 0.d_{31}d_{32}d_{33}\dots \text{ and so on}$$

Construct $r = 0.d_1d_2d_3d_4\dots$ as follows:

Choose d_i different from d_{ii} for all i

Then $r \neq r_i$ for any i .



Theorem

\mathbb{R} is uncountable.

Proof: Suppose not, i.e., \mathbb{R} is countable.

Then $[0, 1]$ is also countable.

Let $f : \mathbb{N} \rightarrow [0, 1]$ be a bijection.

r_1, r_2, r_3, \dots be all real numbers of $[0, 1]$, where $f(n) = r_n$

Use Cantor's diagonalization argument as follows:

$$r_1 = 0.d_{11}d_{12}d_{13}\dots$$

$$r_2 = 0.d_{21}d_{22}d_{23}\dots$$

$$r_3 = 0.d_{31}d_{32}d_{33}\dots \text{ and so on}$$

Construct $r = 0.d_1d_2d_3d_4\dots$ as follows:

Choose d_i different from d_{ii} for all i

Then $r \neq r_i$ for any i .



Define $|\mathbb{R}| := c$

If $|S| = n$ then $|P(S)| = 2^n$

If $|S| = n$ then $|P(S)| = 2^n$
What if S is infinite?

If $|S| = n$ then $|P(S)| = 2^n$

What if S is infinite?

Theorem

Let S be a set. Then $|S| < |P(S)|$

If $|S| = n$ then $|P(S)| = 2^n$

What if S is infinite?

Theorem

Let S be a set. Then $|S| < |P(S)|$

Proof.

Define $f : S \rightarrow P(S)$ as

$$f(a) = \{a\}$$

If $|S| = n$ then $|P(S)| = 2^n$

What if S is infinite?

Theorem

Let S be a set. Then $|S| < |P(S)|$

Proof.

Define $f : S \rightarrow P(S)$ as

$$f(a) = \{a\}$$

Clearly, f is an injection. Hence $|S| \leq |P(S)|$

If $|S| = n$ then $|P(S)| = 2^n$

What if S is infinite?

Theorem

Let S be a set. Then $|S| < |P(S)|$

Proof.

Define $f : S \rightarrow P(S)$ as

$$f(a) = \{a\}$$

Clearly, f is an injection. Hence $|S| \leq |P(S)|$

Claim: Any function $g : S \rightarrow P(S)$ is not surjective (hence not bijective).

If $|S| = n$ then $|P(S)| = 2^n$

What if S is infinite?

Theorem

Let S be a set. Then $|S| < |P(S)|$

Proof.

Define $f : S \rightarrow P(S)$ as

$$f(a) = \{a\}$$

Clearly, f is an injection. Hence $|S| \leq |P(S)|$

Claim: Any function $g : S \rightarrow P(S)$ is not surjective (hence not bijective).

$$A = \{x | x \notin g(x)\} \in P(S)$$

If $|S| = n$ then $|P(S)| = 2^n$

What if S is infinite?

Theorem

Let S be a set. Then $|S| < |P(S)|$

Proof.

Define $f : S \rightarrow P(S)$ as

$$f(a) = \{a\}$$

Clearly, f is an injection. Hence $|S| \leq |P(S)|$

Claim: Any function $g : S \rightarrow P(S)$ is not surjective (hence not bijective).

$$A = \{x | x \notin g(x)\} \in P(S)$$

Suppose g is surjective.

$$A = g(a) \text{ for some } a \in S$$

If $|S| = n$ then $|P(S)| = 2^n$

What if S is infinite?

Theorem

Let S be a set. Then $|S| < |P(S)|$

Proof.

Define $f : S \rightarrow P(S)$ as

$$f(a) = \{a\}$$

Clearly, f is an injection. Hence $|S| \leq |P(S)|$

Claim: Any function $g : S \rightarrow P(S)$ is not surjective (hence not bijective).

$$A = \{x | x \notin g(x)\} \in P(S)$$

Suppose g is surjective.

$$\begin{aligned} A &= g(a) \text{ for some } a \in S \\ a \in A &\Leftrightarrow a \in \{x | x \notin g(x)\} \end{aligned}$$

If $|S| = n$ then $|P(S)| = 2^n$

What if S is infinite?

Theorem

Let S be a set. Then $|S| < |P(S)|$

Proof.

Define $f : S \rightarrow P(S)$ as

$$f(a) = \{a\}$$

Clearly, f is an injection. Hence $|S| \leq |P(S)|$

Claim: Any function $g : S \rightarrow P(S)$ is not surjective (hence not bijective).

$$A = \{x | x \notin g(x)\} \in P(S)$$

Suppose g is surjective.

$$\begin{aligned} A &= g(a) \text{ for some } a \in S \\ a \in A &\Leftrightarrow a \in \{x | x \notin g(x)\} \\ &\Leftrightarrow a \notin g(a) \end{aligned}$$

If $|S| = n$ then $|P(S)| = 2^n$

What if S is infinite?

Theorem

Let S be a set. Then $|S| < |P(S)|$

Proof.

Define $f : S \rightarrow P(S)$ as

$$f(a) = \{a\}$$

Clearly, f is an injection. Hence $|S| \leq |P(S)|$

Claim: Any function $g : S \rightarrow P(S)$ is not surjective (hence not bijective).

$$A = \{x | x \notin g(x)\} \in P(S)$$

Suppose g is surjective.

$$\begin{aligned} A &= g(a) \text{ for some } a \in S \\ a \in A &\Leftrightarrow a \in \{x | x \notin g(x)\} \\ &\Leftrightarrow a \notin g(a) \\ &\Leftrightarrow a \notin A \end{aligned}$$

If $|S| = n$ then $|P(S)| = 2^n$

What if S is infinite?

Theorem

Let S be a set. Then $|S| < |P(S)|$

Proof.

Define $f : S \rightarrow P(S)$ as

$$f(a) = \{a\}$$

Clearly, f is an injection. Hence $|S| \leq |P(S)|$

Claim: Any function $g : S \rightarrow P(S)$ is not surjective (hence not bijective).

$$A = \{x | x \notin g(x)\} \in P(S)$$

Suppose g is surjective.

$$\begin{aligned} A &= g(a) \text{ for some } a \in S \\ a \in A &\Leftrightarrow a \in \{x | x \notin g(x)\} \\ &\Leftrightarrow a \notin g(a) \\ &\Leftrightarrow a \notin A \end{aligned}$$

Contradiction to g is surjective.

If $|S| = n$ then $|P(S)| = 2^n$

What if S is infinite?

Theorem

Let S be a set. Then $|S| < |P(S)|$

Proof.

Define $f : S \rightarrow P(S)$ as

$$f(a) = \{a\}$$

Clearly, f is an injection. Hence $|S| \leq |P(S)|$

Claim: Any function $g : S \rightarrow P(S)$ is not surjective (hence not bijective).

$$A = \{x | x \notin g(x)\} \in P(S)$$

Suppose g is surjective.

$$\begin{aligned} A &= g(a) \text{ for some } a \in S \\ a \in A &\Leftrightarrow a \in \{x | x \notin g(x)\} \\ &\Leftrightarrow a \notin g(a) \\ &\Leftrightarrow a \notin A \end{aligned}$$

Contradiction to g is surjective.

$$\text{Hence } |S| \neq |P(S)|$$