

Determinant & It's Properties

Let $A_{n \times n} = [a_{ij}]$ be a $n \times n$ matrix with entries from \mathbb{R} — set of real numbers.

Determinant & It's Properties

Let $A_{n \times n} = [a_{ij}]$ be a $n \times n$ matrix with entries from \mathbb{R} – set of real numbers. You know

Definition:
$$\det(A) := \sum_{j=1}^n (-1)^{1+j} a_{1j} \det(A_{1j})$$

where A_{ij} is the matrix obtained from A by deleting i^{th} row and j^{th} column.

Determinant & It's Properties

Let $A_{n \times n} = [a_{ij}]$ be a $n \times n$ matrix with entries from \mathbb{R} – set of real numbers. You know

Definition:
$$\det(A) := \sum_{j=1}^n (-1)^{1+j} a_{1j} \det(A_{1j})$$

where A_{ij} is the matrix obtained from A by deleting i^{th} row and j^{th} column.

$$\det\left(\begin{bmatrix} a & b \\ c & d \end{bmatrix}\right) = a(d) - b(c)$$

Let $A = [a_{ij}]$ be a $n \times n$ matrix. Define cofactors as, for each i, j ,

$$C_{ij} := (-1)^{i+j} \det(A_{ij})$$

Theorem

The determinant of an $n \times n$ matrix A can be computed by a cofactor expansion across any row or down any column. The expansion across the i th row using the cofactors in (4) is

$$\det A = a_{i1}C_{i1} + a_{i2}C_{i2} + \cdots + a_{in}C_{in}$$

The cofactor expansion down the j th column is

$$\det A = a_{1j}C_{1j} + a_{2j}C_{2j} + \cdots + a_{nj}C_{nj}$$

Let $A = [a_{ij}]$ be a $n \times n$ matrix. Define cofactors as, for each i, j ,

$$C_{ij} := (-1)^{i+j} \det(A_{ij})$$

Theorem

The determinant of an $n \times n$ matrix A can be computed by a cofactor expansion across any row or down any column. The expansion across the i th row using the cofactors in (4) is

$$\det A = a_{i1}C_{i1} + a_{i2}C_{i2} + \cdots + a_{in}C_{in}$$

The cofactor expansion down the j th column is

$$\det A = a_{1j}C_{1j} + a_{2j}C_{2j} + \cdots + a_{nj}C_{nj}$$

Corollary

$$\det(A) = \det(A^T)$$

Theorem

If A is a triangular matrix then $\det(A)$ is the product of diagonal entries.

Theorem

If A is a triangular matrix then $\det(A)$ is the product of diagonal entries.

Note: If a computer performs one trillion multiplications per second, it would have to run for more than 500,000 years to compute determinant of a 25×25 matrix.

We need better "definition" or properties of determinant to calculate determinant easily.

Let's start with new definition of determinant:

Let's start with new definition of determinant:

Definition (Determinant)

Let $M_n(\mathbb{R})$ denote the set of all $n \times n$ matrix with entries from \mathbb{R} .

Then $\det: M_n(\mathbb{R}) \rightarrow \mathbb{R}$ is a function which satisfies

- (1) For each i , \det is a linear function of the i^{th} row, when other $n - 1$ rows are fixed.

Let's start with new definition of determinant:

Definition (Determinant)

Let $M_n(\mathbb{R})$ denote the set of all $n \times n$ matrix with entries from \mathbb{R} .

Then $\det: M_n(\mathbb{R}) \rightarrow \mathbb{R}$ is a function which satisfies

- (1) For each i , \det is a linear function of the i^{th} row, when other $n - 1$ rows are fixed.

Let's start with new definition of determinant:

Definition (Determinant)

Let $M_n(\mathbb{R})$ denote the set of all $n \times n$ matrix with entries from \mathbb{R} . Then $\det: M_n(\mathbb{R}) \rightarrow \mathbb{R}$ is a function which satisfies

- (1) For each i , \det is a linear function of the i^{th} row, when other $n - 1$ rows are fixed.
- (2) $\det(A') = -\det(A)$, where A' is obtained from A by interchanging 2 rows.

Let's start with new definition of determinant:

Definition (Determinant)

Let $M_n(\mathbb{R})$ denote the set of all $n \times n$ matrix with entries from \mathbb{R} . Then $\det: M_n(\mathbb{R}) \rightarrow \mathbb{R}$ is a function which satisfies

- (1) For each i , \det is a linear function of the i^{th} row, when other $n - 1$ rows are fixed.
- (2) $\det(A') = -\det(A)$, where A' is obtained from A by interchanging 2 rows.
- (3) $\det(I) = 1$, where I denotes $n \times n$ identity matrix.

Let's start with new definition of determinant:

Definition (Determinant)

Let $M_n(\mathbb{R})$ denote the set of all $n \times n$ matrix with entries from \mathbb{R} . Then $\det: M_n(\mathbb{R}) \rightarrow \mathbb{R}$ is a function which satisfies

- (1) For each i , \det is a linear function of the i^{th} row, when other $n - 1$ rows are fixed.
- (2) $\det(A') = -\det(A)$, where A' is obtained from A by interchanging 2 rows.
- (3) $\det(I) = 1$, where I denotes $n \times n$ identity matrix.

Theorem

There exists unique function $\det: M_n(\mathbb{R}) \rightarrow \mathbb{R}$ satisfying above three properties.

This function is called determinant function.

Let's start with new definition of determinant:

Definition (Determinant)

Let $M_n(\mathbb{R})$ denote the set of all $n \times n$ matrix with entries from \mathbb{R} . Then $\det: M_n(\mathbb{R}) \rightarrow \mathbb{R}$ is a function which satisfies

- (1) For each i , \det is a linear function of the i^{th} row, when other $n - 1$ rows are fixed.
- (2) $\det(A') = -\det(A)$, where A' is obtained from A by interchanging 2 rows.
- (3) $\det(I) = 1$, where I denotes $n \times n$ identity matrix.

Theorem

There exists unique function $\det: M_n(\mathbb{R}) \rightarrow \mathbb{R}$ satisfying above three properties.

This function is called determinant function.

Verify your definition of determinant satisfies all 3 properties.

Properties of Determinant

1. If 2 rows of A are equal then $\det(A) = 0$.

Properties of Determinant

1. If 2 rows of A are equal then $\det(A) = 0$.
2. Adding/subtracting a scalar multiple of one row from another row leaves determinant unchanged.

Properties of Determinant

1. If 2 rows of A are equal then $\det(A) = 0$.
2. Adding/subtracting a scalar multiple of one row from another row leaves determinant unchanged.
3. If A has a row of zeros then $\det(A) = 0$.

Properties of Determinant

1. If 2 rows of A are equal then $\det(A) = 0$.
2. Adding/subtracting a scalar multiple of one row from another row leaves determinant unchanged.
3. If A has a row of zeros then $\det(A) = 0$.
4. If A is a triangular matrix then $\det(A) = \text{product of diagonal entries}$.

Properties of Determinant

1. If 2 rows of A are equal then $\det(A) = 0$.
2. Adding/subtracting a scalar multiple of one row from another row leaves determinant unchanged.
3. If A has a row of zeros then $\det(A) = 0$.
4. If A is a triangular matrix then $\det(A) = \text{product of diagonal entries}$.
5. $\det(AB) = \det(A) \cdot \det(B)$

Properties of Determinant

1. If 2 rows of A are equal then $\det(A) = 0$.
2. Adding/subtracting a scalar multiple of one row from another row leaves determinant unchanged.
3. If A has a row of zeros then $\det(A) = 0$.
4. If A is a triangular matrix then $\det(A) = \text{product of diagonal entries}$.
5. $\det(AB) = \det(A) \cdot \det(B)$
6. If A is invertible with inverse A^{-1} , then $\det(A^{-1}) = \frac{1}{\det(A)}$.

Properties of Determinant

1. If 2 rows of A are equal then $\det(A) = 0$.
2. Adding/subtracting a scalar multiple of one row from another row leaves determinant unchanged.
3. If A has a row of zeros then $\det(A) = 0$.
4. If A is a triangular matrix then $\det(A) = \text{product of diagonal entries}$.
5. $\det(AB) = \det(A) \cdot \det(B)$
6. If A is invertible with inverse A^{-1} , then $\det(A^{-1}) = \frac{1}{\det(A)}$.

Another definition of determinant: Let $A = [a_{ij}]$ be a $n \times n$ matrix.

$$\det(A) = \sum_{\sigma \in S_n} \text{sign}(\sigma) a_{1\sigma(1)} a_{2\sigma(2)} \cdots a_{n\sigma(n)}$$

Applications of determinant

Area/Volume:

Applications of determinant

Area/Volume:

(1) If A is a $n \times n$ matrix with columns denoted as C_1, \dots, C_n .
Then $|\det(A)|$ gives the volume of a box formed by C_1, \dots, C_n .

Applications of determinant

Area/Volume:

(1) If A is a $n \times n$ matrix with columns denoted as C_1, \dots, C_n . Then $|\det(A)|$ gives the volume of a box formed by C_1, \dots, C_n .

Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be the linear transformation determined by a 2×2 matrix A . If S is a parallelogram in \mathbb{R}^2 , then

$$\{\text{area of } T(S)\} = |\det A| \cdot \{\text{area of } S\} \quad (5)$$

If T is determined by a 3×3 matrix A , and if S is a parallelepiped in \mathbb{R}^3 , then

$$\{\text{volume of } T(S)\} = |\det A| \cdot \{\text{volume of } S\} \quad (6)$$

Applications of determinant

Area/Volume:

(1) If A is a $n \times n$ matrix with columns denoted as C_1, \dots, C_n . Then $|\det(A)|$ gives the volume of a box formed by C_1, \dots, C_n .

Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be the linear transformation determined by a 2×2 matrix A . If S is a parallelogram in \mathbb{R}^2 , then

$$\{\text{area of } T(S)\} = |\det A| \cdot \{\text{area of } S\} \quad (5)$$

If T is determined by a 3×3 matrix A , and if S is a parallelepiped in \mathbb{R}^3 , then

$$\{\text{volume of } T(S)\} = |\det A| \cdot \{\text{volume of } S\} \quad (6)$$

(2) Computation of A^{-1} :

$$A^{-1} = \frac{1}{\det(A)} [c_{ij}]^T,$$

where $c_{ij} = (-1)^{i+j} \det(A_{ij})$ and A_{ij} is a matrix obtained from A by deleting i^{th} row and j^{th} column.

Applications of determinant

Area/Volume:

(1) If A is a $n \times n$ matrix with columns denoted as C_1, \dots, C_n . Then $|\det(A)|$ gives the volume of a box formed by C_1, \dots, C_n .

Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be the linear transformation determined by a 2×2 matrix A . If S is a parallelogram in \mathbb{R}^2 , then

$$\{\text{area of } T(S)\} = |\det A| \cdot \{\text{area of } S\} \quad (5)$$

If T is determined by a 3×3 matrix A , and if S is a parallelepiped in \mathbb{R}^3 , then

$$\{\text{volume of } T(S)\} = |\det A| \cdot \{\text{volume of } S\} \quad (6)$$

(2) Computation of A^{-1} :

$$A^{-1} = \frac{1}{\det(A)} [c_{ij}]^T,$$

where $c_{ij} = (-1)^{i+j} \det(A_{ij})$ and A_{ij} is a matrix obtained from A by deleting i^{th} row and j^{th} column.

(3) It also helps you to find solution to the linear system $AX = b$, when A is invertible matrix.

Suppose a square matrix A has been reduced to an echelon form U by row replacements and row interchanges only.

Determinant and REF

Suppose a square matrix A has been reduced to an echelon form U by row replacements and row interchanges only.

Then

$$\det(A) = (-1)^r \det(U),$$

where r is the no. of row interchanges required to get $U = \text{REF}(A)$.

Determinant of Block diagonal matrices

Let $A = \begin{bmatrix} A_1 & 0 & 0 & 0 & 0 \\ 0 & A_2 & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & 0 & A_r \end{bmatrix}$ be a matrix in block diagonal form, where A_i are submatrices of A .

Determinant of Block diagonal matrices

Let $A = \begin{bmatrix} A_1 & 0 & 0 & 0 & 0 \\ 0 & A_2 & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & 0 & A_r \end{bmatrix}$ be a matrix in block diagonal form, where A_i are submatrices of A .

Then

$$\det(A) = \det(A_1) \cdot \det(A_2) \cdots \det(A_r)$$

Determinant of Block diagonal matrices

Let $A = \begin{bmatrix} A_1 & 0 & 0 & 0 & 0 \\ 0 & A_2 & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & 0 & A_r \end{bmatrix}$ be a matrix in block diagonal form, where A_i are submatrices of A .

Then

$$\det(A) = \det(A_1) \cdot \det(A_2) \cdots \det(A_r)$$

Let $X = \begin{bmatrix} A & B \\ 0 & D \end{bmatrix}$. Then

$$\det(X) = \det(A) \cdot \det(D)$$

Determinant of Block diagonal matrices

Let $A = \begin{bmatrix} A_1 & 0 & 0 & 0 & 0 \\ 0 & A_2 & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & 0 & A_r \end{bmatrix}$ be a matrix in block diagonal form, where A_i are submatrices of A .

Then

$$\det(A) = \det(A_1) \cdot \det(A_2) \cdots \det(A_r)$$

Let $X = \begin{bmatrix} A & B \\ 0 & D \end{bmatrix}$. Then

$$\det(X) = \det(A) \cdot \det(D)$$

$$M = \begin{bmatrix} I & 0 \\ 0 & D \end{bmatrix} \cdot \begin{bmatrix} I & B \\ 0 & I \end{bmatrix} \cdot \begin{bmatrix} A & 0 \\ 0 & I \end{bmatrix}$$

Therefore, $\det(X) = \det(A) \cdot \det(D)$.

If $X = \begin{bmatrix} A & B \\ C & D \end{bmatrix}$ and A is invertible then

$$\det(X) = \det(A) \cdot \det(D - CA^{-1}B)$$

If $X = \begin{bmatrix} A & B \\ C & D \end{bmatrix}$ and A is invertible then

$$\det(X) = \det(A) \cdot \det(D - CA^{-1}B)$$

$X = \begin{bmatrix} I & 0 \\ CA^{-1} & I \end{bmatrix} \cdot \begin{bmatrix} A & 0 \\ 0 & D - CA^{-1}B \end{bmatrix} \cdot \begin{bmatrix} I & A^{-1}B \\ 0 & I \end{bmatrix}$ If D is invertible then

$$\det(X) = \det(D) \cdot \det(A - BD^{-1}C)$$

If $X = \begin{bmatrix} A & B \\ C & D \end{bmatrix}$ and A is invertible then

$$\det(X) = \det(A) \cdot \det(D - CA^{-1}B)$$

$X = \begin{bmatrix} I & 0 \\ CA^{-1} & I \end{bmatrix} \cdot \begin{bmatrix} A & 0 \\ 0 & D - CA^{-1}B \end{bmatrix} \cdot \begin{bmatrix} I & A^{-1}B \\ 0 & I \end{bmatrix}$ If D is invertible then

$$\det(X) = \det(D) \cdot \det(A - BD^{-1}C)$$

$$X = \begin{bmatrix} I & BD^{-1} \\ 0 & I \end{bmatrix} \cdot \begin{bmatrix} A - BD^{-1}C & 0 \\ 0 & D \end{bmatrix} \cdot \begin{bmatrix} I & 0 \\ D^{-1}C & I \end{bmatrix}$$