

Linear System

Definition

A linear System of m equations in n variables- X_1, X_2, \dots, X_n is

$$a_{11}X_1 + a_{12}X_2 + \dots + a_{1n}X_n = b_1$$

$$a_{21}X_1 + a_{22}X_2 + \dots + a_{2n}X_n = b_2$$

$$\vdots$$
$$\vdots$$
$$\vdots$$

$$a_{m1}X_1 + a_{m2}X_2 + \dots + a_{mn}X_n = b_m$$

where $a_{ij}, b_j \in \mathbb{R}$

Linear System

Definition

A linear System of m equations in n variables- X_1, X_2, \dots, X_n is

$$a_{11}X_1 + a_{12}X_2 + \dots + a_{1n}X_n = b_1$$

$$a_{21}X_1 + a_{22}X_2 + \dots + a_{2n}X_n = b_2$$

$$\vdots$$
$$\vdots$$
$$\vdots$$

$$a_{m1}X_1 + a_{m2}X_2 + \dots + a_{mn}X_n = b_m$$

where $a_{ij}, b_j \in \mathbb{R}$

Let's look at linear system of 2 equations in 2 variables:

Solve the system: (1) $x + 2y = 3$, (2) $3x + y = 4$.

Elimination of variables:

Eliminate x by $(2) - 3 \times (1)$ to get $y = 1$.

Cramer's Rule (determinant): $y = \frac{\begin{vmatrix} 1 & 3 \\ 3 & 4 \end{vmatrix}}{\begin{vmatrix} 1 & 2 \\ 3 & 1 \end{vmatrix}} = \frac{4-9}{1-6} = 1$

In either case, back substitution gives $x = 1$

We could also solve for x first and use back substitution for y .

Comparison: For a large system, say 100 equations in 100 variables, elimination method is preferred, since computing the determinants of a 101 matrices of size 100×100 is time-consuming.


Geometry of Linear Equations

$$x+2y=3$$

and

$$3x+y=4$$

represent lines in \mathbb{R}^2 passing through $(0, 3/2)$ and $(3, 0)$ and through $(0, 4)$ and $(4/3, 0)$ respectively.




The intersection of the two lines is the unique point $(1, 1)$. Hence $x = 1$ and $y = 1$ is the solution of above system of linear equations.

3 Equations in 3 Variables

A linear equation in 3 variables represents a plane in a 3-dimensional space \mathbb{R}^3 .

$$x + 2y + 3z = 6$$

passes through $(0, 0, 2)$, $(0, 3, 0)$, $(6, 0, 0)$.



$x + 2y + 3z = 12$ passes through $(0, 0, 4)$, $(0, 6, 0)$, $(12, 0, 0)$.
which is parallel to above plane.

Suppose we have 3 equations in 3 variables, means we have to find intersection of 3 planes, say P_1, P_2, P_3 .

Suppose we have 3 equations in 3 variables, means we have to find intersection of 3 planes, say P_1, P_2, P_3 .

This is same as finding intersection of line L with P_3 (Intersection of P_1, P_2 is L , if P_1, P_2 are not parallel).

Suppose we have 3 equations in 3 variables, means we have to find intersection of 3 planes, say P_1, P_2, P_3 .

This is same as finding intersection of line L with P_3 (Intersection of P_1, P_2 is L , if P_1, P_2 are not parallel).

- ▶ If the line L does intersects with the plane P_3 , then the linear system has **unique** solution.

Suppose we have 3 equations in 3 variables, means we have to find intersection of 3 planes, say P_1, P_2, P_3 .

This is same as finding intersection of line L with P_3 (Intersection of P_1, P_2 is L , if P_1, P_2 are not parallel).

- ▶ If the line L does intersects with the plane P_3 , then the linear system has **unique** solution.
- ▶ If the line L does not intersects with the plane P_3 , then the linear system has **no** solution.

Suppose we have 3 equations in 3 variables, means we have to find intersection of 3 planes, say P_1, P_2, P_3 .

This is same as finding intersection of line L with P_3 (Intersection of P_1, P_2 is L , if P_1, P_2 are not parallel).

- ▶ If the line L does intersects with the plane P_3 , then the linear system has **unique** solution.
- ▶ If the line L does not intersects with the plane P_3 , then the linear system has **no** solution.
- ▶ If the line L is contained in the plane P_3 , then the system has **infinitely many** solutions.

In this case, every point of L is a solution.

Suppose we have 3 equations in 3 variables, means we have to find intersection of 3 planes, say P_1, P_2, P_3 .

This is same as finding intersection of line L with P_3 (Intersection of P_1, P_2 is L , if P_1, P_2 are not parallel).

- ▶ If the line L does intersects with the plane P_3 , then the linear system has **unique** solution.
- ▶ If the line L does not intersects with the plane P_3 , then the linear system has **no** solution.
- ▶ If the line L is contained in the plane P_3 , then the system has **infinitely many** solutions.

In this case, every point of L is a solution.

Question: Can we do the same when number of variables are > 3 ?

Gaussian Elimination: Unique solution

Example: $2x + y + z = 5$, $4x - 6y = -2$, $-2x + 7y + 2z = 9$.

Algorithm: Eliminate x from last 2 equations by $(2) - 2(1)$, and $(3) + (1)$ to get the *equivalent system*:

$$2x + y + z = 5, \quad -8y - 2z = -12, \quad 8y + 3z = 14$$

The first *pivot* is 2, second pivot is -8 . Eliminate y from the last equation to get an equivalent *triangular system*:

$$2x + y + z = 5, \quad -8y - 2z = -12, \quad z = 2$$

Solve this triangular system by *back substitution*, we get

$$z = 2, \quad y = 1, \quad x = 1$$

Observe: This is the only possible solution!

Gaussian Elimination: No solution

Example: $2x + y + z = 5$, $4x - 6y = -2$, $-2x + 7y + z = 9$.

Step 1 Eliminate x (using the 1st pivot 2) to get:

$$2x + y + z = 5, \quad -8y - 2z = -12, \quad 8y + 2z = 14$$

Step 2: Eliminate y (using the 2nd pivot -8) to get:

$$2x + y + z = 5, \quad -8y - 2z = -12, \quad 0 = 2.$$

The last equation shows that there is no solution, i.e., the system is *inconsistent*.

Geometric reasoning: In Step 1, notice we get two distinct parallel planes $8y + 2z = 12$ and $8y + 2z = 14$.

They have no point in common.

Note: The planes in the original system were not parallel, but in an equivalent system, we get two distinct parallel planes!

Gaussian Elimination: Infinitely solution

Example: $2x + y + z = 5$, $4x - 6y = -2$, $-2x + 7y + z = 7$.

Step 1 Eliminate x (using the 1st pivot 2) to get:

$$2x + y + z = 5, \quad -8y - 2z = -12, \quad 8y + 2z = 12$$

Step 2: Eliminate y (using the 2nd pivot -8) to get:

$$2x + y + z = 5, \quad -8y - 2z = -12, \quad 0 = 0.$$

There are only two equations. For every value of z , values for x and y are obtained by back-substitution, e.g. $(1, 1, 2)$ or $(\frac{7}{4}, \frac{3}{2}, 0)$. Hence the system has infinitely many solutions.

Geometric reasoning: In Step 1, notice we get two parallel planes $-8y - 2z = 12$ and $8y + 2z = 12$.

They give the same plane. Hence we are looking at the intersection of the two planes, $2x + y + z = 5$ and $8x + 2z = 12$, which is a line.

A system of linear equations has exactly one of following

1. no solution; (**inconsistent system**)
2. exactly one solution; (**consistent system**)
3. infinitely many solutions. (**consistent system**)

A system of linear equations has exactly one of following

1. no solution; (**inconsistent system**)
2. exactly one solution; (**consistent system**)
3. infinitely many solutions. (**consistent system**)

Solve

Gaussian Elimination: Matrix form $AX=b$

Example: $2x + y + z = 5$, $4x - 6y = -2$, $-2x + 7y + 2z = 9$.

Note that the last column is the RHS column vector b .

$$\left(\begin{array}{ccc|c} 2 & 1 & 1 & 5 \\ 4 & -6 & 0 & -2 \\ -2 & 7 & 2 & 9 \end{array} \right) \rightarrow \left(\begin{array}{ccc|c} 2 & 1 & 1 & 5 \\ 0 & -8 & -2 & -12 \\ 0 & 8 & 3 & 14 \end{array} \right) \rightarrow$$

$$\left(\begin{array}{ccc|c} 2 & 1 & 1 & 5 \\ 0 & -8 & -2 & -12 \\ 0 & 0 & 1 & 2 \end{array} \right)$$

Gaussian Elimination: Matrix form $AX=b$

Example: $2x + y + z = 5$, $4x - 6y = -2$, $-2x + 7y + 2z = 9$.

Note that the last column is the RHS column vector b .

$$\left(\begin{array}{ccc|c} 2 & 1 & 1 & 5 \\ 4 & -6 & 0 & -2 \\ -2 & 7 & 2 & 9 \end{array} \right) \rightarrow \left(\begin{array}{ccc|c} 2 & 1 & 1 & 5 \\ 0 & -8 & -2 & -12 \\ 0 & 8 & 3 & 14 \end{array} \right) \rightarrow$$

$$\left(\begin{array}{ccc|c} 2 & 1 & 1 & 5 \\ 0 & -8 & -2 & -12 \\ 0 & 0 & 1 & 2 \end{array} \right)$$

The last matrix corresponds to

$$z = 2,$$

$$-8y - 2z = -12,$$

$$2x + y + z = 5.$$

The last matrix corresponds to

$$z = 2,$$

$$-8y - 2z = -12,$$

$$2x + y + z = 5. \text{ Solving this gives } z = 2, y = 1, x = 1.$$

The last matrix corresponds to

$$z = 2,$$

$$-8y - 2z = -12,$$

$$2x + y + z = 5. \text{ Solving this gives } z = 2, y = 1, x = 1.$$

Gaussian Algorithm

For Row Echelon Form

1. Find the leftmost nonzero column.
2. Select a nonzero entry in that column (**pivot**). Bring it on top by interchanging rows
3. Use Elementary row operations to make all entries below that nonzero entry 0.
4. Ignore that row and columns before (including of leading entry) above found column. Apply steps 1-3 for remaining submatrix. Go on till all rows are exhausted.

Gaussian Algorithm

For Row Echelon Form

1. Find the leftmost nonzero column.
2. Select a nonzero entry in that column (**pivot**). Bring it on top by interchanging rows
3. Use Elementary row operations to make all entries below that nonzero entry 0.
4. Ignore that row and columns before (including of leading entry) above found column. Apply steps 1-3 for remaining submatrix. Go on till all rows are exhausted.

For Reduced Row Echelon Form

5. Make all leading entries 1 by elementary row operation ($R_i \rightarrow cR_i$)
6. Make all entries in a column above leading 1 zero by elementary row operation ($R_i \rightarrow R_i - cR_j$)

$$A = \begin{bmatrix} 0 & 3 & -6 & 6 \\ 3 & -7 & 8 & -5 \\ 3 & -9 & 12 & -9 \end{bmatrix}$$

Reduced Row Echelon Form (RREF) of A is

$$A' = \begin{bmatrix} 1 & 0 & -2 & 3 \\ 0 & 1 & -2 & 2 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

$$A = \begin{bmatrix} 0 & 3 & -6 & 6 \\ 3 & -7 & 8 & -5 \\ 3 & -9 & 12 & -9 \end{bmatrix}$$

Reduced Row Echelon Form (RREF) of A is

$$A' = \begin{bmatrix} 1 & 0 & -2 & 3 \\ 0 & 1 & -2 & 2 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Theorem

1. A linear system is inconsistent if and only if REF or RREF of its augmented matrix has a row of the form $0 \ 0 \ \cdots \ 0 \ b$, for some nonzero b .
2. A linear system has unique solution if and only if no. of nonzero rows and no. of columns of REF or RREF of its augmented matrix are same.

Triangular factorization: $A = LU$ Given a square matrix A , we can find L a lower Triangular matrix with 1s on the diagonal and U upper Triangular matrix such that $A = LU$.

For $A = \begin{bmatrix} 1 & 2 \\ 3 & 8 \end{bmatrix}$, $L = \begin{bmatrix} 1 & 0 \\ 3 & 1 \end{bmatrix}$ and $U = \begin{bmatrix} 1 & 2 \\ 0 & 2 \end{bmatrix}$

Calculation of A^{-1} : Gauss-Jordan Method

Let $AA^{-1} = I$

If X_1 is a first column of A^{-1} then $AX_1 = e_1$, where $e_1 = [1 \ 0 \ \cdots \ 0]^T$.

We can use Gaussian Elimination to find X_1 —first column of A^{-1} .

Calculation of A^{-1} : Gauss-Jordan Method

Let $AA^{-1} = I$

If X_1 is a first column of A^{-1} then $AX_1 = e_1$, where $e_1 = [1 \ 0 \ \cdots \ 0]^T$.

We can use Gaussian Elimination to find X_1 —first column of A^{-1} .

Let $A = \begin{bmatrix} 1 & 2 \\ 3 & 8 \end{bmatrix}$.

Consider

$$[A|I] = \left[\begin{array}{cc|cc} 1 & 2 & 1 & 0 \\ 3 & 8 & 0 & 1 \end{array} \right]$$

Perform elementary row operations on $[A|I]$ to convert A into I .
Then I will change to A^{-1} .

Vectors in \mathbb{R}^n :


$$\begin{bmatrix} a_1 \\ a_2 \\ \vdots \\ a_n \end{bmatrix}$$

where $a_1, a_2, \dots, a_n \in \mathbb{R}$.

$$\begin{bmatrix} 2 \\ 3 \end{bmatrix} \text{ is a vector in } \mathbb{R}^2.$$

$$\begin{bmatrix} 2 \\ 3 \\ 0 \end{bmatrix} \text{ is a vector in } \mathbb{R}^3.$$

Vector addition and scalar multiplication are done like matrices.



Linear Combinations

Consider a linear system

$$x_2 - 4x_3 = 8$$

$$2x_1 - 3x_2 + 2x_3 = 1$$

$$5x_1 - 8x_2 + 7x_3 = 1$$

It can be written as

$$x_1 \begin{bmatrix} 0 \\ 2 \\ 5 \end{bmatrix} + x_2 \begin{bmatrix} 1 \\ -3 \\ -8 \end{bmatrix} + x_3 \begin{bmatrix} -4 \\ 2 \\ 7 \end{bmatrix} = \begin{bmatrix} 8 \\ 1 \\ 1 \end{bmatrix}$$

LHS of $=$ is called linear combination of

$$u = \begin{bmatrix} 0 \\ 2 \\ 5 \end{bmatrix}, v = \begin{bmatrix} 1 \\ -3 \\ -8 \end{bmatrix} \& w = \begin{bmatrix} -4 \\ 2 \\ 7 \end{bmatrix}$$

Above equation is called **vector equation**.

Linear system and Vector Equation

A linear system $AX = b$ is same as $x_1 C_1 + x_2 C_2 + \cdots + x_n C_n = b$ where C_i is i^{th} column of A .

Linear system and Vector Equation

A linear system $AX = b$ is same as $x_1 C_1 + x_2 C_2 + \cdots + x_n C_n = b$ where C_i is i^{th} column of A .

Theorem

$AX = b$ has a solution if and only if b is a linear combination of columns of A .

Linear system and Vector Equation

A linear system $AX = b$ is same as $x_1 C_1 + x_2 C_2 + \cdots + x_n C_n = b$ where C_i is i^{th} column of A .

Theorem

$AX = b$ has a solution if and only if b is a linear combination of columns of A .

Hence we need to study linear combination of columns of A .

Linear system and Vector Equation

A linear system $AX = b$ is same as $x_1 C_1 + x_2 C_2 + \cdots + x_n C_n = b$ where C_i is i^{th} column of A .

Theorem

$AX = b$ has a solution if and only if b is a linear combination of columns of A .

Hence we need to study linear combination of columns of A .

Column space of A = set of all linear combinations of columns of A .

$$\text{Col}(A) = \{x_1 C_1 + x_2 C_2 + \cdots + x_n C_n \mid x_i \in \mathbb{R} \text{ for all } i\}$$

$\therefore AX = b$ has a solution iff $b \in \text{Col}(A)$.

If $v_1, v_2, \dots, v_p \in \mathbb{R}^n$, then

$\text{Span} \{v_1, v_2, \dots, v_p\} = \{c_1 v_1 + c_2 v_2 + \dots + c_p v_p \mid c_1, c_2, \dots, c_p \in \mathbb{R}\}$,
the set of all linear combinations of v_1, v_2, \dots, v_p .

If $v_1, v_2, \dots, v_p \in \mathbb{R}^n$, then

$\text{Span} \{v_1, v_2, \dots, v_p\} = \{c_1 v_1 + c_2 v_2 + \dots + c_p v_p \mid c_1, c_2, \dots, c_p \in \mathbb{R}\}$,
the set of all linear combinations of v_1, v_2, \dots, v_p .

Let

$$u = \begin{bmatrix} 2 \\ 3 \end{bmatrix}, v = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

$$\text{Span} \{v\} = \{c.v \mid c \in \mathbb{R}\} = \left\{ \begin{bmatrix} c \\ 0 \end{bmatrix} \mid c \in \mathbb{R} \right\}$$

If $v_1, v_2, \dots, v_p \in \mathbb{R}^n$, then

$\text{Span} \{v_1, v_2, \dots, v_p\} = \{c_1 v_1 + c_2 v_2 + \dots + c_p v_p \mid c_1, c_2, \dots, c_p \in \mathbb{R}\}$,
the set of all linear combinations of v_1, v_2, \dots, v_p .

Let

$$u = \begin{bmatrix} 2 \\ 3 \end{bmatrix}, v = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

$\text{Span} \{v\} = \{c.v \mid c \in \mathbb{R}\} = \left\{ \begin{bmatrix} c \\ 0 \end{bmatrix} \mid c \in \mathbb{R} \right\}$ Let $u \in \mathbb{R}^n$. Then

$\text{span}\{u\} = \{x.u \mid x \in \mathbb{R}\}$.

Geometrically it is a line passing through origin and point u .

Let $u, v \in \mathbb{R}^2$. Then $\text{span}\{u, v\} = \{x.u + y.v \mid x, y \in \mathbb{R}\}$.

Geometrically, what will it be?

If $v_1, v_2, \dots, v_p \in \mathbb{R}^n$, then

$\text{Span} \{v_1, v_2, \dots, v_p\} = \{c_1 v_1 + c_2 v_2 + \dots + c_p v_p \mid c_1, c_2, \dots, c_p \in \mathbb{R}\}$,
the set of all linear combinations of v_1, v_2, \dots, v_p .

Let

$$u = \begin{bmatrix} 2 \\ 3 \end{bmatrix}, v = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

$\text{Span} \{v\} = \{c.v \mid c \in \mathbb{R}\} = \left\{ \begin{bmatrix} c \\ 0 \end{bmatrix} \mid c \in \mathbb{R} \right\}$ Let $u \in \mathbb{R}^n$. Then

$\text{span}\{u\} = \{x.u \mid x \in \mathbb{R}\}$.

Geometrically it is a line passing through origin and point u .

Let $u, v \in \mathbb{R}^2$. Then $\text{span}\{u, v\} = \{x.u + y.v \mid x, y \in \mathbb{R}\}$.

Geometrically, what will it be?

For a matrix A ,

$\text{Col}(A) := \text{span}\{C_1, C_2, \dots, C_n\} \subseteq \mathbb{R}^m$

The equation $Ax = b$ has a solution if and only if b is a linear combination of the columns of A .

The equation $Ax = b$ has a solution if and only if b is a linear combination of the columns of A .

Theorem


Let A be an $m \times n$ matrix. Then the following statements are logically equivalent

- 1. For each $b \in \mathbb{R}^m$, the equation $AX = b$ has a solution.*
- 2. Each $b \in \mathbb{R}^m$, is a linear combination of the columns of A .*
- 3. $\text{Col}(A) = \mathbb{R}^m$.*
- 4. A has a pivot position in every row.*

Homogeneous Linear System

The homogeneous equation $AX = 0$ has a nontrivial solution if and only if the $\text{REF}(A)$ has at least one free variable.

Suppose augmented form $[A|0]$ is




7_homo.PNG


Homogeneous Linear System

The **homogeneous equation** $AX = 0$ has a nontrivial solution if and only if the $\text{REF}(A)$ has at least one free variable.

Suppose augmented form $[A|0]$ is



7_homo.PNG



7_ex3.PNG

Non homogeneous linear system

Theorem

Suppose the equation $Ax = b$ is consistent for some given b , and let p be a solution (i.e., $Ap = b$). Then the solution set of $Ax = b$ is the set of all vectors of the form $w = p + v_h$, where v_h is any solution of the homogeneous equation $Ax = 0$.

Non homogeneous linear system

Theorem

Suppose the equation $Ax = b$ is consistent for some given b , and let p be a solution (i.e., $Ap = b$). Then the solution set of $Ax = b$ is the set of all vectors of the form $w = p + v_h$, where v_h is any solution of the homogeneous equation $Ax = 0$.




Write down the solutions of $AX = 0$ in parametric form where $A =$

7_para1.PNG

Chemical Equations

Consider a chemical reaction of propane gas with oxygen to form carbon dioxide and water.



where x_i are the no. of molecules

Network flow

