

# MA102: Linear Algebra and Matrices: Course Content

**Matrices and Linear systems:** The System of Linear Equations:  $Ax=b$ , Row Reduction and Echelon forms (Gaussian Elimination), Matrix operations (addition, multiplication), Block-Partitioned Matrices and Block Operations, Elementary Row and Column Operations, Determinant and its Properties, Cofactor Expansion, Rank of a Matrix, Gauss Jordan Method for matrix inversion

**Canonical Factorizations:** Eigenvalues and Eigenvectors, Companion Matrices and Characteristic Polynomial, diagonalization of Matrices with a Full-Set of Eigenvectors, The Cayley-Hamilton Theorem, Triangulization and Unitary Diagonalization of a Matrix, Schur's Lemma and the Spectral Theorem, QR-Decomposition, QR-Algorithm, Singular Value Decomposition.

**Vector Spaces:** Vector Space over the set real numbers (Field), Linear Independence of Vectors, Bases in a Vector Space, Dimension of a Vector Space, Direct Sum Decomposition of a Vector Space, Linear Transformation (LT), Change of Bases, Canonical forms, Rank of a LT.

**Numerical methods:** Iterative methods (Jacobi, Gauss-Seidel, Relaxation) for linear systems, computing eigenvalues and eigenvectors.

# Linear Algebra and Matrices

## Evaluation and Grading policy:

Assignments and Quizzes: 5%

Tutorial: 15%

Mini-Project: 5%

Mid-semester Exam.: 30%

End-semester Exam.: 45% [Text & Reference books:](#)

*Linear Algebra and its Applications*, David C. Lay, 4th Ed, Pearson, 2016.

*Introduction to Linear Algebra*, Gilbert Strang, 5th Ed, SIAM, 2016.

*Linear Algebra*, Kunze Ray, Hoffman Kenneth, 2nd Ed, Phi Learning, 2014.

*Fundamentals of Matrix Computations*, David S. Watkins, 3rd ed, Wiley.

**Question:** Why a computer science student should learn Linear Algebra?

**Question:** Why a computer science student should learn Linear Algebra?

ChatGPT:

1. **Foundational for Computer Graphics and Computer Vision:** Linear algebra is fundamental in understanding and implementing algorithms for computer graphics, computer vision, and image processing. Concepts like transformations, rotations, and scaling can be efficiently represented and manipulated using linear algebra.
2. **Machine Learning and Data Science:** Many machine learning algorithms rely on linear algebra for operations such as matrix multiplication, eigenvalue decomposition, and singular value decomposition. Understanding linear algebra is crucial for designing and implementing algorithms in data science and machine learning.
3. **Computer Networking and Distributed Systems:** In computer networking and distributed systems, linear algebra is used to model and analyze various aspects, such as network flows, connectivity, and optimization problems. It is particularly important for understanding algorithms in graph theory.

- 4 Algorithm Analysis and Complexity: Understanding linear algebra helps in analyzing the time and space complexity of algorithms. Matrices and vectors often appear in algorithmic analysis, and knowledge of linear algebra enables students to analyze the efficiency of algorithms in terms of computational resources.
- 5 Computer Simulations and Numerical Methods: Linear algebra is used extensively in computer simulations and numerical methods. Solving systems of linear equations, eigenvalue problems, and numerical optimization are common tasks in scientific computing, physics simulations, and engineering applications.
- 6 Database Systems and Information Retrieval: In database systems and information retrieval, linear algebra is employed for tasks such as indexing, similarity calculations, and data representation. Understanding linear algebra is beneficial for developing efficient and scalable data storage and retrieval systems.

- 7 Coding and Software Development: Linear algebra provides a powerful and concise way to express mathematical and computational concepts in coding. Libraries and frameworks in various programming languages often leverage linear algebra operations to optimize code and enhance performance.
- 8 Signal Processing: Signal processing, used in areas like audio processing and image processing, heavily relies on linear algebra. Understanding concepts like Fourier transforms, convolution, and signal representation requires a solid foundation in linear algebra.
- 9 Robotics and Control Systems: In robotics and control systems, linear algebra is used for modeling the dynamics of robotic systems, designing control algorithms, and analyzing the stability of control systems.

- 10 Problem-solving and Critical Thinking: Learning linear algebra helps develop problem-solving skills and encourages abstract and critical thinking. It provides a structured and formalized way to represent and solve a wide range of problems, fostering a deeper understanding of mathematical concepts. In summary, linear algebra is a versatile and foundational mathematical tool that has widespread applications in various areas of computer science. It forms the basis for understanding and solving complex problems across different domains within the field.

Suppose you have sufficient no. of currency notes of denominations Rs. 10, Rs. 20, Rs. 50 and Rs. 100. Your canteen bill is Rs. 260. How many notes of each denomination will you need to pay?



Suppose you have sufficient no. of currency notes of denominations Rs. 10, Rs. 20, Rs. 50 and Rs. 100. Your canteen bill is Rs. 260. How many notes of each denomination will you need to pay?

**Mathematical model:** Let  $t$ ,  $u$ ,  $v$  and  $w$  be number of notes of denominations Rs. 10, Rs. 20, Rs. 50 and Rs. 100 respectively that you will pay.

Suppose you have sufficient no. of currency notes of denominations Rs. 10, Rs. 20, Rs. 50 and Rs. 100. Your canteen bill is Rs. 260. How many notes of each denomination will you need to pay?

**Mathematical model:** Let  $t, u, v$  and  $w$  be number of notes of denominations Rs. 10, Rs. 20, Rs. 50 and Rs. 100 respectively that you will pay.

$$10t + 20u + 50v + 100w = 260 \dots (*)$$
$$t, u, v \geq 0$$

Suppose you have sufficient no. of currency notes of denominations Rs. 10, Rs. 20, Rs. 50 and Rs. 100. Your canteen bill is Rs. 260. How many notes of each denomination will you need to pay?

**Mathematical model:** Let  $t, u, v$  and  $w$  be number of notes of denominations Rs. 10, Rs. 20, Rs. 50 and Rs. 100 respectively that you will pay.

$$10t + 20u + 50v + 100w = 260 \dots (*)$$
$$t, u, v \geq 0$$

Find all solutions to equation  $(*)$  without non-negative constraints/conditions.

# Covid Pandemic

Suppose 1 cr cases of Covid 19 are found in city X which has a population of 301 cr. Hence city was locked and nobody is allowed to go in or out. Every day 20% are getting infected and 10% were getting cured. How many will be infected (but not cured) after 10 days?

Suppose 1 cr cases of Covid 19 are found in city X which has a population of 301 cr. Hence city was locked and nobody is allowed to go in or out. Every day 20% are getting infected and 10% were getting cured. How many will be infected (but not cured) after 10 days?

$$\text{Infected: } 300 \times (0.20) + 1 \times (0.90) = 60.9$$

$$\text{Healthy: } 300 \times (0.80) + 1 \times (0.10) = 240.1$$

# Covid Pandemic

Suppose 1 cr cases of Covid 19 are found in city X which has a population of 301 cr. Hence city was locked and nobody is allowed to go in or out. Every day 20% are getting infected and 10% were getting cured. How many will be infected (but not cured) after 10 days?

$$\text{Infected: } 300 \times (0.20) + 1 \times (0.90) = 60.9$$

$$\text{Healthy: } 300 \times (0.80) + 1 \times (0.10) = 240.1$$

$$\begin{bmatrix} 0.2 & 0.9 \\ 0.8 & 0.1 \end{bmatrix} \cdot \begin{bmatrix} 300 \\ 1 \end{bmatrix} = \begin{bmatrix} 60.9 \\ 240.1 \end{bmatrix}$$

$$AX_0 = X_1$$

$$X_{10} = A^{10}X_0$$

# System of linear equations

Let  $M$  be a packet which contains 5kg of wheat & 2 kg of rice and  $N$  be a packet which contains 2 kg of wheat & 5 kg of rice. How many packets of  $M$  and  $N$  should you buy to get 19kg of wheat and 15kgs of rice?

# System of linear equations

Let  $M$  be a packet which contains 5kg of wheat & 2 kg of rice and  $N$  be a packet which contains 2 kg of wheat & 5 kg of rice. How many packets of  $M$  and  $N$  should you buy to get 19kg of wheat and 15kgs of rice?

**Question:** Is it possible to buy 39 kg of wheat and 39kg of rice?



# System of linear equations

Let  $M$  be a packet which contains 5kg of wheat & 2 kg of rice and  $N$  be a packet which contains 2 kg of wheat & 5 kg of rice. How many packets of  $M$  and  $N$  should you buy to get 19kg of wheat and 15kgs of rice?

**Question:** Is it possible to buy 39 kg of wheat and 39kg of rice?

**Question:** What if we have  $M$  &  $N$  containing 3 types of grains?

Entry of matrices.

# System of linear equations

Let  $M$  be a packet which contains 5kg of wheat & 2 kg of rice and  $N$  be a packet which contains 2 kg of wheat & 5 kg of rice. How many packets of  $M$  and  $N$  should you buy to get 19kg of wheat and 15kgs of rice?

**Question:** Is it possible to buy 39 kg of wheat and 39kg of rice?

**Question:** What if we have  $M$  &  $N$  containing 3 types of grains?

Entry of matrices.

# Matrix operations(addition, multiplication)

**Size/Dimension of a matrix:** A matrix with  $m$  rows and  $n$  columns is called an  $m \times n$  matrix, where  $m$  &  $n$  are called its dimensions.

# Matrix operations(addition, multiplication)

**Size/Dimension of a matrix:** A matrix with  $m$  rows and  $n$  columns is called an  $m \times n$  matrix, where  $m$  &  $n$  are called its dimensions.

Addition of two matrices:

**Question:** When can we add two matrices?

# Matrix operations(addition, multiplication)

**Size/Dimension of a matrix:** A matrix with  $m$  rows and  $n$  columns is called an  $m \times n$  matrix, where  $m$  &  $n$  are called its dimensions.

Addition of two matrices:

**Question:** When can we add two matrices?

**Ans:** If they have same size/dimension.

# Matrix operations(addition, multiplication)

**Size/Dimension of a matrix:** A matrix with  $m$  rows and  $n$  columns is called an  $m \times n$  matrix, where  $m$  &  $n$  are called its dimensions.

Addition of two matrices:

**Question:** When can we add two matrices?

**Ans:** If they have same size/dimension.

If  $A = [a_{ij}]$ ,  $B = [b_{ij}]$  then  $A + B = [c_{ij}]$ , where  $c_{ij} = a_{ij} + b_{ij}$

# Matrix operations(addition, multiplication)

**Size/Dimension of a matrix:** A matrix with  $m$  rows and  $n$  columns is called an  $m \times n$  matrix, where  $m$  &  $n$  are called its dimensions.

Addition of two matrices:

**Question:** When can we add two matrices?

**Ans:** If they have same size/dimension.

If  $A = [a_{ij}]$ ,  $B = [b_{ij}]$  then  $A + B = [c_{ij}]$ , where  $c_{ij} = a_{ij} + b_{ij}$

Multiplication of two matrices:

**Question:** When can we multiply two matrices?

# Matrix operations(addition, multiplication)

**Size/Dimension of a matrix:** A matrix with  $m$  rows and  $n$  columns is called an  $m \times n$  matrix, where  $m$  &  $n$  are called its dimensions.

Addition of two matrices:

**Question:** When can we add two matrices?

**Ans:** If they have same size/dimension.

If  $A = [a_{ij}]$ ,  $B = [b_{ij}]$  then  $A + B = [c_{ij}]$ , where  $c_{ij} = a_{ij} + b_{ij}$

Multiplication of two matrices:

**Question:** When can we multiply two matrices?

**Ans:** Yes, if no. of columns of first matrix = no. of rows of second matrix.



# Matrix operations(addition, multiplication)

**Size/Dimension of a matrix:** A matrix with  $m$  rows and  $n$  columns is called an  $m \times n$  matrix, where  $m$  &  $n$  are called its dimensions.

Addition of two matrices:

**Question:** When can we add two matrices?

**Ans:** If they have same size/dimension.

If  $A = [a_{ij}]$ ,  $B = [b_{ij}]$  then  $A + B = [c_{ij}]$ , where  $c_{ij} = a_{ij} + b_{ij}$

Multiplication of two matrices:

**Question:** When can we multiply two matrices?

**Ans:** Yes, if no. of columns of first matrix = no. of rows of second matrix.

If  $A = [a_{ij}]$  of order  $m \times n$  &  $B = [b_{ij}]$  of order  $n \times p$  then  $AB$  is of size  $m \times p$  with

$$AB = [c_{ij}], \text{ where } c_{ij} = \sum_{k=1}^n a_{ik} b_{kj}$$

# Matrix operations(addition, multiplication)

**Size/Dimension of a matrix:** A matrix with  $m$  rows and  $n$  columns is called an  $m \times n$  matrix, where  $m$  &  $n$  are called its dimensions.

Addition of two matrices:

**Question:** When can we add two matrices?

**Ans:** If they have same size/dimension.

If  $A = [a_{ij}]$ ,  $B = [b_{ij}]$  then  $A + B = [c_{ij}]$ , where  $c_{ij} = a_{ij} + b_{ij}$

Multiplication of two matrices:

**Question:** When can we multiply two matrices?

**Ans:** Yes, if no. of columns of first matrix = no. of rows of second matrix.

If  $A = [a_{ij}]$  of order  $m \times n$  &  $B = [b_{ij}]$  of order  $n \times p$  then  $AB$  is of size  $m \times p$  with

$$AB = [c_{ij}], \text{ where } c_{ij} = \sum_{k=1}^n a_{ik} b_{kj}$$

**Question** Is  $AB = BA$  for any two matrices  $A, B$ ? 

Let  $A, B, C$  be matrices of same dimension and  $r, s \in \mathbb{R}$ . Then

1.  $A+B=B+A$
2.  $r.(A+B)=r.A+r.B$
3.  $(A+B)+C=A+(B+C)$
4.  $(r+s).A=r.A+s.A$
5.  $r.(s.A)=(rs).A$

Let  $A, B, C$  be matrices of same dimension and  $r, s \in \mathbb{R}$ . Then

1.  $A+B=B+A$
2.  $r.(A+B)=r.A+r.B$
3.  $(A+B)+C=A+(B+C)$
4.  $(r+s).A=r.A+s.A$
5.  $r.(s.A)=(rs).A$

If  $b_1, b_2, \dots, b_p$  are columns of  $B$  then columns of  $AB$  are

Let  $A, B, C$  be matrices of same dimension and  $r, s \in \mathbb{R}$ . Then

1.  $A+B=B+A$
2.  $r.(A+B)=r.A+r.B$
3.  $(A+B)+C=A+(B+C)$
4.  $(r+s).A=r.A+s.A$
5.  $r.(s.A)=(rs).A$

If  $b_1, b_2, \dots, b_p$  are columns of  $B$  then columns of  $AB$  are  $Ab_1, Ab_2, \dots, Ab_p$  respectively.

Let  $A, B, C$  be matrices of same dimension and  $r, s \in \mathbb{R}$ . Then

1.  $A+B=B+A$
2.  $r.(A+B)=r.A+r.B$
3.  $(A+B)+C=A+(B+C)$
4.  $(r+s).A=r.A+s.A$
5.  $r.(s.A)=(rs).A$

If  $b_1, b_2, \dots, b_p$  are columns of  $B$  then columns of  $AB$  are  $Ab_1, Ab_2, \dots, Ab_p$  respectively.

What can you say about rows of  $AB$ ?

Let  $A, B, C$  be matrices of same dimension and  $r, s \in \mathbb{R}$ . Then

1.  $A+B=B+A$
2.  $r.(A+B)=r.A+r.B$
3.  $(A+B)+C=A+(B+C)$
4.  $(r+s).A=r.A+s.A$
5.  $r.(s.A)=(rs).A$

If  $b_1, b_2, \dots, b_p$  are columns of  $B$  then columns of  $AB$  are  $Ab_1, Ab_2, \dots, Ab_p$  respectively.

What can you say about rows of  $AB$ ?

6.  $A.(B.C)=(A.B).C$
7.  $A.(B+C)=A.B+A.C$
8.  $(B+C).A=B.A+C.A$
9.  $r.(A.B)=(r.A).B=A.(r.B)$
10.  $I.A=A.I=A$

Let  $A, B, C$  be matrices of same dimension and  $r, s \in \mathbb{R}$ . Then

1.  $A+B=B+A$
2.  $r.(A+B)=r.A+r.B$
3.  $(A+B)+C=A+(B+C)$
4.  $(r+s).A=r.A+s.A$
5.  $r.(s.A)=(rs).A$

If  $b_1, b_2, \dots, b_p$  are columns of  $B$  then columns of  $AB$  are  $Ab_1, Ab_2, \dots, Ab_p$  respectively.

What can you say about rows of  $AB$ ?

6.  $A.(B.C)=(A.B).C$
7.  $A.(B+C)=A.B+A.C$
8.  $(B+C).A=B.A+C.A$
9.  $r.(A.B)=(r.A).B=A.(r.B)$
10.  $I.A=A.I=A$

Question: Does  $AB = 0$  mean either  $A = 0$  or  $B = 0$ ?



**Transpose of a matrix:** If  $A = [a_{ij}]$  is a matrix then  $(i, j)^{th}$  entry of  $A^T$  is  $a_{ji}$ .

**Transpose of a matrix:** If  $A = [a_{ij}]$  is a matrix then  $(i, j)^{th}$  entry of  $A^T$  is  $a_{ji}$ .

1.  $(A^T)^T = A.$

2.  $(A + B)^T = A^T + B^T$

3.  $(rA)^T = r.A^T$

4.  $(AB)^T =$

**Transpose of a matrix:** If  $A = [a_{ij}]$  is a matrix then  $(i, j)^{th}$  entry of  $A^T$  is  $a_{ji}$ .

1.  $(A^T)^T = A.$

2.  $(A + B)^T = A^T + B^T$

3.  $(rA)^T = r.A^T$

4.  $(AB)^T = B^T A^T.$

**Transpose of a matrix:** If  $A = [a_{ij}]$  is a matrix then  $(i, j)^{th}$  entry of  $A^T$  is  $a_{ji}$ .

1.  $(A^T)^T = A$ .
2.  $(A + B)^T = A^T + B^T$
3.  $(rA)^T = r.A^T$
4.  $(AB)^T = B^T A^T$ .

**Inverse of a Matrix:** A matrix  $A$  is said to be invertible if there exists a matrix  $B$  such that

$$AB = BA = I.$$

$B$  is called inverse of  $A$ .

**Transpose of a matrix:** If  $A = [a_{ij}]$  is a matrix then  $(i, j)^{th}$  entry of  $A^T$  is  $a_{ji}$ .

1.  $(A^T)^T = A$ .

2.  $(A + B)^T = A^T + B^T$

3.  $(rA)^T = r.A^T$

4.  $(AB)^T = B^T A^T$ .

**Inverse of a Matrix:** A matrix  $A$  is said to be invertible if there exists a matrix  $B$  such that

$$AB = BA = I.$$

$B$  is called inverse of  $A$ .

If  $A$  is invertible then  $A$  must be square matrix.

**Transpose of a matrix:** If  $A = [a_{ij}]$  is a matrix then  $(i, j)^{th}$  entry of  $A^T$  is  $a_{ji}$ .

1.  $(A^T)^T = A$ .
2.  $(A + B)^T = A^T + B^T$
3.  $(rA)^T = r.A^T$
4.  $(AB)^T = B^T A^T$ .

**Inverse of a Matrix:** A matrix  $A$  is said to be invertible if there exists a matrix  $B$  such that

$$AB = BA = I.$$

$B$  is called inverse of  $A$ .

If  $A$  is invertible than  $A$  must be square matrix.

5.  $(A^{-1})^{-1} = A$ .
6.  $(AB)^{-1} = B^{-1}A^{-1}$ .
7.  $(A^T)^{-1} = (A^{-1})^T$ .

Question: If  $A, B, C$  are square matrices of same dimension such that  $AB = CA = I$  then is  $B = C$ ?

Let  $A, B$  be matrices of same dimension.  
Does  $AB = I$  imply  $BA = I$ , i.e.,  $A$  is invertible?

Let  $A, B$  be matrices of same dimension.

Does  $AB = I$  imply  $BA = I$ , i.e.,  $A$  is invertible?

We will see another characterization of invertible matrices using linear system, elementary matrices, linear map, etc.



Let  $A, B$  be matrices of same dimension.

Does  $AB = I$  imply  $BA = I$ , i.e.,  $A$  is invertible?

We will see another characterization of invertible matrices using linear system, elementary matrices, linear map, etc.

# Block-Partitioned Matrices

We can partition a matrix into smaller matrices called blocks such as

**Note:** There are many ways to partition given matrix.

# Block-Partitioned Matrices

We can partition a matrix into smaller matrices called blocks such as

**Note:** There are many ways to partition given matrix.

**Question:** What is the advantage of partitioning?

# Block-Partitioned Matrices

We can partition a matrix into smaller matrices called blocks such as

**Note:** There are many ways to partition given matrix.

**Question:** What is the advantage of partitioning?

$$M^2 = \begin{bmatrix} A^2 + BC & AB + BD \\ CA + DC & CB + D^2 \end{bmatrix}$$

## Block Matrix Inversion

$$\begin{bmatrix} A & B \\ C & D \end{bmatrix}^{-1} = \begin{bmatrix} A^{-1}(I + B(D - CA^{-1}B)^{-1}CA^{-1}) & -A^{-1}B(D - CA^{-1}B)^{-1} \\ -(D - CA^{-1}B)^{-1}CA^{-1} & (D - CA^{-1}B)^{-1} \end{bmatrix}$$

## Block Matrix Inversion

$$\begin{bmatrix} A & B \\ C & D \end{bmatrix}^{-1} = \begin{bmatrix} A^{-1}(I + B(D - CA^{-1}B)^{-1}CA^{-1}) & -A^{-1}B(D - CA^{-1}B)^{-1} \\ -(D - CA^{-1}B)^{-1}CA^{-1} & (D - CA^{-1}B)^{-1} \end{bmatrix}$$

A **block diagonal matrix** is a block matrix that is a square matrix such that the main diagonal blocks are square matrices and all off diagonal blocks are zero matrices. A block diagonal matrix and its inverse have the form:

$$A = \begin{bmatrix} A_1 & 0 & \cdots & 0 \\ 0 & A_2 & \cdots & 0 \\ \vdots & \vdots & & \vdots \\ 0 & 0 & \cdots & A_n \end{bmatrix} \quad A^{-1} = \begin{bmatrix} A_1^{-1} & 0 & \cdots & 0 \\ 0 & A_2^{-1} & \cdots & 0 \\ \vdots & \vdots & & \vdots \\ 0 & 0 & \cdots & A_n^{-1} \end{bmatrix}$$

## Block Matrix Inversion

$$\begin{bmatrix} A & B \\ C & D \end{bmatrix}^{-1} = \begin{bmatrix} A^{-1}(I + B(D - CA^{-1}B)^{-1}CA^{-1}) & -A^{-1}B(D - CA^{-1}B)^{-1} \\ -(D - CA^{-1}B)^{-1}CA^{-1} & (D - CA^{-1}B)^{-1} \end{bmatrix}$$

A **block diagonal matrix** is a block matrix that is a square matrix such that the main diagonal blocks are square matrices and all off diagonal blocks are zero matrices. A block diagonal matrix and its inverse have the form:

$$A = \begin{bmatrix} A_1 & 0 & \cdots & 0 \\ 0 & A_2 & \cdots & 0 \\ \vdots & \vdots & & \vdots \\ 0 & 0 & \cdots & A_n \end{bmatrix} \quad A^{-1} = \begin{bmatrix} A_1^{-1} & 0 & \cdots & 0 \\ 0 & A_2^{-1} & \cdots & 0 \\ \vdots & \vdots & & \vdots \\ 0 & 0 & \cdots & A_n^{-1} \end{bmatrix}$$

Block partition is useful in many computer science applications, VLSI chip design, the Strassen algorithm for fast matrix multiplication, coding theory.

# Linear System

## Definition

A linear System of  $m$  equations in  $n$  variables-  $X_1, X_2, \dots, X_n$  is

$$a_{11}X_1 + a_{12}X_2 + \cdots + a_{1n}X_n = b_1$$

$$a_{21}X_1 + a_{22}X_2 + \cdots + a_{2n}X_n = b_2$$

$$\vdots$$
$$\vdots$$
$$\vdots$$

$$a_{m1}X_1 + a_{m2}X_2 + \cdots + a_{mn}X_n = b_m$$

where  $a_{ij}, b_j \in \mathbb{R}$



# Linear System

## Definition

A linear System of  $m$  equations in  $n$  variables-  $X_1, X_2, \dots, X_n$  is

$$a_{11}X_1 + a_{12}X_2 + \cdots + a_{1n}X_n = b_1$$

$$a_{21}X_1 + a_{22}X_2 + \cdots + a_{2n}X_n = b_2$$

$$\vdots$$
$$\vdots$$
$$\vdots$$

$$a_{m1}X_1 + a_{m2}X_2 + \cdots + a_{mn}X_n = b_m$$

where  $a_{ij}, b_j \in \mathbb{R}$

Let's look at linear system of 2 equations in 2 variables:

Solve the system: (1)  $x + 2y = 3$ , (2)  $3x + y = 4$ .

**Elimination of variables:**

Eliminate  $x$  by  $(2) - 3 \times (1)$  to get  $y = 1$ .

**Cramer's Rule (determinant):**  $y = \frac{\begin{vmatrix} 1 & 3 \\ 3 & 4 \end{vmatrix}}{\begin{vmatrix} 1 & 2 \\ 3 & 1 \end{vmatrix}} = \frac{4-9}{1-6} = 1$

In either case, back substitution gives  $x = 1$

We could also solve for  $x$  first and use back substitution for  $y$ .

**Comparison:** For a large system, say 100 equations in 100 variables, elimination method is preferred, since computing the determinants of a 101 matrices of size  $100 \times 100$  is time-consuming.


# Geometry of Linear Equations

$$x+2y=3$$

and

$$3x+y=4$$

represent lines in  $\mathbb{R}^2$  passing through  $(0, 3/2)$  and  $(3, 0)$  and through  $(0, 4)$  and  $(4/3, 0)$  respectively.



The intersection of the two lines is the unique point  $(1, 1)$ .


Hence  $x = 1$  and  $y = 1$  is the solution of above system of linear equations.

# 3 Equations in 3 Variables

A linear equation in 3 variables represents a plane in a 3-dimensional space  $\mathbb{R}^3$ .

$$x + 2y + 3z = 6$$

passes through  $(0, 0, 2)$ ,  $(0, 3, 0)$ ,  $(6, 0, 0)$ .



$x + 2y + 3z = 12$  passes through  $(0, 0, 4)$ ,  $(0, 6, 0)$ ,  $(12, 0, 0)$ .  
which is parallel to above plane.

Suppose we have 3 equations in 3 variables, means we have to find intersection of 3 planes, say  $P_1, P_2, P_3$ .

Suppose we have 3 equations in 3 variables, means we have to find intersection of 3 planes, say  $P_1, P_2, P_3$ .

This is same as finding intersection of line  $L$  with  $P_3$  (Intersection of  $P_1, P_2$  is  $L$ , if  $P_1, P_2$  are not parallel).

Suppose we have 3 equations in 3 variables, means we have to find intersection of 3 planes, say  $P_1, P_2, P_3$ .

This is same as finding intersection of line  $L$  with  $P_3$  (Intersection of  $P_1, P_2$  is  $L$ , if  $P_1, P_2$  are not parallel).

- ▶ If the line  $L$  does intersects with the plane  $P_3$  , then the linear system has **unique** solution.

Suppose we have 3 equations in 3 variables, means we have to find intersection of 3 planes, say  $P_1, P_2, P_3$ .

This is same as finding intersection of line  $L$  with  $P_3$  (Intersection of  $P_1, P_2$  is  $L$ , if  $P_1, P_2$  are not parallel).

- ▶ If the line  $L$  does intersects with the plane  $P_3$  , then the linear system has **unique** solution.
- ▶ If the line  $L$  does not intersects with the plane  $P_3$  , then the linear system has **no** solution.



Suppose we have 3 equations in 3 variables, means we have to find intersection of 3 planes, say  $P_1, P_2, P_3$ .

This is same as finding intersection of line  $L$  with  $P_3$  (Intersection of  $P_1, P_2$  is  $L$ , if  $P_1, P_2$  are not parallel).

- ▶ If the line  $L$  does intersects with the plane  $P_3$  , then the linear system has **unique** solution.
- ▶ If the line  $L$  does not intersects with the plane  $P_3$  , then the linear system has **no** solution.
- ▶ If the line  $L$  is contained in the plane  $P_3$ , then the system has **infinitely many** solutions.

In this case, every point of  $L$  is a solution.

Suppose we have 3 equations in 3 variables, means we have to find intersection of 3 planes, say  $P_1, P_2, P_3$ .

This is same as finding intersection of line  $L$  with  $P_3$  (Intersection of  $P_1, P_2$  is  $L$ , if  $P_1, P_2$  are not parallel).

- ▶ If the line  $L$  does intersects with the plane  $P_3$  , then the linear system has **unique** solution.
- ▶ If the line  $L$  does not intersects with the plane  $P_3$  , then the linear system has **no** solution.
- ▶ If the line  $L$  is contained in the plane  $P_3$ , then the system has **infinitely many** solutions.

In this case, every point of  $L$  is a solution.

**Question:** Can we do the same when number of variables are  $> 3$ ?

# Gaussian Elimination: Unique solution

**Example:**  $2x + y + z = 5$ ,  $4x - 6y = -2$ ,  $-2x + 7y + 2z = 9$ .

**Algorithm:** Eliminate  $x$  from last 2 equations by  $(2) - 2(1)$ , and  $(3) + (1)$  to get the *equivalent system*:

$$2x + y + z = 5, \quad -8y - 2z = -12, \quad 8y + 3z = 14$$

The first *pivot* is 2, second pivot is  $-8$ . Eliminate  $y$  from the last equation to get an equivalent *triangular system*:

$$2x + y + z = 5, \quad -8y - 2z = -12, \quad z = 2$$

Solve this triangular system by *back substitution*, we get

$$z = 2, \quad y = 1, \quad x = 1$$

**Observe:** This is the only possible solution!

# Gaussian Elimination: No solution

**Example:**  $2x + y + z = 5$ ,  $4x - 6y = -2$ ,  $-2x + 7y + z = 9$ .

**Step 1** Eliminate  $x$  (using the 1st pivot 2) to get:

$$2x + y + z = 5, \quad -8y - 2z = -12, \quad 8y + 2z = 14$$

**Step 2**: Eliminate  $y$  (using the 2nd pivot -8) to get:

$$2x + y + z = 5, \quad -8y - 2z = -12, \quad 0 = 2.$$

The last equation shows that there is no solution, i.e., the system is *inconsistent*.

**Geometric reasoning:** In Step 1, notice we get two distinct parallel planes  $8y + 2z = 12$  and  $8y + 2z = 14$ .

They have no point in common.

**Note:** The planes in the original system were not parallel, but in an equivalent system, we get two distinct parallel planes!

# Gaussian Elimination: Infinitely solution

**Example:**  $2x + y + z = 5$ ,  $4x - 6y = -2$ ,  $-2x + 7y + z = 7$ .

**Step 1** Eliminate  $x$  (using the 1st pivot 2) to get:

$$2x + y + z = 5, \quad -8y - 2z = -12, \quad 8y + 2z = 12$$

**Step 2**: Eliminate  $y$  (using the 2nd pivot -8) to get:

$$2x + y + z = 5, \quad -8y - 2z = -12, \quad 0 = 0.$$

There are only two equations. For every value of  $z$ , values for  $x$  and  $y$  are obtained by back-substitution, e.g.  $(1, 1, 2)$  or  $(\frac{7}{4}, \frac{3}{2}, 0)$ . Hence the system has infinitely many solutions.

**Geometric reasoning:** In Step 1, notice we get two parallel planes  $-8y - 2z = 12$  and  $8y + 2z = 12$ .

They give the same plane. Hence we are looking at the intersection of the two planes,  $2x + y + z = 5$  and  $8x + 2z = 12$ , which is a line.

A system of linear equations has exactly one of following

1. no solution; (**inconsistent system**)
2. exactly one solution; (**consistent system**)
3. infinitely many solutions. (**consistent system**)

A system of linear equations has exactly one of following

1. no solution; (**inconsistent system**)
2. exactly one solution; (**consistent system**)
3. infinitely many solutions. (**consistent system**)

Solve