

Problems In Group Theory

Basu Dev Karki

Contents

1	Basic Group Theory	1
1.1	Introduction	1

Chapter 1

Basic Group Theory

1.1 Introduction

Problem 1.1.1. Determine which of the following binary operations are associative:

1. the operation on $\mathbb{Z}$ defined by $a * b = a - b$
2. the operation on $\mathbb{R}$ defined by $a * b = a + b + ab$
3. the operation on $\mathbb{Q}$ defined by $a * b = \frac{a+b}{5}$
4. the operation on $\mathbb{Z} \times \mathbb{Z}$ defined by $(a, b) * (c, d) = (ad + bc, bd)$
5. the operation on $\mathbb{Q} - \{0\}$ defined by $a * b = \frac{a}{b}$

Proof. This exercise is pretty trivial as we just have to compute $a * (b * c) = (a * b) * c$. For 1., $a * (b * c) = a * (b - c) = a - (b - c) = a - b + c$ but $(a * b) * c = (a - b) * c = (a - b) - c = a - b - c$. Rest of them are pretty similar. $\square$

Problem 1.1.2. Prove that addition of residue classes in $\mathbb{Z}/n\mathbb{Z}$ is associative.

Proof. By definition of residue addition,

$$[a] + [b] = [a + b]$$

Therefore, we get the following.

$$\begin{aligned} [a] + ([b] + [c]) &= [a] + [b + c] \\ &= [a + (b + c)] \\ &= [(a + b) + c] \\ &= [a + b] + [c] \\ &= ([a] + [b]) + [c] \end{aligned}$$

$\square$

Problem 1.1.3. Prove that multiplication of residue classes in $\mathbb{Z}/n\mathbb{Z}$ is associative.

Proof. By definition of residue multiplication,

$$[a] \cdot [b] = [a \cdot b]$$

Therefore, we get the following.

$$\begin{aligned} [a] \cdot ([b] \cdot [c]) &= [a] \cdot [b \cdot c] \\ &= [a \cdot (b \cdot c)] \\ &= [(a \cdot b) \cdot c] \\ &= [a \cdot b] \cdot [c] \\ &= ([a] \cdot [b]) \cdot [c] \end{aligned}$$

□

Problem 1.1.4. Prove for all $n > 1$ that $\mathbb{Z}/n\mathbb{Z}$ is not a group under multiplication of residue classes.

Proof. There is no multiplicative inverse for $[a]$ with $\gcd(a, n) > 1$. The proof for it pretty trivial, if there was a inverse, it would mean that

$$a \cdot b \equiv 1 \pmod{n} \implies 0 \equiv 1 \pmod{\gcd(a, n)}$$

That's a contradiction.

□

Problem 1.1.5. Determine which of the following sets are groups under addition:

1. the set of rational numbers (including $0 = 0/1$) in lowest terms whose denominators are odd
2. the set of rational numbers (including $0 = 0/1$) in lowest terms whose denominators are even
3. the set of rational numbers of absolute value < 1
4. the set of rational numbers of absolute value ≥ 1 together with 0
5. the set of rational numbers with denominators equal to 1 or 2
6. the set of rational numbers with denominators equal to 1, 2 or 3.

Proof. Below are the answers:

1. Group.
2. Not a group. Because it is not closed. (Eg: $5/2 + 7/2 = 6$)
3. Not a group. Because again it is not closed. (Eg : $0.99 + 0.98 = 1.97$)
4. Not a group. Because again it is not closed. (Eg : $4 + (-3.1) = 0.9$)
5. Group.
6. Not a group. Because its not closed. (Eg : $1/2 + 1/3 = 5/6$)

□

Problem 1.1.6. Let $G = \{x \in \mathbb{R} \mid 0 \leq x < 1\}$ and for $x, y \in G$ let $x * y$ be the fractional part of $x + y$ (i.e., $x * y = x + y - \lfloor x + y \rfloor$ where $\lfloor a \rfloor$ is the greatest integer less than or equal to a). Prove that $*$ is a well defined binary operation on G and that G is an abelian group under $*$ (called the real numbers mod 1).

Proof. It is well defined as,

$$\begin{aligned} x_1 = x_2, y_1 = y_2 &\implies x_1 * y_1 = x_1 + y_1 - \lfloor x_1 + y_1 \rfloor \\ &\implies x_1 + y_1 - \lfloor x_1 + y_1 \rfloor = x_2 + y_2 - \lfloor x_2 + y_2 \rfloor = x_2 * y_2 \end{aligned}$$

One can check that inverses and identity exists and its closed under addition. For associativity,

$$\begin{aligned} x * (y * z) &= x * (y + z - \lfloor y + z \rfloor) \\ &= \{x + y + z - \lfloor y + z \rfloor\} \\ &= \{x + y + z\} \\ &= \{x + y - \lfloor x + y \rfloor + z\} \\ &= (x * y) * z \end{aligned}$$

□