

# Econometrics III

## Assignment Part 3, 4, 5

### Tinbergen Insitute

Stanislav Avdeev  
590050sa  
stnavdeev@gmail.com

Bas Machielsens  
590049bm  
590049bm@eur.nl

March 29, 2021

## Question 1

### Part 1:


First, we plot the two time series:

```
df3 <- readr::read_csv("./data/data_assign_p3.csv")

twostocks <- c("apple", "netflix")


df3_twostocks <- df3 %>%
  janitor::clean_names() %>%
  pivot_longer(-date) %>%
  filter(is.element(name, twostocks)) %>%
  mutate(date = lubridate::dmy(date))

df3_twostocks %>%
  ggplot(aes(x = date, y = value,
             group = name, color = name)) + geom_line()
```



Then, we show the acf and pacf-functions:

```
n1 <- df3$NETFLIX %>%  
  acf(lag.max = 12, plot = F)  
  
n2 <- df3$NETFLIX %>%  
  pacf(lag.max = 12, plot = F)  
  
a1 <- df3$APPLE %>%  
  acf(lag.max = 12, plot = F)  
  
a2 <- df3$APPLE %>%  
  pacf(lag.max = 12, plot = F)  
  
par(mfrow=c(2,2))  
  
plot(n1, main = "Netflix"); plot(n2, main = "Netflix")  
plot(a1, main = "Apple"); plot(a2, main = "Apple")
```



The ACF's tell us that the stock price is highly dependent on the past stock price, and this dependence decays only very slowly: even the 50 or 100-period lag still shows significant autocorrelation.

## Part 2

We now implement a general to specific unit root test function:

```
unit_root_test <- function(column, order){  
  
  # Make the dataset  
  series <- ts(column)  
  first_differences <- diff(series, differences = 1)  
  laggedvar <- stats::lag(series, -1)  
  
  #other lagged first differences, delta x_{t-1}, ..., delta x_{t-p+1}  
  lagged_fds <- list()  
  
  for(i in 1:(order-1)){  
  
    lagged_fds[[i]] <- stats::lag(first_differences, k = -i)  
  
  }  
  
  df <- cbind(first_differences, laggedvar, purrr::reduce(lagged_fds, cbind)) %>%  
    as_tibble()  
  
  colnames(df) <- c("dxt", "xtm1", paste("dxtm", 1:(order-1), sep = ""))  
  
  df <- df %>%  
    na.omit()  
  
  # run the stepwise regression - find the best model  
  null = lm(data = df, formula = "dxt ~ xtm1")  
  full = lm(data = df, formula = paste("dxt ~ xtm1 +",  
                                       paste(paste("dxtm", 1:(order-1), sep = ""),  
                                             collapse = ' + '),  
                                       collapse = " "))  
  )  
  
  bestmodel <- step(full,  
    scope = list(lower = null, upper = full),  
    direction = "backward",  
    criterion = "BIC",  
    k = log(nrow(df)))  
  
  # perform the unit root test (MacKinnon, 2010)  
  b_critical <- -1.6156  
  
  t_value <- bestmodel %>%  
    summary() %>%  
    .$coefficients %>%  
    .[,3] %>%  
    .["xtm1"]  
  
  significant = abs(t_value) > abs(b_critical)  
  
  data.frame(stock = deparse(substitute(column)),  
    best_model = as.character(bestmodel$call[2]),  
    t_value = t_value,  
    sig = significant)  
}
```

```
summary_tests <- list()

for(i in 1:length(colnames(df3[, -1]))){
  df <- unit_root_test(df3[, i+1], 12)
  summary_tests[[i]] <- df %>%
    mutate(name = colnames(df3[, -1][i]))
}

summary_tests <- summary_tests %>%
  purrr::reduce(rbind) %>%
  select(-stock)

rownames(summary_tests) <- NULL

knitr::kable(summary_tests)
```

best_model	t_value	sig	name
dxt ~ xtm1 + dxtm1 + dxtm3 + dxtm7	-0.6984454	FALSE	APPLE
dxt ~ xtm1 + dxtm1 + dxtm2	-2.0475011	TRUE	EXXON_MOBIL
dxt ~ xtm1	-1.1619768	FALSE	FORD
dxt ~ xtm1 + dxtm1 + dxtm3	-1.5455395	FALSE	GEN_ELECTRIC
dxt ~ xtm1	-2.2598273	TRUE	INTEL
dxt ~ xtm1	-2.3999971	TRUE	MICROSOFT
dxt ~ xtm1 + dxtm2	-1.0642387	FALSE	NETFLIX
dxt ~ xtm1 + dxtm2 + dxtm3	-0.6373731	FALSE	NOKIA
dxt ~ xtm1 + dxtm1	-1.3045653	FALSE	SP500
dxt ~ xtm1 + dxtm1 + dxtm6 + dxtm11	-2.6283893	TRUE	YAHOO

The test seems to be significant for a number of stocks, indicating that for these stocks, the null hypothesis of a unit root is rejected in favor of stationarity. With a 10%  $\alpha$ -level, we expect to see a type I-error (rejecting the null while it is true) about one tenth of the time for every test. This means that the probability of having at least 1 type-I error is very large:  $(1 - 0.9^{10}) = 0.6513216$ . It would be better to use some kind of Bonferroni correction to correct for these compounding type I-errors, but alternatively, we could also lower the  $\alpha$ -level.

### Part 3

The forecast  $\mathbb{E}[P_{t+1}|D_t] = \mathbb{E}[P_{t+1}|P_t] = \mathbb{E}[P_t + \epsilon_t] = P_t$ . Similarly, the forecast  $\mathbb{E}[P_{t+2}] = \mathbb{E}[P_{t+1} + \epsilon_{t+1}] = \mathbb{E}[P_{t+1}] = P_t$ . Generalizing this pattern, the forecast for  $P_{t+h} = P_t$ . The variance of the forecast is derived using the distribution:

$$\begin{aligned}
 P_{t+1} &= P_t + \epsilon_t \Rightarrow P_{t+1}|P_t \sim N(P_t, \sigma^2) \\
 P_{t+2} &= P_{t+1} + \epsilon_{t+1} = \\
 &= P_t + \epsilon_t + \epsilon_{t+1} \Rightarrow P_{t+2}|P_t \sim N(P_t, 2\sigma^2)
 \end{aligned}$$

Generalizing this pattern, we can see that  $\text{Var}(P_{t+h}) = h \cdot \sigma^2$ . Hence, we can implement our forecasts in the following way:

```
#forecast code
p_tplush <- df3 %>%
  slice_tail(n=1) %>%
```

```

select(c("APPLE", "MICROSOFT"))

var_apple <- lm(data = df3 %>%
  select("APPLE"),
  formula = APPLE ~ lag(APPLE, 1)) %>%
  .$residuals %>%
  var()


var_microsoft <- lm(data = df3 %>%
  select("MICROSOFT"),
  formula = MICROSOFT ~ lag(MICROSOFT, 1)) %>%
  .$residuals %>%
  var()

forecasts_apple <- data.frame(time = 1:5,
  value = rep(p_tplush %>%
    pull(1), 5),
  var = var_apple * 1:5)

forecasts_microsoft <- data.frame(time = 1:5,
  value = rep(p_tplush %>%
    pull(2), 5),
  var = var_microsoft * 1:5)

forecasts_apple %>%
  ggplot(aes(x = time)) +
  geom_line(aes(y = value)) +
  geom_line(aes(y = value + 1.96*sqrt(var)), lty = "dashed") +
  geom_line(aes(y = value - 1.96*sqrt(var)), lty = "dashed")


```



```

forecasts_microsoft %>%
  ggplot(aes(x = time)) +
  geom_line(aes(y = value)) +
  geom_line(aes(y = value + 1.96*sqrt(var)), lty = "dashed") +
  geom_line(aes(y = value - 1.96*sqrt(var)), lty = "dashed")

```



Hence, there is no investment advice that we can give: the value is predicted to remain constant, and for all predictions, the probability of an increase in stock price equals the probability of a decrease in stock price. The expected value of any investment strategy is 0, and the value is neither expected to increase, nor to decrease.

#### Part 4

*Do you find a statistically significant contemporaneous relation between Microsoft and Exxon Mobile stock prices?*

```
lm(df3, formula = MICROSOFT ~ EXXON_MOBIL) %>%
  stargazer(header = F, omit.stat = c("adj.rsq", "ser", "f"))
```

Table 2:

<i>Dependent variable:</i>	
MICROSOFT	
EXXON_MOBIL	0.203*** (0.009)
Constant	11.245*** (0.713)
Observations	1,499
R <sup>2</sup>	0.251
<i>Note:</i> *p<0.1; **p<0.05; ***p<0.01	

*Do you agree that changes in Microsoft stock prices are largely explained by fluctuations in the stock price of Exxon Mobile?*

We do not agree that changes in Microsoft stock prices are largely explained by fluctuations in the stock price of Exxon Mobile. It is likely that these results are driven by a shared stochastic trend in both of variables, in other words, the variables might be cointegrated. It can be shown that if two variables share a stochastic trend, the t-value of the estimated coefficient tends to infinity, and the probability of obtaining statistical significance to 1, even under the assumption of completely unrelated trends.

## Question 4


```
df4 <- read_csv("data/data_assign_p4.csv")

df4$obs <- ts(df4$obs,
             frequency = 4,
             start = c(1988, 1))

p_cons <- ggplot(data = df4, aes(x = as.Date(obs), y = CONS)) +
  geom_line() +
  scale_x_date(date_labels = "%b %Y")

p_inc <- ggplot(data = df4, aes(x = as.Date(obs), y = INC)) +
  geom_line() +
  scale_x_date(date_labels = "%b %Y")

cowplot::plot_grid(p_cons, p_inc)
```



```
inc <- ts(df4$INC,
         frequency = 4,
         start = c(1988, 1))

cons <- ts(df4$CONS,
         frequency = 4,
         start = c(1988, 1))

inc_acf <- acf(inc, plot = F); inc_pcf <- pacf(inc, plot = F)
cons_acf <- acf(cons, plot = F); cons_pcf <- pacf(cons, plot = F)

par(mfrow=c(2,2))
plot(inc_acf); plot(inc_pcf); plot(cons_acf); plot(cons_pcf)
```

