
Confidence intervals: coverage [Wilson 95% interval] and mean width w


small centroid gap / homoscedastic / balanced / rho = (-0.8, -0.8)



Bar: mean lower to mean upper bound over the simulated bulks; point: mean estimate; dashed: mean target.
Methods without intervals are not shown.

Confidence intervals: coverage [Wilson 95% interval] and mean width w


small centroid gap / homoscedastic / balanced / rho = (+0.0, +0.0)



Bar: mean lower to mean upper bound over the simulated bulks; point: mean estimate; dashed: mean target.
Methods without intervals are not shown.

Confidence intervals: coverage [Wilson 95% interval] and mean width w


small centroid gap / homoscedastic / balanced / rho = (-0.8, +0.8)



Bar: mean lower to mean upper bound over the simulated bulks; point: mean estimate; dashed: mean target.
Methods without intervals are not shown.

Confidence intervals: coverage [Wilson 95% interval] and mean width w

small centroid gap / homoscedastic / balanced / rho = (+0.8, +0.8)




Bar: mean lower to mean upper bound over the simulated bulks; point: mean estimate; dashed: mean target.
Methods without intervals are not shown.

Confidence intervals: coverage [Wilson 95% interval] and mean width w

small centroid gap / homoscedastic / moderately unbalanced / rho = (-0.8, -0.8)


```
Wald (delta)            0.94 [ 0.9, 0.97]   w =  1.3
ILR, mapped back           1 [0.98,    1]   w = 0.929
Profile likelihood      0.98 [0.95, 0.99]   w = 0.828
Oracle (Fisher at p*)   0.94 [ 0.9, 0.97]   w = 1.27
Wald (delta)            0.95 [0.91, 0.97]   w = 1.29
ILR, mapped back           1 [0.98,    1]   w = 0.913
Wald (delta)               1 [0.98,    1]   w = 1.28
ILR, mapped back           1 [0.98,    1]   w = 0.874
```



Bar: mean lower to mean upper bound over the simulated bulks; point: mean estimate; dashed: mean target.
Methods without intervals are not shown.

Confidence intervals: coverage [Wilson 95% interval] and mean width w


small centroid gap / homoscedastic / moderately unbalanced / rho = (+0.0, +0.0)



Bar: mean lower to mean upper bound over the simulated bulks; point: mean estimate; dashed: mean target.
Methods without intervals are not shown.

Confidence intervals: coverage [Wilson 95% interval] and mean width w


small centroid gap / homoscedastic / moderately unbalanced / rho = (-0.8, +0.8)



Bar: mean lower to mean upper bound over the simulated bulks; point: mean estimate; dashed: mean target.
Methods without intervals are not shown.

Confidence intervals: coverage [Wilson 95% interval] and mean width w


small centroid gap / homoscedastic / moderately unbalanced / rho = (+0.8, +0.8)



Bar: mean lower to mean upper bound over the simulated bulks; point: mean estimate; dashed: mean target.
Methods without intervals are not shown.

Confidence intervals: coverage [Wilson 95% interval] and mean width w


small centroid gap / homoscedastic / highly unbalanced / rho = (-0.8, -0.8)



Bar: mean lower to mean upper bound over the simulated bulks; point: mean estimate; dashed: mean target.
Methods without intervals are not shown.

Confidence intervals: coverage [Wilson 95% interval] and mean width w

small centroid gap / homoscedastic / highly unbalanced / rho = (+0.0, +0.0)



Bar: mean lower to mean upper bound over the simulated bulks; point: mean estimate; dashed: mean target.
Methods without intervals are not shown.

Confidence intervals: coverage [Wilson 95% interval] and mean width w

small centroid gap / homoscedastic / highly unbalanced / rho = (-0.8, +0.8)

Wald (delta) 0.84 [0.79, 0.89] w = 1.15

ILR, mapped back 0.64 [0.57, 0.7] w = 0.884

Profile likelihood 0.96 [0.92, 0.98] w = 0.504


Oracle (Fisher at p*) 0.95 [0.91, 0.97] w = 1.36

Wald (delta) 0.78 [0.7, 0.84] w = 1.06

ILR, mapped back 0.46 [0.38, 0.55] w = 0.829

Wald (delta) 0.7 [0.62, 0.77] w = 0.875

ILR, mapped back 0.0074 [0.0013, 0.04] w = 0.717




Bar: mean lower to mean upper bound over the simulated bulks; point: mean estimate; dashed: mean target.
Methods without intervals are not shown.

Confidence intervals: coverage [Wilson 95% interval] and mean width w

small centroid gap / homoscedastic / highly unbalanced / rho = (+0.8, +0.8)

```
Wald (delta)            0.94 [ 0.9, 0.97]   w = 0.556
ILR, mapped back        0.74 [0.67, 0.79]   w = 0.819
Profile likelihood      0.96 [0.92, 0.98]   w = 0.273
Oracle (Fisher at p*)   0.96 [0.93, 0.98]   w = 0.586
Wald (delta)            0.91 [0.84, 0.95]   w = 0.532
ILR, mapped back        0.55 [0.47, 0.64]   w = 0.697
Wald (delta)            0.85 [0.77,  0.9]   w = 0.501
ILR, mapped back          0  [3.5e-18, 0.031]   w = 0.51
```




Bar: mean lower to mean upper bound over the simulated bulks; point: mean estimate; dashed: mean target.
Methods without intervals are not shown.

Confidence intervals: coverage [Wilson 95% interval] and mean width w

small centroid gap / heteroscedastic / balanced / rho = (-0.8, -0.8)

```
Wald (delta)              1  [0.98,    1]   w = 1.34
ILR, mapped back          1  [0.98,    1]   w = 0.906
Profile likelihood     0.96 [0.93, 0.98]   w = 0.943
Oracle (Fisher at p*)     1  [0.98,    1]   w = 1.37
Wald (delta)              1  [0.98,    1]   w = 1.34
ILR, mapped back          1  [0.98,    1]   w = 0.899
Wald (delta)              1  [0.98,    1]   w = 1.34
ILR, mapped back          1  [0.98,    1]   w = 0.882
```



Bar: mean lower to mean upper bound over the simulated bulks; point: mean estimate; dashed: mean target.
Methods without intervals are not shown.


Confidence intervals: coverage [Wilson 95% interval] and mean width w

small centroid gap / heteroscedastic / balanced / rho = (+0.0, +0.0)

```
Wald (delta)              1  [0.98,    1]   w = 1.14
ILR, mapped back          1  [0.98,    1]   w = 0.847
Profile likelihood     0.98 [0.94, 0.99]   w = 0.908
Oracle (Fisher at p*)     1  [0.98,    1]   w = 1.13

Wald (delta)              1  [0.98,    1]   w = 1.13
ILR, mapped back          1  [0.98,    1]   w = 0.842


Wald (delta)              1  [0.98,    1]   w = 1.13
ILR, mapped back          1  [0.98,    1]   w = 0.819
```



Bar: mean lower to mean upper bound over the simulated bulks; point: mean estimate; dashed: mean target.
Methods without intervals are not shown.

Confidence intervals: coverage [Wilson 95% interval] and mean width w


small centroid gap / heteroscedastic / balanced / rho = (-0.8, +0.8)



Bar: mean lower to mean upper bound over the simulated bulks; point: mean estimate; dashed: mean target.
Methods without intervals are not shown.

Confidence intervals: coverage [Wilson 95% interval] and mean width w


small centroid gap / heteroscedastic / balanced / rho = (+0.8, +0.8)



Bar: mean lower to mean upper bound over the simulated bulks; point: mean estimate; dashed: mean target.
Methods without intervals are not shown.

Confidence intervals: coverage [Wilson 95% interval] and mean width w


small centroid gap / heteroscedastic / moderately unbalanced / rho = (-0.8, -0.8)



Bar: mean lower to mean upper bound over the simulated bulks; point: mean estimate; dashed: mean target.
Methods without intervals are not shown.

Confidence intervals: coverage [Wilson 95% interval] and mean width w


small centroid gap / heteroscedastic / moderately unbalanced / rho = (+0.0, +0.0)



Bar: mean lower to mean upper bound over the simulated bulks; point: mean estimate; dashed: mean target.
Methods without intervals are not shown.

Confidence intervals: coverage [Wilson 95% interval] and mean width w


small centroid gap / heteroscedastic / moderately unbalanced / rho = (-0.8, +0.8)



Bar: mean lower to mean upper bound over the simulated bulks; point: mean estimate; dashed: mean target.
Methods without intervals are not shown.

Confidence intervals: coverage [Wilson 95% interval] and mean width w


small centroid gap / heteroscedastic / moderately unbalanced / rho = (+0.8, +0.8)



Bar: mean lower to mean upper bound over the simulated bulks; point: mean estimate; dashed: mean target.
Methods without intervals are not shown.

Confidence intervals: coverage [Wilson 95% interval] and mean width w

small centroid gap / heteroscedastic / highly unbalanced / rho = (-0.8, -0.8)




Bar: mean lower to mean upper bound over the simulated bulks; point: mean estimate; dashed: mean target.
Methods without intervals are not shown.

Confidence intervals: coverage [Wilson 95% interval] and mean width w

small centroid gap / heteroscedastic / highly unbalanced / rho = (+0.0, +0.0)


```
Wald (delta)            0.94 [0.89, 0.96]   w = 1.18
ILR, mapped back        0.64 [0.58, 0.71]   w = 0.933
Profile likelihood      0.97 [0.94, 0.99]   w = 0.634
Oracle (Fisher at p*)   0.94 [ 0.9, 0.97]   w = 1.23
Wald (delta)            0.92 [0.86, 0.96]   w = 1.15
ILR, mapped back        0.41 [0.32,  0.5]   w = 0.889
Wald (delta)            0.94 [0.88, 0.97]   w = 1.13
ILR, mapped back        0.05 [0.023,  0.1]   w = 0.831
```



Bar: mean lower to mean upper bound over the simulated bulks; point: mean estimate; dashed: mean target.
Methods without intervals are not shown.

Confidence intervals: coverage [Wilson 95% interval] and mean width w

small centroid gap / heteroscedastic / highly unbalanced / rho = (-0.8, +0.8)




Bar: mean lower to mean upper bound over the simulated bulks; point: mean estimate; dashed: mean target.
Methods without intervals are not shown.

Confidence intervals: coverage [Wilson 95% interval] and mean width w

small centroid gap / heteroscedastic / highly unbalanced / rho = (+0.8, +0.8)


```
Wald (delta)            0.9  [0.85, 0.93]   w = 0.635
ILR, mapped back        0.7  [0.63, 0.75]   w = 0.837
Profile likelihood      0.96 [0.92, 0.98]   w = 0.316
Oracle (Fisher at p*)   0.96 [0.92, 0.98]   w = 0.674
Wald (delta)            0.83 [0.75, 0.89]   w = 0.604
ILR, mapped back        0.48 [0.39, 0.57]   w = 0.724
Wald (delta)            0.75 [0.66, 0.82]   w = 0.569
ILR, mapped back          0  [   0, 0.032]   w = 0.558
```



Bar: mean lower to mean upper bound over the simulated bulks; point: mean estimate; dashed: mean target.
Methods without intervals are not shown.

Confidence intervals: coverage [Wilson 95% interval] and mean width w


large centroid gap / homoscedastic / balanced / rho = (-0.8, -0.8)



Bar: mean lower to mean upper bound over the simulated bulks; point: mean estimate; dashed: mean target.
Methods without intervals are not shown.

Confidence intervals: coverage [Wilson 95% interval] and mean width w


large centroid gap / homoscedastic / balanced / rho = (+0.0, +0.0)



Bar: mean lower to mean upper bound over the simulated bulks; point: mean estimate; dashed: mean target.
Methods without intervals are not shown.

Confidence intervals: coverage [Wilson 95% interval] and mean width w


large centroid gap / homoscedastic / balanced / rho = (-0.8, +0.8)



Bar: mean lower to mean upper bound over the simulated bulks; point: mean estimate; dashed: mean target.
Methods without intervals are not shown.

Confidence intervals: coverage [Wilson 95% interval] and mean width w


large centroid gap / homoscedastic / balanced / rho = (+0.8, +0.8)



Bar: mean lower to mean upper bound over the simulated bulks; point: mean estimate; dashed: mean target.
Methods without intervals are not shown.

Confidence intervals: coverage [Wilson 95% interval] and mean width w


large centroid gap / homoscedastic / moderately unbalanced / rho = (-0.8, -0.8)



Bar: mean lower to mean upper bound over the simulated bulks; point: mean estimate; dashed: mean target.
Methods without intervals are not shown.

Confidence intervals: coverage [Wilson 95% interval] and mean width w


large centroid gap / homoscedastic / moderately unbalanced / rho = (+0.0, +0.0)



Bar: mean lower to mean upper bound over the simulated bulks; point: mean estimate; dashed: mean target.
Methods without intervals are not shown.

Confidence intervals: coverage [Wilson 95% interval] and mean width w


large centroid gap / homoscedastic / moderately unbalanced / rho = (-0.8, +0.8)



Bar: mean lower to mean upper bound over the simulated bulks; point: mean estimate; dashed: mean target.
Methods without intervals are not shown.

Confidence intervals: coverage [Wilson 95% interval] and mean width w


large centroid gap / homoscedastic / moderately unbalanced / rho = (+0.8, +0.8)



Bar: mean lower to mean upper bound over the simulated bulks; point: mean estimate; dashed: mean target.
Methods without intervals are not shown.

Confidence intervals: coverage [Wilson 95% interval] and mean width w


large centroid gap / homoscedastic / highly unbalanced / rho = (-0.8, -0.8)



Bar: mean lower to mean upper bound over the simulated bulks; point: mean estimate; dashed: mean target.
Methods without intervals are not shown.

Confidence intervals: coverage [Wilson 95% interval] and mean width w


large centroid gap / homoscedastic / highly unbalanced / rho = (+0.0, +0.0)



Bar: mean lower to mean upper bound over the simulated bulks; point: mean estimate; dashed: mean target.
Methods without intervals are not shown.

Confidence intervals: coverage [Wilson 95% interval] and mean width w


large centroid gap / homoscedastic / highly unbalanced / rho = (-0.8, +0.8)



Bar: mean lower to mean upper bound over the simulated bulks; point: mean estimate; dashed: mean target.
Methods without intervals are not shown.

Confidence intervals: coverage [Wilson 95% interval] and mean width w


large centroid gap / homoscedastic / highly unbalanced / rho = (+0.8, +0.8)



Bar: mean lower to mean upper bound over the simulated bulks; point: mean estimate; dashed: mean target.
Methods without intervals are not shown.

Confidence intervals: coverage [Wilson 95% interval] and mean width w


large centroid gap / heteroscedastic / balanced / rho = (-0.8, -0.8)



Bar: mean lower to mean upper bound over the simulated bulks; point: mean estimate; dashed: mean target.
Methods without intervals are not shown.

Confidence intervals: coverage [Wilson 95% interval] and mean width w


large centroid gap / heteroscedastic / balanced / rho = (+0.0, +0.0)



Bar: mean lower to mean upper bound over the simulated bulks; point: mean estimate; dashed: mean target.
Methods without intervals are not shown.

Confidence intervals: coverage [Wilson 95% interval] and mean width w


large centroid gap / heteroscedastic / balanced / rho = (-0.8, +0.8)



Bar: mean lower to mean upper bound over the simulated bulks; point: mean estimate; dashed: mean target.
Methods without intervals are not shown.

Confidence intervals: coverage [Wilson 95% interval] and mean width w


large centroid gap / heteroscedastic / balanced / rho = (+0.8, +0.8)



Bar: mean lower to mean upper bound over the simulated bulks; point: mean estimate; dashed: mean target.
Methods without intervals are not shown.

Confidence intervals: coverage [Wilson 95% interval] and mean width w


large centroid gap / heteroscedastic / moderately unbalanced / rho = (-0.8, -0.8)



Bar: mean lower to mean upper bound over the simulated bulks; point: mean estimate; dashed: mean target.
Methods without intervals are not shown.

Confidence intervals: coverage [Wilson 95% interval] and mean width w


large centroid gap / heteroscedastic / moderately unbalanced / rho = (+0.0, +0.0)



Bar: mean lower to mean upper bound over the simulated bulks; point: mean estimate; dashed: mean target.
Methods without intervals are not shown.

Confidence intervals: coverage [Wilson 95% interval] and mean width w


large centroid gap / heteroscedastic / moderately unbalanced / rho = (-0.8, +0.8)



Bar: mean lower to mean upper bound over the simulated bulks; point: mean estimate; dashed: mean target.
Methods without intervals are not shown.

Confidence intervals: coverage [Wilson 95% interval] and mean width w


large centroid gap / heteroscedastic / moderately unbalanced / rho = (+0.8, +0.8)



Bar: mean lower to mean upper bound over the simulated bulks; point: mean estimate; dashed: mean target.
Methods without intervals are not shown.

Confidence intervals: coverage [Wilson 95% interval] and mean width w


large centroid gap / heteroscedastic / highly unbalanced / rho = (-0.8, -0.8)



Bar: mean lower to mean upper bound over the simulated bulks; point: mean estimate; dashed: mean target.
Methods without intervals are not shown.

Confidence intervals: coverage [Wilson 95% interval] and mean width w


large centroid gap / heteroscedastic / highly unbalanced / rho = (+0.0, +0.0)



Bar: mean lower to mean upper bound over the simulated bulks; point: mean estimate; dashed: mean target.
Methods without intervals are not shown.

Confidence intervals: coverage [Wilson 95% interval] and mean width w

large centroid gap / heteroscedastic / highly unbalanced / rho = (-0.8, +0.8)




Bar: mean lower to mean upper bound over the simulated bulks; point: mean estimate; dashed: mean target.
Methods without intervals are not shown.

Confidence intervals: coverage [Wilson 95% interval] and mean width w

large centroid gap / heteroscedastic / highly unbalanced / rho = (+0.8, +0.8)

```
Wald (delta)            0.98 [0.96, 0.99]   w = 0.0715
ILR, mapped back        0.89 [0.84, 0.93]   w = 0.49
Profile likelihood      0.98 [0.96, 0.99]   w = 0.045
Oracle (Fisher at p*)   0.98 [0.96, 0.99]   w = 0.0717
Wald (delta)            0.98 [0.94, 0.99]   w = 0.0712
ILR, mapped back        0.85 [0.78,  0.9]   w = 0.297
Wald (delta)            0.92 [0.87, 0.96]   w = 0.0703
ILR, mapped back        0.68 [ 0.6, 0.75]   w = 0.0897
```



Bar: mean lower to mean upper bound over the simulated bulks; point: mean estimate; dashed: mean target.
Methods without intervals are not shown.