



balanced / low OVL




$W = l_z(p)^{1/2}$ ($\hat{z} - z^*$) versus $N(0, 1)$; identity line. Tail-sensitive band (Aldor-Noiman et al.) with $\mu = 0$, $\sigma = 1$. LBFGS, Marquardt-Levenberg and Newton-Raphson only; not detrended.

Whitened ILR Monte Carlo quantiles




balanced / high OVL



$W = L_z(p)^{1/2} (\hat{z} - z^*)$ versus $N(0, 1)$; identity line. Tail-sensitive band (Aldor-Noiman et al.) with $\mu = 0$, $\sigma = 1$. LBFGS, Marquardt-Levenberg and Newton-Raphson only; not detrended.


moderately unbalanced / low OVL



Algorithm LBFGS Marquardt-Levenberg Newton-Raphson


$W = L_z(p)^{1/2} (\hat{z} - z^*)$ versus $N(0, 1)$; identity line. Tail-sensitive band (Aldor-Noiman et al.) with $\mu = 0$, $\sigma = 1$. LBFGS, Marquardt-Levenberg and Newton-Raphson only; not detrended.

moderately unbalanced / moderate OVL




$W = L_z(p^*)^{1/2} (\hat{z} - z^*)$ versus $N(0, 1)$; identity line. Tail-sensitive band (Aldor-Noiman et al.) with $\mu = 0$, $\sigma = 1$. LBFGS, Marquardt-Levenberg and Newton-Raphson only; not detrended.


moderately unbalanced / high OVL



highly unbalanced / low OVL




highly unbalanced / moderate OVL



$W = I_z(p)^{1/2} (\hat{z} - z^*)$ versus $N(0, 1)$; identity line. Tail-sensitive band (Aldor-Noiman et al.) with $\mu = 0$, $\sigma = 1$. LBFGS, Marquardt-Levenberg and Newton-Raphson only; not detrended.

highly unbalanced / high OVL



$W = I_z(p^*)^{1/2} (\hat{z} - z^*)$ versus $N(0, 1)$; identity line. Tail-sensitive band (Aldor-Noiman et al.) with $\mu = 0$, $\sigma = 1$. LBFGS, Marquardt-Levenberg and Newton-Raphson only; not detrended.