
Wald approximation: whitened ILR estimates against N(0, 1)

small centroid gap / homoscedastic / balanced




ILR coordinate ● z1 ● Converged, local maximum ● Converged, not a local maximum Numerical failure

I_z(p*)^(1/2) (z-hat - z*), information at the true composition; estimates on a face have no ILR coordinates and are not shown.
Assesses the Wald approximation only; profile and bootstrap intervals do not rely on it.

Wald approximation: whitened ILR estimates against N(0, 1)

small centroid gap / homoscedastic / moderately unbalanced




ILR coordinate ● z1 ● Converged, local maximum ● Converged, not a local maximum Numerical failure

I_z(p*)^(1/2) (z-hat - z*), information at the true composition; estimates on a face have no ILR coordinates and are not shown.
Assesses the Wald approximation only; profile and bootstrap intervals do not rely on it.

Wald approximation: whitened ILR estimates against N(0, 1)

small centroid gap / homoscedastic / highly unbalanced




ILR coordinate ● z1 ● Converged, local maximum ● Converged, not a local maximum Numerical failure

I_z(p*)^(1/2) (z-hat - z*), information at the true composition; estimates on a face have no ILR coordinates and are not shown.
Assesses the Wald approximation only; profile and bootstrap intervals do not rely on it.

Wald approximation: whitened ILR estimates against N(0, 1)

small centroid gap / heteroscedastic / balanced




ILR coordinate ● z1 ● Converged, local maximum ● Converged, not a local maximum Numerical failure

I_z(p*)^(1/2) (z-hat - z*), information at the true composition; estimates on a face have no ILR coordinates and are not shown.
Assesses the Wald approximation only; profile and bootstrap intervals do not rely on it.

Wald approximation: whitened ILR estimates against N(0, 1)

small centroid gap / heteroscedastic / moderately unbalanced




ILR coordinate ● z1 ● Converged, local maximum ● Converged, not a local maximum Numerical failure

I_z(p*)^(1/2) (z-hat - z*), information at the true composition; estimates on a face have no ILR coordinates and are not shown.
Assesses the Wald approximation only; profile and bootstrap intervals do not rely on it.

Wald approximation: whitened ILR estimates against N(0, 1)

small centroid gap / heteroscedastic / highly unbalanced




ILR coordinate ● z1 ● Converged, local maximum ● Converged, not a local maximum Numerical failure

I_z(p*)^(1/2) (z-hat - z*), information at the true composition; estimates on a face have no ILR coordinates and are not shown.
Assesses the Wald approximation only; profile and bootstrap intervals do not rely on it.

Wald approximation: whitened ILR estimates against N(0, 1)

large centroid gap / homoscedastic / balanced




ILR coordinate ● z1 ● Converged, local maximum ● Converged, not a local maximum ● Numerical failure

I_z(p*)^(1/2) (z-hat - z*), information at the true composition; estimates on a face have no ILR coordinates and are not shown.
Assesses the Wald approximation only; profile and bootstrap intervals do not rely on it.

Wald approximation: whitened ILR estimates against N(0, 1)

large centroid gap / homoscedastic / moderately unbalanced




ILR coordinate ● z1 ● Converged, local maximum ● Converged, not a local maximum Numerical failure

I_z(p*)^(1/2) (z-hat - z*), information at the true composition; estimates on a face have no ILR coordinates and are not shown.
Assesses the Wald approximation only; profile and bootstrap intervals do not rely on it.

Wald approximation: whitened ILR estimates against N(0, 1)

large centroid gap / homoscedastic / highly unbalanced




ILR coordinate ● z1 ● Converged, local maximum ● Converged, not a local maximum Numerical failure

I_z(p*)^(1/2) (z-hat - z*), information at the true composition; estimates on a face have no ILR coordinates and are not shown.
Assesses the Wald approximation only; profile and bootstrap intervals do not rely on it.

Wald approximation: whitened ILR estimates against N(0, 1)

large centroid gap / heteroscedastic / balanced




ILR coordinate ● z1 ● Converged, local maximum Converged, not a local maximum Numerical failure

I_z(p*)^(1/2) (z-hat - z*), information at the true composition; estimates on a face have no ILR coordinates and are not shown.
Assesses the Wald approximation only; profile and bootstrap intervals do not rely on it.

Wald approximation: whitened ILR estimates against N(0, 1)

large centroid gap / heteroscedastic / moderately unbalanced




ILR coordinate ● z1 ● Converged, local maximum ● Converged, not a local maximum Numerical failure

I_z(p*)^(1/2) (z-hat - z*), information at the true composition; estimates on a face have no ILR coordinates and are not shown.
Assesses the Wald approximation only; profile and bootstrap intervals do not rely on it.

Wald approximation: whitened ILR estimates against N(0, 1)

large centroid gap / heteroscedastic / highly unbalanced



ILR coordinate ● z1 ● Converged, local maximum ● Converged, not a local maximum Numerical failure

I_z(p*)^(1/2) (z-hat - z*), information at the true composition; estimates on a face have no ILR coordinates and are not shown.
Assesses the Wald approximation only; profile and bootstrap intervals do not rely on it.