
Purified profiles in two latent directions


low overlap / CT1 Scale-free / CT2 Scale-free



Equal numbers of draws per type; ellipses: 95% normal regions. PCA uses the pooled variance, MclustDR (EDDA, VVV) the class means and covariances.

Purified profiles in two latent directions


low overlap / CT1 Scale-free / CT2 SBM



Equal numbers of draws per type; ellipses: 95% normal regions. PCA uses the pooled variance, MclustDR (EDDA, VVV) the class means and covariances.

Purified profiles in two latent directions


low overlap / CT1 SBM / CT2 Scale-free



Equal numbers of draws per type; ellipses: 95% normal regions. PCA uses the pooled variance, MclustDR (EDDA, VVV) the class means and covariances.

Purified profiles in two latent directions


low overlap / CT1 SBM / CT2 SBM



Equal numbers of draws per type; ellipses: 95% normal regions. PCA uses the pooled variance, MclustDR (EDDA, VVV) the class means and covariances.

Purified profiles in two latent directions


moderate overlap / CT1 Scale-free / CT2 Scale-free



Equal numbers of draws per type; ellipses: 95% normal regions. PCA uses the pooled variance, MclustDR (EDDA, VVV) the class means and covariances.

Purified profiles in two latent directions


moderate overlap / CT1 Scale-free / CT2 SBM



Equal numbers of draws per type; ellipses: 95% normal regions. PCA uses the pooled variance, MclustDR (EDDA, VVV) the class means and covariances.

Purified profiles in two latent directions


moderate overlap / CT1 SBM / CT2 Scale-free



Equal numbers of draws per type; ellipses: 95% normal regions. PCA uses the pooled variance, MclustDR (EDDA, VVV) the class means and covariances.

Purified profiles in two latent directions


moderate overlap / CT1 SBM / CT2 SBM



Equal numbers of draws per type; ellipses: 95% normal regions. PCA uses the pooled variance, MclustDR (EDDA, VVV) the class means and covariances.

Purified profiles in two latent directions


high overlap / CT1 Scale-free / CT2 Scale-free



Equal numbers of draws per type; ellipses: 95% normal regions. PCA uses the pooled variance, MclustDR (EDDA, VVV) the class means and covariances.

Purified profiles in two latent directions


high overlap / CT1 Scale-free / CT2 SBM



Equal numbers of draws per type; ellipses: 95% normal regions. PCA uses the pooled variance, MclustDR (EDDA, VVV) the class means and covariances.

Purified profiles in two latent directions


high overlap / CT1 SBM / CT2 Scale-free



Equal numbers of draws per type; ellipses: 95% normal regions. PCA uses the pooled variance, MclustDR (EDDA, VVV) the class means and covariances.

Purified profiles in two latent directions

high overlap / CT1 SBM / CT2 SBM



Equal numbers of draws per type; ellipses: 95% normal regions. PCA uses the pooled variance, MclustDR (EDDA, VVV) the class means and covariances.