
Wald approximation: D^2 of the ILR estimate against chi-square


balanced / low overlap



Grey band: 95% pointwise Monte Carlo envelope of sorted chi-square samples of the same size.
Log axes keep the bulk of the points visible next to the large values of estimates near a face.

Wald approximation: D^2 of the ILR estimate against chi-square

balanced / moderate overlap




Converged, local maximum Converged, not a local maximum Numerical failure

Grey band: 95% pointwise Monte Carlo envelope of sorted chi-square samples of the same size.
Log axes keep the bulk of the points visible next to the large values of estimates near a face.

Wald approximation: D^2 of the ILR estimate against chi-square


balanced / high overlap



Grey band: 95% pointwise Monte Carlo envelope of sorted chi-square samples of the same size.
Log axes keep the bulk of the points visible next to the large values of estimates near a face.

Wald approximation: D^2 of the ILR estimate against chi-square


moderately unbalanced / low overlap



Grey band: 95% pointwise Monte Carlo envelope of sorted chi-square samples of the same size.
Log axes keep the bulk of the points visible next to the large values of estimates near a face.

Wald approximation: D^2 of the ILR estimate against chi-square


moderately unbalanced / moderate overlap



Grey band: 95% pointwise Monte Carlo envelope of sorted chi-square samples of the same size.
Log axes keep the bulk of the points visible next to the large values of estimates near a face.

Wald approximation: D^2 of the ILR estimate against chi-square


moderately unbalanced / high overlap



Grey band: 95% pointwise Monte Carlo envelope of sorted chi-square samples of the same size.
Log axes keep the bulk of the points visible next to the large values of estimates near a face.

Wald approximation: D^2 of the ILR estimate against chi-square


highly unbalanced / low overlap



Grey band: 95% pointwise Monte Carlo envelope of sorted chi-square samples of the same size.
Log axes keep the bulk of the points visible next to the large values of estimates near a face.

Wald approximation: D^2 of the ILR estimate against chi-square


highly unbalanced / moderate overlap



Grey band: 95% pointwise Monte Carlo envelope of sorted chi-square samples of the same size.
Log axes keep the bulk of the points visible next to the large values of estimates near a face.

Wald approximation: D^2 of the ILR estimate against chi-square

highly unbalanced / high overlap



Grey band: 95% pointwise Monte Carlo envelope of sorted chi-square samples of the same size.
Log axes keep the bulk of the points visible next to the large values of estimates near a face.