
Purified profiles in two latent directions


G = 20 / low sibling overlap / equal cell sizes



Equal numbers of draws per type; ellipses: 95% normal regions. PCA uses the pooled variance, MclustDR (EDDA, VVV) the class means and covariances.

Purified profiles in two latent directions


G = 20 / low sibling overlap / lineage cell sizes



Equal numbers of draws per type; ellipses: 95% normal regions. PCA uses the pooled variance, MclustDR (EDDA, VVV) the class means and covariances.

Purified profiles in two latent directions


G = 20 / high sibling overlap / equal cell sizes



Equal numbers of draws per type; ellipses: 95% normal regions. PCA uses the pooled variance, MclustDR (EDDA, VVV) the class means and covariances.

Purified profiles in two latent directions


G = 20 / high sibling overlap / lineage cell sizes



Equal numbers of draws per type; ellipses: 95% normal regions. PCA uses the pooled variance, MclustDR (EDDA, VVV) the class means and covariances.

Purified profiles in two latent directions


G = 200 / low sibling overlap / equal cell sizes



Equal numbers of draws per type; ellipses: 95% normal regions. PCA uses the pooled variance, MclustDR (EDDA, VVV) the class means and covariances.

Purified profiles in two latent directions


G = 200 / low sibling overlap / lineage cell sizes



Equal numbers of draws per type; ellipses: 95% normal regions. PCA uses the pooled variance, MclustDR (EDDA, VVV) the class means and covariances.

Purified profiles in two latent directions


G = 200 / high sibling overlap / equal cell sizes



Equal numbers of draws per type; ellipses: 95% normal regions. PCA uses the pooled variance, MclustDR (EDDA, VVV) the class means and covariances.

Purified profiles in two latent directions

G = 200 / high sibling overlap / lineage cell sizes



Equal numbers of draws per type; ellipses: 95% normal regions. PCA uses the pooled variance, MclustDR (EDDA, VVV) the class means and covariances.