
Run time of every method


G = 20 / 1000 cells per bulk / balanced / low sibling overlap



Point estimate only; interval computations are not included. Competitors: batch time divided by the bulks.

Run time of every method


G = 20 / 1000 cells per bulk / balanced / high sibling overlap



Point estimate only; interval computations are not included. Competitors: batch time divided by the bulks.

Run time of every method


G = 20 / 1000 cells per bulk / unbalanced / low sibling overlap



Point estimate only; interval computations are not included. Competitors: batch time divided by the bulks.

Run time of every method


G = 20 / 1000 cells per bulk / unbalanced / high sibling overlap



Point estimate only; interval computations are not included. Competitors: batch time divided by the bulks.

Run time of every method


G = 20 / 1e+05 cells per bulk / balanced / low sibling overlap



Point estimate only; interval computations are not included. Competitors: batch time divided by the bulks.

Run time of every method


G = 20 / 1e+05 cells per bulk / balanced / high sibling overlap



Point estimate only; interval computations are not included. Competitors: batch time divided by the bulks.

Run time of every method


G = 20 / 1e+05 cells per bulk / unbalanced / low sibling overlap



Point estimate only; interval computations are not included. Competitors: batch time divided by the bulks.

Run time of every method


G = 20 / 1e+05 cells per bulk / unbalanced / high sibling overlap



Point estimate only; interval computations are not included. Competitors: batch time divided by the bulks.

Run time of every method


G = 200 / 1000 cells per bulk / balanced / low sibling overlap



Point estimate only; interval computations are not included. Competitors: batch time divided by the bulks.

Run time of every method


G = 200 / 1000 cells per bulk / balanced / high sibling overlap



Point estimate only; interval computations are not included. Competitors: batch time divided by the bulks.

Run time of every method


G = 200 / 1000 cells per bulk / unbalanced / low sibling overlap



Point estimate only; interval computations are not included. Competitors: batch time divided by the bulks.

Run time of every method


G = 200 / 1000 cells per bulk / unbalanced / high sibling overlap



Point estimate only; interval computations are not included. Competitors: batch time divided by the bulks.

Run time of every method


G = 200 / 1e+05 cells per bulk / balanced / low sibling overlap



Point estimate only; interval computations are not included. Competitors: batch time divided by the bulks.

Run time of every method


G = 200 / 1e+05 cells per bulk / balanced / high sibling overlap



Point estimate only; interval computations are not included. Competitors: batch time divided by the bulks.

Run time of every method


G = 200 / 1e+05 cells per bulk / unbalanced / low sibling overlap



Point estimate only; interval computations are not included. Competitors: batch time divided by the bulks.

Run time of every method

G = 200 / 1e+05 cells per bulk / unbalanced / high sibling overlap



Point estimate only; interval computations are not included. Competitors: batch time divided by the bulks.