
Expected log-likelihood

small centroid gap / homoscedastic / balanced




Expected log-likelihood at the true composition, maximised at p* exactly. Fill: log10(L / max L), floored at -10.

◇ true p* ● Monte Carlo mean of p-hat (Newton-Raphson)

Expected log-likelihood

small centroid gap / homoscedastic / moderately unbalanced




Expected log-likelihood at the true composition, maximised at p* exactly. Fill: log10(L / max L), floored at -10.

◇ true p* ● Monte Carlo mean of p-hat (Newton-Raphson)

Expected log-likelihood

small centroid gap / homoscedastic / highly unbalanced




Expected log-likelihood at the true composition, maximised at p* exactly. Fill: log10(L / max L), floored at -10.

◇ true p* ● Monte Carlo mean of p-hat (Newton-Raphson)

Expected log-likelihood

small centroid gap / heteroscedastic / balanced




Expected log-likelihood at the true composition, maximised at p* exactly. Fill: log10(L / max L), floored at -10.

◇ true p* ● Monte Carlo mean of p-hat (Newton-Raphson)

Expected log-likelihood

small centroid gap / heteroscedastic / moderately unbalanced




Expected log-likelihood at the true composition, maximised at p* exactly. Fill: log10(L / max L), floored at -10.

◇ true p* ● Monte Carlo mean of p-hat (Newton-Raphson)

Expected log-likelihood

small centroid gap / heteroscedastic / highly unbalanced




Expected log-likelihood at the true composition, maximised at p* exactly. Fill: log10(L / max L), floored at -10.

◇ true p* ● Monte Carlo mean of p-hat (Newton-Raphson)

Expected log-likelihood

large centroid gap / homoscedastic / balanced




Expected log-likelihood at the true composition, maximised at p* exactly. Fill: log10(L / max L), floored at -10.

◇ true p* ● Monte Carlo mean of p-hat (Newton-Raphson)

Expected log-likelihood

large centroid gap / homoscedastic / moderately unbalanced




Expected log-likelihood at the true composition, maximised at p* exactly. Fill: log10(L / max L), floored at -10.

◇ true p* ● Monte Carlo mean of p-hat (Newton-Raphson)

Expected log-likelihood

large centroid gap / homoscedastic / highly unbalanced




Expected log-likelihood at the true composition, maximised at p* exactly. Fill: log10(L / max L), floored at -10.

◇ true p* ● Monte Carlo mean of p-hat (Newton-Raphson)

Expected log-likelihood

large centroid gap / heteroscedastic / balanced




Expected log-likelihood at the true composition, maximised at p* exactly. Fill: log10(L / max L), floored at -10.

◇ true p* ● Monte Carlo mean of p-hat (Newton-Raphson)

Expected log-likelihood

large centroid gap / heteroscedastic / moderately unbalanced




Expected log-likelihood at the true composition, maximised at p* exactly. Fill: log10(L / max L), floored at -10.

◇ true p* ● Monte Carlo mean of p-hat (Newton-Raphson)

Expected log-likelihood

large centroid gap / heteroscedastic / highly unbalanced



Expected log-likelihood at the true composition, maximised at p* exactly. Fill: log10(L / max L), floored at -10.

◇ true p* ● Monte Carlo mean of p-hat (Newton-Raphson)