
Wald approximation: D^2 of the ILR estimate against chi-square


small centroid gap / homoscedastic / balanced



Grey band: 95% pointwise Monte Carlo envelope of sorted chi-square samples of the same size.
Log axes keep the bulk of the points visible next to the large values of estimates near a face.

Wald approximation: D^2 of the ILR estimate against chi-square


small centroid gap / homoscedastic / moderately unbalanced



Grey band: 95% pointwise Monte Carlo envelope of sorted chi-square samples of the same size.
Log axes keep the bulk of the points visible next to the large values of estimates near a face.

Wald approximation: D^2 of the ILR estimate against chi-square


small centroid gap / homoscedastic / highly unbalanced



Grey band: 95% pointwise Monte Carlo envelope of sorted chi-square samples of the same size.
Log axes keep the bulk of the points visible next to the large values of estimates near a face.

Wald approximation: D^2 of the ILR estimate against chi-square


small centroid gap / heteroscedastic / balanced



Grey band: 95% pointwise Monte Carlo envelope of sorted chi-square samples of the same size.
Log axes keep the bulk of the points visible next to the large values of estimates near a face.

Wald approximation: D^2 of the ILR estimate against chi-square


small centroid gap / heteroscedastic / moderately unbalanced



Grey band: 95% pointwise Monte Carlo envelope of sorted chi-square samples of the same size.
Log axes keep the bulk of the points visible next to the large values of estimates near a face.

Wald approximation: D^2 of the ILR estimate against chi-square


small centroid gap / heteroscedastic / highly unbalanced



Grey band: 95% pointwise Monte Carlo envelope of sorted chi-square samples of the same size.
Log axes keep the bulk of the points visible next to the large values of estimates near a face.

Wald approximation: D^2 of the ILR estimate against chi-square


large centroid gap / homoscedastic / balanced



Grey band: 95% pointwise Monte Carlo envelope of sorted chi-square samples of the same size.
Log axes keep the bulk of the points visible next to the large values of estimates near a face.

Wald approximation: D^2 of the ILR estimate against chi-square


large centroid gap / homoscedastic / moderately unbalanced



Grey band: 95% pointwise Monte Carlo envelope of sorted chi-square samples of the same size.
Log axes keep the bulk of the points visible next to the large values of estimates near a face.

Wald approximation: D^2 of the ILR estimate against chi-square


large centroid gap / homoscedastic / highly unbalanced



Grey band: 95% pointwise Monte Carlo envelope of sorted chi-square samples of the same size.
Log axes keep the bulk of the points visible next to the large values of estimates near a face.

Wald approximation: D^2 of the ILR estimate against chi-square


large centroid gap / heteroscedastic / balanced




Grey band: 95% pointwise Monte Carlo envelope of sorted chi-square samples of the same size.
Log axes keep the bulk of the points visible next to the large values of estimates near a face.

Wald approximation: D^2 of the ILR estimate against chi-square

large centroid gap / heteroscedastic / moderately unbalanced



Grey band: 95% pointwise Monte Carlo envelope of sorted chi-square samples of the same size.
Log axes keep the bulk of the points visible next to the large values of estimates near a face.



Grey band: 95% pointwise Monte Carlo envelope of sorted chi-square samples of the same size.
Log axes keep the bulk of the points visible next to the large values of estimates near a face.