



small_CLD / homoscedastic / balanced

$\rho = (0, 0)$




$\rho = (-0.8, -0.8)$




Y-axis is $L / \max(L) = \exp(\text{ell} - \max \text{ell})$ on a log10 scale (ggplot2::annotation_logticks).

$\rho = (0.8, 0.8)$



Y-axis is $L / \max(L) = \exp(\text{ell} - \max \text{ell})$ on a log10 scale (ggplot2::annotation_logticks).


$\rho = (-0.8, 0.8)$




Y-axis is $L / \max(L) = \exp(\text{ell} - \max \text{ell})$ on a log10 scale (ggplot2::annotation_logticks).

small_CLD / homoscedastic / moderately unbalanced

$\rho = (0, 0)$




$\rho = (-0.8, -0.8)$




Y-axis is $L / \max(L) = \exp(\text{ell} - \max \text{ell})$ on a log10 scale (ggplot2::annotation_logticks).

Y-axis is $L / \max(L) = \exp(\text{ell} - \max \text{ell})$ on a log10 scale (ggplot2::annotation_logticks).

$\rho = (0.8, 0.8)$



$\rho = (-0.8, 0.8)$




Y-axis is $L / \max(L) = \exp(\text{ell} - \max \text{ell})$ on a log10 scale (ggplot2::annotation_logticks).


Y-axis is $L / \max(L) = \exp(\text{ell} - \max \text{ell})$ on a log10 scale (ggplot2::annotation_logticks).

small_CLD / homoscedastic

$\rho = (0, 0)$




$\rho = (-0.8, -0.8)$




Y-axis is $L / \max(L) = \exp(\text{ell} - \max \text{ell})$ on a log10 scale (ggplot2::annotation_logticks).

Y-axis is $L / \max(L) = \exp(\text{ell} - \max \text{ell})$ on a log10 scale (ggplot2::annotation_logticks).

$\rho = (0.8, 0.8)$



$\rho = (-0.8, 0.8)$




Y-axis is $L / \max(L) = \exp(\text{ell} - \max \text{ell})$ on a log10 scale (ggplot2::annotation_logticks).


Y-axis is $L / \max(L) = \exp(\text{ell} - \max \text{ell})$ on a log10 scale (ggplot2::annotation_logticks).

small_CLD / heteroscedastic / balanced

$\rho = (0, 0)$




$\rho = (-0.8, -0.8)$




Y-axis is $L / \max(L) = \exp(\text{ell} - \max \text{ell})$ on a log10 scale (ggplot2::annotation_logticks).

Y-axis is $L / \max(L) = \exp(\text{ell} - \max \text{ell})$ on a log10 scale (ggplot2::annotation_logticks).

$\rho = (0.8, 0.8)$



$\rho = (-0.8, 0.8)$




Y-axis is $L / \max(L) = \exp(\text{ell} - \max \text{ell})$ on a log10 scale (ggplot2::annotation_logticks).


Y-axis is $L / \max(L) = \exp(\text{ell} - \max \text{ell})$ on a log10 scale (ggplot2::annotation_logticks).

small_CLD / heteroscedastic / moderately unbalanced

$\rho = (0, 0)$




$\rho = (-0.8, -0.8)$




Y-axis is $L / \max(L) = \exp(\text{ell} - \max \text{ell})$ on a log10 scale (ggplot2::annotation_logticks).

Y-axis is $L / \max(L) = \exp(\text{ell} - \max \text{ell})$ on a log10 scale (ggplot2::annotation_logticks).

$\rho = (0.8, 0.8)$



$\rho = (-0.8, 0.8)$




Y-axis is $L / \max(L) = \exp(\text{ell} - \max \text{ell})$ on a log10 scale (ggplot2::annotation_logticks).


Y-axis is $L / \max(L) = \exp(\text{ell} - \max \text{ell})$ on a log10 scale (ggplot2::annotation_logticks).

small_CLD / heteroscedastic

$\rho = (0, 0)$




$\rho = (-0.8, -0.8)$




Y-axis is $L / \max(L) = \exp(\text{ell} - \max \text{ell})$ on a log10 scale (ggplot2::annotation_logticks).

Y-axis is $L / \max(L) = \exp(\text{ell} - \max \text{ell})$ on a log10 scale (ggplot2::annotation_logticks).

$\rho = (0.8, 0.8)$



$\rho = (-0.8, 0.8)$




Y-axis is $L / \max(L) = \exp(\text{ell} - \max \text{ell})$ on a log10 scale (ggplot2::annotation_logticks).

Y-axis is $L / \max(L) = \exp(\text{ell} - \max \text{ell})$ on a log10 scale (ggplot2::annotation_logticks).


large_CLD / homoscedastic / balanced

$\rho = (0, 0)$




Y-axis is $L / \max(L) = \exp(\text{ell} - \max \text{ell})$ on a log10 scale (ggplot2::annotation_logticks).

$\rho = (-0.8, -0.8)$




Y-axis is $L / \max(L) = \exp(\text{ell} - \max \text{ell})$ on a log10 scale (ggplot2::annotation_logticks).

$\rho = (0.8, 0.8)$



Y-axis is $L / \max(L) = \exp(\text{ell} - \max \text{ell})$ on a log10 scale (ggplot2::annotation_logticks).


$\rho = (-0.8, 0.8)$




Y-axis is $L / \max(L) = \exp(\text{ell} - \max \text{ell})$ on a log10 scale (ggplot2::annotation_logticks).

large_CLD / homoscedastic / moderately unbalanced

$\rho = (0, 0)$




$\rho = (-0.8, -0.8)$




Y-axis is $L / \max(L) = \exp(\text{ell} - \max \text{ell})$ on a log10 scale (ggplot2::annotation_logticks).

Y-axis is $L / \max(L) = \exp(\text{ell} - \max \text{ell})$ on a log10 scale (ggplot2::annotation_logticks).

$\rho = (0.8, 0.8)$



$\rho = (-0.8, 0.8)$




Y-axis is $L / \max(L) = \exp(\text{ell} - \max \text{ell})$ on a log10 scale (ggplot2::annotation_logticks).

Y-axis is $L / \max(L) = \exp(\text{ell} - \max \text{ell})$ on a log10 scale (ggplot2::annotation_logticks).


large_CLD / homoscedastic / highly unbalanced

$\rho = (0, 0)$




Y-axis is $L / \max(L) = \exp(\text{ell} - \max \text{ell})$ on a log10 scale (ggplot2::annotation_logticks).

$\rho = (-0.8, -0.8)$




Y-axis is $L / \max(L) = \exp(\text{ell} - \max \text{ell})$ on a log10 scale (ggplot2::annotation_logticks).

$\rho = (0.8, 0.8)$




Y-axis is $L / \max(L) = \exp(\text{ell} - \max \text{ell})$ on a log10 scale (ggplot2::annotation_logticks).

$\rho = (-0.8, 0.8)$




Y-axis is $L / \max(L) = \exp(\text{ell} - \max \text{ell})$ on a log10 scale (ggplot2::annotation_logticks).


large_CLD / heteroscedastic / balanced




Y-axis is $L / \max(L) = \exp(\text{ell} - \max \text{ell})$ on a log10 scale (ggplot2::annotation_logticks).



Y-axis is $L / \max(L) = \exp(\text{ell} - \max \text{ell})$ on a log10 scale (ggplot2::annotation_logticks).




Y-axis is $L / \max(L) = \exp(\text{ell} - \max \text{ell})$ on a log10 scale (ggplot2::annotation_logticks).



Y-axis is $L / \max(L) = \exp(\text{ell} - \max \text{ell})$ on a log10 scale (ggplot2::annotation_logticks).


large_CLD / heteroscedastic / moderately unbalanced

$\rho = (0, 0)$




Y-axis is $L / \max(L) = \exp(\text{ell} - \max \text{ell})$ on a log10 scale (ggplot2::annotation_logticks).

$\rho = (-0.8, -0.8)$




Y-axis is $L / \max(L) = \exp(\text{ell} - \max \text{ell})$ on a log10 scale (ggplot2::annotation_logticks).

$\rho = (0.8, 0.8)$



Y-axis is $L / \max(L) = \exp(\text{ell} - \max \text{ell})$ on a log10 scale (ggplot2::annotation_logticks).


$\rho = (-0.8, 0.8)$



Y-axis is $L / \max(L) = \exp(\text{ell} - \max \text{ell})$ on a log10 scale (ggplot2::annotation_logticks).


large_CLD / heteroscedastic / highly unbalanced

$\rho = (0, 0)$




Y-axis is $L / \max(L) = \exp(\text{ell} - \max \text{ell})$ on a log10 scale (ggplot2::annotation_logticks).

$\rho = (-0.8, -0.8)$




Y-axis is $L / \max(L) = \exp(\text{ell} - \max \text{ell})$ on a log10 scale (ggplot2::annotation_logticks).

$\rho = (0.8, 0.8)$



Y-axis is $L / \max(L) = \exp(\text{ell} - \max \text{ell})$ on a log10 scale (ggplot2::annotation_logticks).

$\rho = (-0.8, 0.8)$



Y-axis is $L / \max(L) = \exp(\text{ell} - \max \text{ell})$ on a log10 scale (ggplot2::annotation_logticks).