



balanced / low OVL / SF/SF




A: Independent Thomson factanal() on purified type 1 (regression scores).
B: Independent Thomson factanal() on purified type 2 (regression scores).
C: Independent Thomson factanal() on purified type 3 (regression scores).
D: MCFA fitted to convolution bulk draws (weights p_j^2); purified slices scored in that plane.
E: MCFA fitted to unsupervised mixture draws (one type per draw, weights p_j).
F: Supervised MclustDA (EDDA) then MclustDR on labelled mixture draws. Insets: first 10 eigenvalues.

balanced / low OVL / SF/SBM




A: Independent Thomson factanal() on purified type 1 (regression scores).
B: Independent Thomson factanal() on purified type 2 (regression scores).
C: Independent Thomson factanal() on purified type 3 (regression scores).
D: MCFA fitted to convolution bulk draws (weights p_j^2); purified slices scored in that plane.
E: MCFA fitted to unsupervised mixture draws (one type per draw, weights p_j).
F: Supervised MclustDA (EDDA) then MclustDR on labelled mixture draws. Insets: first 10 eigenvalues.




A: Independent Thomson factanal() on purified type 1 (regression scores).
B: Independent Thomson factanal() on purified type 2 (regression scores).
C: Independent Thomson factanal() on purified type 3 (regression scores).
D: MCFA fitted to convolution bulk draws (weights p_j^2); purified slices scored in that plane.
E: MCFA fitted to unsupervised mixture draws (one type per draw, weights p_j).
F: Supervised MclustDA (EDDA) then MclustDR on labelled mixture draws. Insets: first 10 eigenvalues.

balanced / low OVL / SBM/SBM




A: Independent Thomson factanal() on purified type 1 (regression scores).
B: Independent Thomson factanal() on purified type 2 (regression scores).
C: Independent Thomson factanal() on purified type 3 (regression scores).
D: MCFA fitted to convolution bulk draws (weights p_j^2); purified slices scored in that plane.
E: MCFA fitted to unsupervised mixture draws (one type per draw, weights p_j).
F: Supervised MclustDA (EDDA) then MclustDR on labelled mixture draws. Insets: first 10 eigenvalues.

balanced / high OVL / SF/SF




A: Independent Thomson factanal() on purified type 1 (regression scores).
B: Independent Thomson factanal() on purified type 2 (regression scores).
C: Independent Thomson factanal() on purified type 3 (regression scores).
D: MCFA fitted to convolution bulk draws (weights p_j^2); purified slices scored in that plane.
E: MCFA fitted to unsupervised mixture draws (one type per draw, weights p_j).
F: Supervised MclustDA (EDDA) then MclustDR on labelled mixture draws. Insets: first 10 eigenvalues.

balanced / high OVL / SF/SBM




A: Independent Thomson factanal() on purified type 1 (regression scores).
B: Independent Thomson factanal() on purified type 2 (regression scores).
C: Independent Thomson factanal() on purified type 3 (regression scores).
D: MCFA fitted to convolution bulk draws (weights p_j^2); purified slices scored in that plane.
E: MCFA fitted to unsupervised mixture draws (one type per draw, weights p_j).
F: Supervised MclustDA (EDDA) then MclustDR on labelled mixture draws. Insets: first 10 eigenvalues.

balanced / high OVL / SBM/SF




A: Independent Thomson factanal() on purified type 1 (regression scores).
B: Independent Thomson factanal() on purified type 2 (regression scores).
C: Independent Thomson factanal() on purified type 3 (regression scores).
D: MCFA fitted to convolution bulk draws (weights p_j^2); purified slices scored in that plane.
E: MCFA fitted to unsupervised mixture draws (one type per draw, weights p_j).
F: Supervised MclustDA (EDDA) then MclustDR on labelled mixture draws. Insets: first 10 eigenvalues.

balanced / high OVL / SBM/SBM




A: Independent Thomson factanal() on purified type 1 (regression scores).
B: Independent Thomson factanal() on purified type 2 (regression scores).
C: Independent Thomson factanal() on purified type 3 (regression scores).
D: MCFA fitted to convolution bulk draws (weights p_j^2); purified slices scored in that plane.
E: MCFA fitted to unsupervised mixture draws (one type per draw, weights p_j).
F: Supervised MclustDA (EDDA) then MclustDR on labelled mixture draws. Insets: first 10 eigenvalues.

highly unbalanced / low OVL / SF/SF




A: Independent Thomson factanal() on purified type 1 (regression scores).
B: Independent Thomson factanal() on purified type 2 (regression scores).
C: Independent Thomson factanal() on purified type 3 (regression scores).
D: MCFA fitted to convolution bulk draws (weights p_j^2); purified slices scored in that plane.
E: MCFA fitted to unsupervised mixture draws (one type per draw, weights p_j).
F: Supervised MclustDA (EDDA) then MclustDR on labelled mixture draws. Insets: first 10 eigenvalues.

highly unbalanced / low OVL / SF/SBM




A: Independent Thomson factanal() on purified type 1 (regression scores).
B: Independent Thomson factanal() on purified type 2 (regression scores).
C: Independent Thomson factanal() on purified type 3 (regression scores).
D: MCFA fitted to convolution bulk draws (weights p_j^2); purified slices scored in that plane.
E: MCFA fitted to unsupervised mixture draws (one type per draw, weights p_j).
F: Supervised MclustDA (EDDA) then MclustDR on labelled mixture draws. Insets: first 10 eigenvalues.

highly unbalanced / low OVL / SBM/SF




A: Independent Thomson factanal() on purified type 1 (regression scores).
B: Independent Thomson factanal() on purified type 2 (regression scores).
C: Independent Thomson factanal() on purified type 3 (regression scores).
D: MCFA fitted to convolution bulk draws (weights p_j^2); purified slices scored in that plane.
E: MCFA fitted to unsupervised mixture draws (one type per draw, weights p_j).
F: Supervised MclustDA (EDDA) then MclustDR on labelled mixture draws. Insets: first 10 eigenvalues.

highly unbalanced / low OVL / SBM/SBM




A: Independent Thomson factanal() on purified type 1 (regression scores).
B: Independent Thomson factanal() on purified type 2 (regression scores).
C: Independent Thomson factanal() on purified type 3 (regression scores).
D: MCFA fitted to convolution bulk draws (weights p_j^2); purified slices scored in that plane.
E: MCFA fitted to unsupervised mixture draws (one type per draw, weights p_j).
F: Supervised MclustDA (EDDA) then MclustDR on labelled mixture draws. Insets: first 10 eigenvalues.

highly unbalanced / high OVL / SF/SF




A: Independent Thomson factanal() on purified type 1 (regression scores).
B: Independent Thomson factanal() on purified type 2 (regression scores).
C: Independent Thomson factanal() on purified type 3 (regression scores).
D: MCFA fitted to convolution bulk draws (weights p_j^2); purified slices scored in that plane.
E: MCFA fitted to unsupervised mixture draws (one type per draw, weights p_j).
F: Supervised MclustDA (EDDA) then MclustDR on labelled mixture draws. Insets: first 10 eigenvalues.

highly unbalanced / high OVL / SF/SBM




A: Independent Thomson factanal() on purified type 1 (regression scores).
B: Independent Thomson factanal() on purified type 2 (regression scores).
C: Independent Thomson factanal() on purified type 3 (regression scores).
D: MCFA fitted to convolution bulk draws (weights p_j^2); purified slices scored in that plane.
E: MCFA fitted to unsupervised mixture draws (one type per draw, weights p_j).
F: Supervised MclustDA (EDDA) then MclustDR on labelled mixture draws. Insets: first 10 eigenvalues.

highly unbalanced / high OVL / SBM/SF



A: Independent Thomson factanal() on purified type 1 (regression scores).
B: Independent Thomson factanal() on purified type 2 (regression scores).
C: Independent Thomson factanal() on purified type 3 (regression scores).
D: MCFA fitted to convolution bulk draws (weights p_j^2); purified slices scored in that plane.
E: MCFA fitted to unsupervised mixture draws (one type per draw, weights p_j).
F: Supervised MclustDA (EDDA) then MclustDR on labelled mixture draws. Insets: first 10 eigenvalues.

highly unbalanced / high OVL / SBM/SBM



A: Independent Thomson factanal() on purified type 1 (regression scores).
B: Independent Thomson factanal() on purified type 2 (regression scores).
C: Independent Thomson factanal() on purified type 3 (regression scores).
D: MCFA fitted to convolution bulk draws (weights p_j^2); purified slices scored in that plane.
E: MCFA fitted to unsupervised mixture draws (one type per draw, weights p_j).
F: Supervised MclustDA (EDDA) then MclustDR on labelled mixture draws. Insets: first 10 eigenvalues.