
pysw Documentation

Release 0.1

Francis Poulin and Ben Storer

November 09, 2015

CONTENTS

Contents:

LAYERED QUASIGEOSTROPHIC MODEL

The N-layer quasigeostrophic (QG) potential vorticity is

$$\begin{aligned} q_1 &= \nabla^2 \psi_1 + \frac{f_0^2}{H_1} \left(\frac{\psi_2 - \psi_1}{g'_1} \right), & i = 1, \\ q_n &= \nabla^2 \psi_n + \frac{f_0^2}{H_n} \left(\frac{\psi_{n-1} - \psi_n}{g'_{n-1}} - \frac{\psi_i - \psi_{n+1}}{g'_i} \right), & i = 2, N-1, \\ q_N &= \nabla^2 \psi_N + \frac{f_0^2}{H_N} \left(\frac{\psi_{N-1} - \psi_N}{g'_{N-1}} \right) + \frac{f_0}{H_N} h_b(x, y), & i = N, \end{aligned}$$

where q_n is the n'th layer QG potential vorticity, and ψ_n is the streamfunction, f_0 is the inertial frequency, n'th H_n is the layer depth, and h_b is the bottom topography. (Note that in QG $h_b/H_N \ll 1$.) Also the n'th buoyancy jump (reduced gravity) is

$$g'_n \equiv g \frac{\rho_n - \rho_{n+1}}{\rho_n},$$

where g is the acceleration due to gravity and ρ_n is the layer density.

The dynamics of the system is given by the evolution of PV. We introduce a background flow that can vary in the horizontal. The streamfunction associated with this flow can be denoted with $\Psi_n(x, y)$ for each layer and geostrophy yields its corresponding velocity $\vec{V}_n = (U_n(x, y), V_n(x, y))$ where $\Psi_{ny} = -U_n$ and $\Psi_{nx} = V_n$. We can perturb the stream function in each layer into a background flow and deviations from that flow as,

$$\psi_n^{\text{tot}} = \Psi_n + \psi_n.$$

With this basic decomposition we can than write out the corresponding decompositions in velocity

$$\begin{aligned} u_n^{\text{tot}} &= U_n - \psi_{ny}, \\ v_n^{\text{tot}} &= V_n + \psi_{nx}, \end{aligned}$$

and

$$q_n^{\text{tot}} = Q_n + \delta_{nN} \frac{f_0}{H_N} h_b + q_n,$$

where $Q_n + \delta_{nN} \frac{f_0}{H_N} h_b$ is n'th layer background PV, we obtain the evolution equations

$$\begin{aligned} q_{nt} + \mathbf{J}(\psi_n, q_n + \delta_{nN} \frac{f_0}{H_N} h_b) + U_n(q_{nx} + \delta_{nN} \frac{f_0}{H_N} h_{bx}) + V_n(q_{ny} + \delta_{nN} \frac{f_0}{H_N} h_{by}) + \\ Q_{ny} \psi_{nx} - Q_{nx} \psi_{ny} = \text{ssd} - r_{ek} \delta_{nN} \nabla^2 \psi_n, \quad n = 1, N, \end{aligned}$$

where ssd is stands for small scale dissipation, which is achieved by an spectral exponential filter or hyperviscosity, and r_{ek} is the linear bottom drag coefficient. The Dirac delta, δ_{nN} , indicates that the drag is only applied in the bottom layer.

1.1 Linear Stability Analysis

In order to study the stability of a jet in the context of our n -layer QG model we focus our attention on basic states that consist of zonal flows. i.e. $\Psi_n(y)$ only. If we assume that the quadratic quantities we can then linearize to obtain in the conservative limit over a flat bottom,

$$q_{nt} + U_n q_{nx} + Q_{ny} \psi_{nx} = 0,$$

for $n = 1, \dots, N$.

We assume that the perturbations are normal modes in the zonal direction and time,

$$\psi_n = \text{Re}[\hat{\psi}_n e^{i(kx - \omega t)}].$$

This implies that the PV will be modified appropriately and we denote it with \hat{q}_n .

We substitute this into the linear equations and then divide by the exponential to obtain,

$$c\hat{q}_n = U_n \hat{q}_n + Q_{ny} \hat{\psi}_n,$$

where the basic state only depends on y , and layer of course, and we have introduced the phase speed $c = \omega/k$. Note that the actual PVs are

$$\begin{aligned} \hat{q}_1 &= (\partial_{yy} - k^2) \hat{\psi}_1 + \frac{f_0^2}{H_1} \left(\frac{\hat{\psi}_2 - \hat{\psi}_1}{g'_1} \right), & i = 1, \\ \hat{q}_n &= (\partial_{yy} - k^2) \hat{\psi}_n + \frac{f_0^2}{H_n} \left(\frac{\hat{\psi}_{n-1} - \hat{\psi}_n}{g'_{n-1}} - \frac{\hat{\psi}_i - \hat{\psi}_{n+1}}{g'_i} \right), & i = 2, N-1, \\ \hat{q}_N &= (\partial_{yy} - k^2) \hat{\psi}_N + \frac{f_0^2}{H_N} \left(\frac{\hat{\psi}_{N-1} - \hat{\psi}_N}{g'_{N-1}} \right), & i = N, \end{aligned}$$

SPECIAL CASE: ONE-LAYER MODEL

In the one-layer case we have

$$c\hat{q}_1 = U_1\hat{q}_1 + Q_{1y}\hat{\psi}_1,$$

$$\hat{q}_1 = \left[\partial_{yy} - k^2 - \frac{f_0^2}{g_1' H_1} \right] \hat{\psi}_1.$$

SPECIAL CASE: TWO-LAYER MODEL

In the two-layer case we have

$$c\hat{q}_n = U_n\hat{q}_n + Q_{ny}\hat{\psi}_n,$$
$$\hat{q}_1 = \left[\partial_{yy} - k^2 - \frac{f_0^2}{g_1' H_1} \right] \hat{\psi}_1 + \frac{f_0^2}{g_1' H_1} \hat{\psi}_2,$$
$$\hat{q}_2 = \frac{f_0^2}{g_1' H_2} \hat{\psi}_1 + \left[\partial_{yy} - k^2 - \frac{f_0^2}{g_1' H_2} \right] \hat{\psi}_2.$$

INDICES AND TABLES

- `genindex`
- `modindex`
- `search`