Urban Agriculture in Nigerian Cities

A spatial analysis of 484 OpenStreetMap features across 775 Local Government Areas

Jerry Bannister Zachary — Statistician, Abubakar Tafawa Balewa University (ATBU

15 October 2026

Urban agriculture is widely promoted as a lever for urban food security in Nigeria, yet its spatial distribution has rarely been measured. This study maps cultivated land in five Nigerian cities — Abuja, Lagos, Kano, Port Harcourt and Gombe — using 484 OpenStreetMap features, and tests whether their density across 775 Local Government Areas is spatially clustered, how it relates to LGA size, and whether apparent patterns survive correction for uneven mapping coverage. Density is strongly clustered (Moran's I = 0.300, z = 17.6, p < 1e-69). A spatial lag model reduces residual autocorrelation from 0.256 to −0.036, and a negative binomial model with area as an exposure offset confirms that density falls by roughly 66% per doubling of LGA area (coefficient -1.551 on log area, p = 8.5e-11). Because OpenStreetMap coverage is itself uneven — and is the main threat to the study's validity — per-LGA building and highway counts were collected as an independent exposure control. The confound is real (Spearman = 0.613 with mapped highways) but the area effect survives it and strengthens (ΔAIC = 20.0). A rainfall hypothesis could not be tested: climate was sampled at city centroids, leaving only five distinct values, so the analysis is reported at the city level where no effect is established. All results, including the corrections that produced them, are reproducible from frozen data.

Table of contents

1 The question **2**
1.1 H1 — Clustering . . . 3
1.2 H2 — Climate . . . 3
1.3 H3 — Scale . . . 3
1.4 H4 — Measurement . . . 3

2 Data **3**
2.1 What the 484 features are 4

3 Methods **5**
3.1 The unit of analysis 5
3.2 The central measurement problem 5
3.3 Spatial weights 6
3.4 The four tests 6

4 Findings **7**
4.1 H1 — Urban agriculture is strongly clustered 7
4.2 H1 at the point scale — clustering beyond the areal frame 9
4.3 H3 — Density falls sharply as LGA area grows 11
4.4 H4 — The mapping-coverage confound, measured 13
4.5 H2 — Withdrawn 15
4.6 Where the intensity actually is 17

5 What the audit changed **18**

6 Limitations **19**

7 Reproducing this **20**

What this document is

A complete, self-contained account of the spatial study: question, data, methods, findings, robustness, and the audit that corrected several of them. Every number is computed from the frozen data in `data/raw/` and can be re-derived with `Rscript analysis/spatial/audit_claims.R`.

This is original work. The OpenStreetMap, NASA POWER, World Bank and GADM data were collected by the author in August and October 2026 using public APIs. It is unrelated to the Bauchi Metropolis survey thesis, which is a separate body of work in the same repository.

1 The question

Urban agriculture in Nigeria is usually studied one city at a time, through household surveys that cannot say where cultivation happens or how it is distributed across a metropolitan area. The policy questions that matter — *where should land be protected, and how much of it is there?* — are spatial, but the evidence base mostly is not.

This study asks four questions of openly available spatial data:

1.1 H1 — Clustering

Is urban agriculture spatially clustered, or scattered at random?

If cultivation were randomly distributed across local government areas, there would be nothing distinctive to protect and no basis for targeting. If it is clustered, the clusters are where policy attention should go.

1.2 H2 — Climate

Does cultivation concentrate in drier areas?

Parts of northern Nigeria are far drier than the south, and irrigation potential differs sharply. If aridity drives where cultivation is found, dry-season water infrastructure becomes the binding constraint.

1.3 H3 — Scale

Does density fall as administrative area grows?

An LGA with ten mapped farms might be large and thinly covered or small and intensely farmed. Whether density depends on area determines whether any of this can be read as intensity at all.

1.4 H4 — Measurement

Does the pattern survive correction for mapping coverage?

OpenStreetMap is contributed unevenly. Kano is the best-mapped state in Nigeria *and* has the most recorded cultivation. If the whole pattern is an artefact of where volunteers have mapped, none of the above can be interpreted as agriculture.

2 Data

Five cities were chosen to span Nigeria's agro-ecological gradient, from the humid Niger Delta to the Sahelian north. All layers are frozen to `data/raw/` with a provenance record of the URL, timestamp, row count and licence for every pull.

Table 1: Data layers. Each was queried once and frozen; analysis never touches the network.

Source	Dates	Records	Purpose
OpenStreetMap Overpass API	25 Aug 2026	0	Cultivated land features: landuse=farmland, farmyard, allotments; leisure=garden
OpenStreetMap Overpass API (aggregate counts)	15 Oct 2026	78	Per-LGA building and highway counts, used as a mapping-coverage control
NASA POWER API	25 Aug 2026	—	Daily temperature, precipitation and humidity, 2023, at five city centroids
World Bank Indicators API	25 Aug 2026	75	National urban population, food production index, undernourishment
GADM 4.1	26 Aug 2026	—	Administrative boundaries, 775 Local Government Areas

2.1 What the 484 features are

Table 2: Feature types in the frozen OSM layer.

kind	Features	Share
farmland	328	67.8%
garden	105	21.7%
farmyard	46	9.5%
allotments	5	1.0%

Farming land (`farmland`, 328 features) dominates, followed by gardens (105) and farmyards (46). True allotments — the land-use class specifically reserved for cultivation — number only five. That asymmetry is itself a finding: cultivation in these cities is overwhelmingly recorded on land whose primary zoning is something else.

Table 3: Features by city. The spread is stark: Kano holds 189, Abuja 32.

city	Features	Share	Per mapped LGA
Kano	189	39.0%	15.8
Lagos	130	26.9%	8.7
Port Harcourt	88	18.2%	22.0
Gombe	45	9.3%	45.0
Abuja	32	6.6%	16.0

3 Methods

3.1 The unit of analysis

The areal unit is the Local Government Area — 775 of them, from GADM v4.1. LGAs are the right unit because zoning and land allocation are LGA responsibilities in Nigeria. They range in area from 11.6 to 10,358 km², a nine-hundredfold spread, which is why raw counts cannot be compared across them and density is used throughout.

Table 4: The analysis frame. Note the last row: every point reconciles to exactly one LGA.

Quantity	Value
Local Government Areas	775
LGAs containing at least one feature	39 (5.0%)
Share of LGAs with any feature	5.0%
Features in total	484
LGAs with density exactly zero	736 (95.0%)
Median LGA area	697 km²
Density range	0.000 – 1.058 per km²
Point-in-LGA reconciliation	484 of 484 assigned

3.2 The central measurement problem

Ninety-five per cent of LGAs have a density of exactly zero. That single fact drives most of the methodological choices that follow, and it is worth stating plainly rather than burying:




Figure 1: Distribution of urban agriculture density across 775 LGAs. The distribution is a spike at zero with a long right tail, skewness 15.5.

This shape rules out ordinary linear regression, which assumes a continuous, unbounded outcome. It does not rule out spatial statistics — Moran's I is a correlation coefficient and is fine here — but it does mean the regression step needed a different model family (see H3 below).

3.3 Spatial weights

Neighbour structure is defined by queen contiguity: two LGAs are neighbours if their boundaries share an edge *or* a vertex. Weights are row-standardised so that each LGA's neighbours sum to one, preventing well-connected LGAs from dominating.

$$w_{ij} = \begin{cases} 1 & \text{if LGAs } i, j \text{ share an edge or vertex} \\ 0 & \text{otherwise} \end{cases} \qquad w^*_{ij} = w_{ij} \Big/ \sum_j w_{ij}$$

Rook contiguity (shared edges only) is used as a robustness check.

3.4 The four tests

Hypothesis	Method	Why this method
H1 clustering	Global Moran's I; LISA with BH-FDR	Moran's I tests clustering of a continuous variable; LISA localises it. Benjamini–Hochberg controls the 775 simultaneous tests.
H1 (point scale)	Join-count test; Ripley's L within cities	The 484 features are a *point* pattern. Areal methods discretise them; these do not.
H2 climate	City-level regression ($K = 5$)	Rainfall was sampled at city centroids, so it cannot vary within a city. See the withdrawal below.
H3 scale	Negative binomial with log(area) as an **offset**	Counts with unequal exposure. Area enters as exposure, not as a regressor on a ratio.
H4 coverage	Per-LGA OSM building and highway counts as a second offset	Independently measures how much of each LGA is mapped at all.

4 Findings

4.1 H1 — Urban agriculture is strongly clustered

Table 5: Global Moran's I, queen contiguity, 775 LGAs.

Variable	Moran's I	Expected I	z	p
Density (per km^2)	0.2999	-0.00129	17.61	$<$ 1e-69
Raw counts	0.1584	-0.00129	8.47	$<$ 1e-16

Density is far more clustered than raw counts (I = 0.300 versus 0.158). That gap is not incidental — it shows that normalising by area reveals structure that counts conceal, because counts conflate intensity with administrative size.

It is not one state's doing. Excluding Kano entirely — the state with the most mapped features — leaves I = 0.274 (z = 16.3, p $<$ 1e-58). The clustering is national, not a single municipality.


Table 6: LISA results before and after multiplicity control.

Criterion	LGAs
Significant at raw $p < 0.05$	23
Benjamini–Hochberg FDR 5%	19
Bonferroni	18
Expected false positives at $= 0.05$	38.8

⚠ The LISA p-values are conservative, not the clustering absent

Twenty-three LGAs reach raw significance across 775 tests, while chance alone would produce about 38.8. Observing *fewer* significant units than chance is the signature of an uncalibrated test — the normal approximation behind LISA is discrete and conservative when 95% of the variable sits at zero. The defensible figure is the Benjamini–Hochberg set of **19**, and the count being below chance expectation is reported alongside it rather than hidden.

There is a second, more substantive caveat in the map below: 18 of those 19 LGAs also simply have any feature at all. The cluster map and the coverage map are, to a first approximation, the same map. That is H4, and it is why the exposure control later in this document matters.



4.2 H1 at the point scale — clustering beyond the areal frame

Areal methods impose an arbitrary spatial frame on what is really a point pattern. Two tests that do not:

Table 7: Point-pattern tests. Both indicate clustering well beyond random.

Test	Result
Join-count, 'has any feature' (z)	30.3 ($p < 2e\text{-}16$)
Ripley's L outside envelope	5 of 5
Cities tested	Abuja, Lagos, Kano, Port Harcourt, Gombe

Within every one of the five cities, the observed Ripley's L curve rises above its 199-simulation envelope — cultivation clusters at scales from a few hundred metres upward in each city independently. The join-count test agrees at the areal level.




Figure 2: The 484 features by city. Kano and Lagos dominate; the four other cities show distinct, separated clusters.

4.3 H3 — Density falls sharply as LGA area grows

Here the original analysis went wrong, and the correction is instructive enough to report in full rather than quietly fix.

Why the outcome is not what it first appears

The earlier version of this analysis fitted `density ~ log(area)` with an ordinary Gaussian model. Two things were wrong with that:

1. **Density is a ratio.** Regressing it on its own denominator partly recovers a mechanical relationship — the ratio fallacy. The magnitude of the "area effect" is not interpretable.
2. **Density is 95% zeros and strictly non-negative,** so a Gaussian is the wrong family. The fitted model returned **265 of 775 negative fitted densities**, which are impossible quantities.

The corrected specification treats the outcome as what it is — a *count* — and puts area in as an **exposure**, not a regressor:

$$n_i \sim \text{NegBin}(\mu_i, \theta), \qquad \log \mu_i = \log(\text{area}_i) + X_i\beta$$

Now $\exp(\beta)$ is an incidence rate ratio, which is exactly the density question asked correctly.

Table 8: Negative binomial, area as exposure offset. The IRR is per one-unit increase in log(area) - an e-fold (x2.718) increase. The effect per doubling is 2^b = 0.341.

Term	b	SE	p	IRR
Intercept	2.8348	1.5552	0.068	17.027
log(area)	-1.5510	0.2389	8.5e-11	0.212

Table 9: Model comparison at the correct family.

Model	AIC	Note
Poisson	3569.3	dispersion 39 — unusable
NegBin, offset only	601.6	no area effect
NegBin, offset + area slope	571.6	LR ^{2} = 32.0, p = 2e-08

H3 holds, and strongly. The area coefficient is −1.551, an incidence rate ratio of 0.212 per e-fold increase in area (p = 8.5e-11), and the likelihood-ratio test against an offset-only model is decisive (^{2} = 32.0, p = 1.5e-8).

Expressed in units a reader can use: **a doubling of LGA area multiplies density by 0.341**, a fall of about 66%. A tenfold increase multiplies it by 0.028. Small LGAs concentrate cultivation; large ones dilute it.

This is the same *direction* the flawed Gaussian model found, which is why the error was not obvious. What changed is that the magnitude is now interpretable and the predictions are physically possible.




Figure 3: Left: the superseded Gaussian model predicts negative densities for 34% of LGAs. Right: a rate model cannot.

4.4 H4 — The mapping-coverage confound, measured

The most serious threat to this study is that OpenStreetMap density measures *mapping effort* as much as it measures agriculture. Kano is both the best-mapped state in Nigerian OSM and the state with the most recorded cultivation. The two facts cannot be separated by wishing them apart, so per-LGA counts of buildings and highways were collected as an independent measure of how much of each LGA is mapped at all.

Table 10: The confound, quantified. LGAs with cultivation have nearly twice the mapped buildings of those without.

Comparison	Statistic
Features vs mapped buildings	= 0.410 (p = 0.05)
Features vs mapped highways	= 0.613 (p = 2.5e-09)
Median mapped buildings — LGAs with features	33,465
Median mapped buildings — LGAs without	17,347

The confound is real. LGAs containing cultivation have a median of 33,465 mapped buildings; those without have 17,347. Features correlate with mapped highway density at = 0.613.

And the finding survives it. Adding the mapped-building count as a second offset, so that the model predicts cultivation *per unit of mapped environment* rather than per km²:

Table 11: Does the area effect survive correction for mapping coverage? It does — and strengthens. ΔAIC = 20.0.

Model	log(area)	p	AIC	n
Area offset only	-1.0010	1e-09	374.2	78
Area + mapped-building offsets	-1.4213	2e-22	354.2	78

Table 12: The same coefficients in interpretable units. Exponents are ratios, so a doubling of area is 2^b, not exp(b).

Model	b on log(area)	Ratio per doubling (2^b)	Fall per doubling
Area offset only	-1.001	0.500	50%
Area + coverage control	-1.421	0.373	63%

The coefficient moves from −1.001 to −1.421 and the model improves by 20 AIC points. Rather than evaporating when coverage is controlled, the area effect gets **stronger**. That is the most important robustness result in this study: the confound is real, it is measured, and the substantive finding outlives it.




Figure 4: Bounding-box leakage: 43 of 484 features (8.9%) fall outside the five declared study states, because the Overpass query used rectangles rather than administrative boundaries.

4.5 H2 — Withdrawn

The climate hypothesis **cannot be tested with these data**, and the earlier claim that it was supported has been withdrawn.

Table 13: Why H2 cannot be tested.

Quantity	Value
Rows in the original 'N = 34' model	34
Distinct cities	5
Distinct rainfall values	5
R^2 of rainfall on city dummies	1.000
Rainfall effect at the honest level (K = 5)	p = 0.576
Spearman, K = 5	= -0.20 (p = 0.78)

Rainfall was sampled at five city centroids and then assigned to every LGA in that city. The predictor therefore had **five distinct values**, not thirty-four — regressing it on city dummies returns $R^2 = 1.000$, meaning rainfall *is* city identity. Treating 34 rows as independent observations inflated the apparent sample size roughly sevenfold, and the significant p-values were an artefact of that. At the level the data actually replicate — five cities — the effect disappears (p = 0.58).




Figure 5: The honest unit of analysis: five cities, not 34 LGAs. The data are consistent with an aridity effect but cannot establish one.

What would be needed: rainfall that varies within cities. CHIRPS (~5 km), ERA5-Land (~9 km) or TAMSAT (4 km) all rasterise to LGA polygons and would give each LGA its own value. Until then this remains an open question, not a finding.

4.6 Where the intensity actually is

Table 14: The ten most intensively cultivated LGAs.

LGA	State	Features	Area (km²)	Density (per km²)	LISA
Tarauni	Kano	25	23.6	1.058	HH
Shomolu	Lagos	10	18.3	0.546	HH
Ungogo	Kano	90	187.4	0.480	HH
Obio/Akpor	Rivers	64	275.5	0.232	not significant
Mushin	Lagos	4	17.4	0.230	HH
Gwale	Kano	4	23.8	0.168	HH
Nasarawa	Kano	3	19.0	0.158	HH
Kano	Kano	4	26.5	0.151	HH
Fagge	Kano	4	27.7	0.145	HH
Ikorodu	Lagos	53	369.5	0.143	not significant

The ranking exposes why counts mislead. Ikorodu records 53 features — more than any LGA except Ungogo — but spreads them across 369 km², giving a density of 0.143 and no significant cluster. Tarauni, with 25 features on 23.6 km², reaches 1.058 per km²: more than seven times Ikorodu's intensity on one-fifteenth the land.

5 What the audit changed

This study audited itself, and four findings changed the analysis. They are recorded here because a result that has survived scrutiny is worth more than one that has not.

Corrections register

1. Wrong model family. Density is 95% zeros and non-negative; a Gaussian spatial model produced 265 impossible negative fitted densities. → Negative binomial with area as an offset. *Effect on the finding: direction unchanged, magnitude now interpretable, predictions valid.*
2. No multiplicity control. LISA flagged 23 LGAs at raw $p < 0.05$ against 38.8 expected false positives. → Benjamini–Hochberg applied; 19 survive. *Effect: modest, and the raw count being below chance expectation is now reported as evidence of conservatism.*
3. Pseudoreplication. The "N = 34" climate model carried five distinct rainfall values. → H2 withdrawn and reported at K = 5. *Effect: a claimed finding removed.*
4. Uncontrolled coverage confound. OSM density partly measures mapping effort. → Per-LGA building and highway counts collected as an exposure control. *Effect: confound confirmed (= 0.613) and the area effect strengthened under control.*
Also corrected: 43 of 484 features (8.9%) fall outside the five declared study states because the Overpass query used rectangular bounding boxes — the "Abuja" box returns more features in Niger State (17) than in the Federal Capital Territory (15).

6 Limitations

OpenStreetMap is a volunteered dataset. Coverage is uneven, and completeness varies by city in ways that are correlated with — but not identical to — urban size. The exposure control addresses this directly for the area effect, but it cannot make the layer complete. The features are also a *proxy* for cultivation: `landuse=farmland` inside a city bounding box includes ordinary farmland.

`farmland` is not `allotments`. Only five of 484 features are recorded as allotments. What this study measures is therefore cultivated land within city regions, not formally designated urban agriculture. The distinction matters for policy, and it is under the data's control rather than the analyst's.

The study extent is rectangular. 43 features lie outside the five intended states. A corrected pull would filter by GADM polygons or an urban-extent mask rather than a bounding box, using the same `poly:` mechanism the exposure control already uses.

Five cities, one year. The features were frozen in August 2026 and climate is a single 2023 window. Neither supports inference about trends.

Cross-sectional, and not causal. The models describe where cultivation is recorded. Nothing here identifies why.

7 Reproducing this

```
# Re-derive every number in this document and check it against the claims
Rscript analysis/spatial/audit_claims.R

# Or run the whole pipeline from frozen data
Rscript build.R

# Render this report
quarto render docs/spatial-report.qmd --to html
```

Analysis reads only the frozen files in `data/raw/` and never the network, so results are stable. Refreshing the mapping-coverage layer is the single networked step: `python scripts/fetch_osm_exposure.py`.

Table 15: Analysis environment. Exact pinned versions are in renv.lock.

Component	Version
R	4.5.2
sf	1.1.3
spdep	1.4.2
spatialreg	1.4.3
MASS	7.3.66
spatstat	3.8.3

Data licensing

OpenStreetMap: ODbL, © OpenStreetMap contributors. GADM 4.1: CC BY 4.0. World Bank Indicators: CC BY 4.0. NASA POWER: public domain. Full provenance with URLs, timestamps and row counts is in `data/raw/provenance.json`.