

A Book of Abstract Algebra | (2nd Edition)

Chapter 32, Problem 4ED

Bookmark

Show all steps: ☒ ON

Problem

If $\alpha = \sqrt[4]{2}$ is a real fourth root of 2, then the four fourth roots of 2 are $\pm\alpha$ and $\pm i\alpha$. Explain parts 1–6, briefly but carefully:

$$[\mathbb{Q}(\alpha, i) : \mathbb{Q}] = 8.$$

Step-by-step solution

Step 1 of 2

The objective is to show that $[\mathbb{Q}(\sqrt[4]{2}, i) : \mathbb{Q}] = 8$.

[Comment](#)

Step 2 of 2

Clearly, $\sqrt[4]{2}$ is the root of $x^4 - 2$.

Also, $x^4 - 2$ is irreducible polynomial of lowest degree 4 over \mathbb{Q} by Eisenstein ($p = 2$).

Therefore, $[\mathbb{Q}(\sqrt[4]{2}) : \mathbb{Q}] = \deg(x^4 - 2) = 4$.

Because $\mathbb{Q}(\sqrt[4]{2})$ is a subfield of the reals and so, $i \notin \mathbb{Q}(\sqrt[4]{2})$.

Hence, $x^2 + 1$ is irreducible over $\mathbb{Q}(\sqrt[4]{2})$.

So, $[\mathbb{Q}(\sqrt[4]{2}, i) : \mathbb{Q}(\sqrt[4]{2})]$ is at least 2.

But i is a root of $x^2 + 1 \in \mathbb{Q}(\sqrt[4]{2})[X]$, so the degree of $\mathbb{Q}(\sqrt[4]{2}, i)$ over $\mathbb{Q}(\sqrt[4]{2})$ is at most 2 and therefore, is exactly 2.

Hence, $[\mathbb{Q}(\sqrt[4]{2}, i) : \mathbb{Q}(\sqrt[4]{2})] = 2$.

Thus , $\left[\mathbb{Q}(\sqrt[4]{2}, i) : \mathbb{Q} \right] = \left[\mathbb{Q}(\sqrt[4]{2}, i) : \mathbb{Q}(\sqrt[4]{2}) \right] \left[\mathbb{Q}(\sqrt[4]{2}) : \mathbb{Q} \right]$

$$= 2 \cdot 4$$

$$= 8.$$

[Comment](#)