

# A Book of Abstract Algebra | (2nd Edition)



Step-by-step solution

	<b>Step 1</b> of 3
Here, objective is to prove that, if $a \in D, a > 0$ , then $\sqrt{a} \in D$ .	
Commer	nt
	<b>Step 2</b> of 3
Construc	etible point:
-	nt is either the end point of given unit segment or it is the intersection of two lines ared by previous constructible points.
Commer	nt
	<b>Step 3</b> of 3
Conside	$a \in D$ if and only if the point $(a,0)$ is constructible from $\{O,I\}$
Now, we	have to prove that length $\sqrt{a}$ is constructed from the length $a$
Conside	r the below figure:



## figure:construction of sqare root of a

AB is the diameter of a circle.

By observing there exist two similar triangles

 $\triangle AOB$  and  $\triangle BOC$ 

Using the property of similar triangles, we have

$$\frac{1}{x} = \frac{x}{a}$$

$$x^2 = a$$

$$x = \sqrt{a}$$

Then, the length  $\sqrt{a}$  is constructed from  $\{O,I\}$  which implies  $\sqrt{a}\in D$ 

Therefore, if  $a \in D, a > 0$ , then  $\sqrt{a} \in D$ .

Hence, proved

Comment

#### **COMPANY**

About Chegg Chegg For Good College Marketing Corporate Development Investor Relations Jobs Join Our Affiliate Program Media Center Site Map

#### **LEGAL & POLICIES**

Advertising Choices
Cookie Notice
General Policies
Intellectual Property Rights
Terms of Use
Global Privacy Policy
Honor Code
Honor Shield

#### **CHEGG PRODUCTS AND SERVICES**

Cheap Textbooks Mobile Apps Chegg Coupon Sell Textbooks Chegg Play Solutions Manual Chegg Study Help Study 101 College Textbooks Textbook Rental eTextbooks **Used Textbooks** Digital Access Codes Flashcards Learn Chegg Money Chegg Math Solver

### **CHEGG NETWORK**

EasyBib Internships.com Thinkful

#### **CUSTOMER SERVICE**

Customer Service
Give Us Feedback
Help with eTextbooks
Help to use EasyBib Plus
Manage Chegg Study
Subscription
Return Your Books
Textbook Return Policy