

## A Book of Abstract Algebra | (2nd Edition)

Chapter 27, Problem 4EF

Bookmark

Show all steps:

ON

Problem

<

Let  $F$  be a finite field, and  $F^*$  the multiplicative group of nonzero elements of  $F$ . Obviously  $H = \{x^2: x \in F^*\}$  is a subgroup of  $F^*$ ; since every square  $x^2$  in  $F^*$  is the square of only two different elements, namely  $\pm x$ , exactly half the elements of  $F^*$  are in  $H$ . Thus,  $H$  has exactly two cosets:  $H$  itself, containing all the squares, and  $aH$  (where  $a \notin H$ ), containing all the nonsquares. If  $a$  and  $b$  are nonsquares, then by Chapter 15, Theorem 5(i),

$$ab^{-1} = \frac{a}{b} \in H$$

Thus: if  $a$  and  $b$  are nonsquares,  $a/b$  is a square. Use these remarks in the following:

If the minimum polynomial of  $a$  over  $F$  has degree 2, we call  $F(a)$  a quadratic extension of  $F$ .

Use part 3 to prove: Any two quadratic extensions of a finite field are isomorphic.

>

Step-by-step solution

Step 1 of 3 ^

Objective is to prove that any two quadratic extensions of a finite field are isomorphic.

Consider the finite field  $F$  and let  $a, b \in F$ . Assume that  $p(x)$ ,  $q(x)$  are arbitrary quadratic irreducible polynomials in  $F[x]$  and  $\sqrt{a}, \sqrt{b}$  are the roots of these polynomials in some extension of  $F$ .

Then  $a/b$  is a square and  $\sqrt{b}$  will be a root of  $p(cx)$  for some  $c$  in  $F$ .

Since  $p(x) \in F[x]$  is an irreducible polynomial and  $\sqrt{b}$  is a root of  $p(cx)$ . Therefore, it follows, from the above result, that

$$\frac{F[x]}{\langle p(cx) \rangle} \cong F(\sqrt{b}).$$

Comment

Step 2 of 3 ^

Next, objective is to prove that  $F(\sqrt{a}) \cong F(\sqrt{b})$ . It is known that  $\sqrt{a}$  is a root of  $p(x)$  and  $\sqrt{b}$  is a root of  $p(cx)$ . Thus,

$$\frac{F[x]}{\langle p(x) \rangle} \cong F(\sqrt{a}), \quad \frac{F[x]}{\langle p(cx) \rangle} \cong F(\sqrt{b}).$$

Since  $F[x]/\langle p(cx) \rangle \cong F[x]/\langle p(x) \rangle$ , therefore

$$F(\sqrt{a}) \cong F(\sqrt{b}).$$

Comment

Step 3 of 3 ^

Since polynomials  $p(x)$  and  $q(x)$  are arbitrary irreducibles in  $F[x]$  of degree 2, therefore it can conclude that any two quadratic extensions of a finite field are isomorphic.

Comment

