



May question

Time left 0:39:10

In the limiting reactant experiment, a precipitate was formed upon the addition of few drops of 0.5 M BaCl_2 solution to the supernatant, this indicates that:

- ☐ a. $\text{Ba}_3(\text{PO}_4)_2$ is the limiting reactant
- ☐ b. Na_3PO_4 is the limiting reactant
- ☐ c. BaCl_2 is the excess reactant
- ☐ d. BaCl_2 is the limiting reactant

Next page



/ Quiz # 1 repeated: Experiments # 1 + # 2 + # 3 + #4
Friday 8/4/2022 at 22:30 - 23:00

Time left 0:36:33

Question **1**

Not yet answered

Marked out of 1.00

Flag question

Which of the following glassware is used to accurately measure volumes?

- ☐ a. **Reagent Bottle**
- ☒ b. **Pipet**
- ☐ c. **Beaker**
- ☐ d. **Erlenmeyer flask**

[Clear my choice](#)

[Next page](#)

← Quiz # 1: 4 experiments:
(Including Empirical Formula of
MgO + L.R. in a salt mixture)
Saturday 1/4/2022 Online at
18:00 PM (6:00 Evening)

[Jump to...](#)





Time left 0:36:30

1.884 g of salt mixture containing the salts Na_3PO_4 (M.M. = 164. 2 g/mol) and CaCl_2 (M.M. 110.0 g/mol) is added to 200 mL water and 0.763g of $\text{Ca}_3(\text{PO}_4)_2$ (M.M. = 310.0 g/mol) precipitates. The reaction equation was:



Tests reveal that Na_3PO_4 is the excess reactant; then the mass percent of excess reactant is

☐ a. 62%

☐ b. 38%

☒ c. 43%

☐ d. 57%



The following data were obtained for an experiment to determine the mass percent of limiting reactant in salt mixture

i) Mass of salt mixture = 8.25g)

ii) Mass of filter paper = 1.88g)

iii) Mass of filter paper and $\text{Ba}_3(\text{PO}_4)_2$ = 4.63 g

Na_3PO_4 was the limiting reactant, the percent by mass of the BaCl_2 is

Molar mass for $\text{BaCl}_2 \cdot 2\text{H}_2\text{O}$ = 244.2g/mol,)

Na_3PO_4 = 164 g/mol., $\text{Ba}_3(\text{PO}_4)_2$ = 602.2 (.g/mol

81.8 %

.a ☐

85.4 %

.b ☐

40.6 %

.c ☐

59.4 %

.d ☐



Time left 0:33:31

The following data were obtained for an experiment to determine the empirical formula of an oxide

(i) Mass of empty crucible and cover = 20.74g

(ii) Mass of metal, crucible and cover = 26.63 g

(iii) Mass of crucible, cover, and final product = 28.22g

What is the empirical formula for the metal oxide?

(Atomic Mass of metal = 118.7 g/mol.; At. Mass of O= 16 g/mol.)

☐ a. MO_2

☒ b. M_2O

☐ c. MO

☐ d. M_2O_3





Not yet answered

Time left 0:29:53

Marked out of 1.00

Flag question

If 43.88 g of metal rod is placed in a graduated cylinder filled initially to 20.1 mL with water, and then rises to a volume of 45.2 mL, the density of the rode is (in g/mL):

☐ a. 8.60☐ b. 6.84☐ c. 1.39☒ d. 1.75[Clear my choice](#)[Next page](#)



الوقت المتبقي 0:39:48

علم هذا السؤال

Which of the following compound have the same empirical formula?

a. $\text{C}_6\text{H}_{12}\text{O}_6$ and C_6H_{12} ☐

b. $\text{C}_3\text{H}_6\text{O}_3$ and $\text{C}_6\text{H}_{12}\text{O}_6$ ☐

c. $\text{C}_2\text{H}_4\text{O}_2$ and $\text{H}_2\text{C}_2\text{O}_4$ ☐

d. $\text{C}_6\text{H}_6\text{O}$ and C_6H_6 ☐



s1.ebalqa.courses

عع





Flag question

Time left 0:24:59

In limiting reactant experiment, $\text{Ba}_3(\text{PO}_4)_2$ was washed with hot water to -----

- ☐ a. Decrease the solubility of precipitate
- ☒ b. Improve the filtration process
- ☐ c. Remove insoluble $\text{Ba}_3(\text{PO}_4)_2$
- ☐ d. Remove remaining NaCl in the wet precipitate

[Next page](#)[← Quiz # 1: 4](#)

experiments: (Including
Empirical Formula of
 MgO + L.R. in a salt
mixture) Saturday
1/4/2022 Online at 18:00
PM (6:00 Evening)





Time left 0:19:22

The following data were obtained for an experiment to determine the mass percent of limiting reactant in salt mixture:

(i) Mass of salt mixture = 8.25g

(ii) Mass of filter paper = 1.88g

(iii) Mass of filter paper and $\text{Ba}_3(\text{PO}_4)_2$ = 4.63 g

Na_3PO_4 was the limiting reactant, the percent by mass of the BaCl_2 is

(Molar mass for $\text{BaCl}_2 \cdot 2\text{H}_2\text{O}$ = 244.2g/mol, Na_3PO_4 = 164 g/mol., $\text{Ba}_3(\text{PO}_4)_2$ = 602.2 g/mol.)

☐ a. 85.4 %

☐ b. 81.8 %

☒ c. 40.6 %

☐ d. 59.4 %



Time left 0:14:50

Question **8**

Not yet answered

Marked out of 1.00

Flag question

In empirical formula experiment, if loss of ash had been lost while heating the magnesium then

- ☐ a. The measured mass of oxygen would be higher.
- ☐ b. The measured mass of Mg would be lower.
- ☒ c. The measured mass of oxygen would be lower.
- ☐ d. The measured mass of Mg would be higher.

[Clear my choice](#)[Next page](#)

← Quiz # 1: 4 experiments:
(Including Empirical Formula of
MgO + L.R. in a salt mixture)
Saturday 1/4/2022 Online at
18:00 PM (6:00 Evening)

[Jump to...](#)



/ Quiz # 1 repeated: Experiments # 1 + # 2 + # 3 + #4
Friday 8/4/2022 at 22:30 - 23:00

Time left 0:07:49

Question **10**

Not yet answered

Marked out of 1.00

🚩 Flag question

In determining of the empirical formula of MgO , water was added to the crucible containing ash in order to:

- ☒ a. **Convert Mg_3N_2 to $\text{Mg}(\text{OH})_2$**
- ☐ b. **Convert Mg_3N_2 to MgO**
- ☐ c. **Convert $\text{Mg}(\text{OH})_2$ to Mg**
- ☐ d. **Convert $\text{Mg}(\text{OH})_2$ to MgO**

[Clear my choice](#)

Finish attempt ...

← Quiz # 1: 4 experiments:
(Including Empirical
Formula of MgO + L.R. in a
salt mixture) Saturday
1/4/2022 Online at 18:00
PM (6:00 Evening)