

CmpE 362 Homework 1 : Getting Started with MATLAB

Prof. Dr. Fatih ALAGOZ
TA : Yekta Said CAN

Deadline : 06 / 03 / 2019 - 23:59 (Sharp)

1 Description

1.1 Problem 1

For problems 1-8, write a script called `signalAndNoise.m` and put all the commands in it. Separate and label different problems using comments.

Let x is vector of real numbers (-100:100)

plot $y_1 = \sin x$, $y_2 = \sin 50x$, $y_3 = 50 \sin x$, $y_4 = \sin x + 50$, $y_5 = \sin(x + 50)$, $y_6 = 50 \sin 50x$, $y_7 = x \sin x$, $y_8 = \sin x / x$

Use 4x2 subplot to fit all subfigures belong to a single figure (Hint: write help for SUBPLOT in MATLAB). For more information about these functions, you can look Chapter 2: Sinusoids in your textbook.

1.2 Problem 2

Let x is vector of real numbers (-20:20)

Plot $y_1 = \sin x$, $y_2 = \sin 50x$, $y_3 = 50 \sin x$, $y_4 = \sin x + 50$, $y_5 = \sin(x + 50)$, $y_6 = 50 \sin 50x$, $y_7 = x \sin x$, $y_8 = \sin x / x$, $y_9 = y_1 + y_2 + y_3 + y_4 + y_5 + y_6 + y_7 + y_8$

Use 5x2 subplot to fit all subfigures belong to a single figure. For more information about these functions, you can look Chapter 2: Sinusoids in your textbook.

1.3 Problem 3

`randn` generates zero-mean, unit variance Gaussian distributed random number in (-,). Generate 41 random numbers following Gaussian distributed random numbers, call this as vector z .

Plot $y_{10} = z$, $y_{11} = z + x$, $y_{12} = z + \sin x$, $y_{13} = z \sin x$, $y_{14} = x \sin z$, $y_{15} = \sin(x + z)$, $y_{16} = z \sin 50x$, $y_{17} = \sin(x + 50z)$, $y_{18} = \sin x / z$, $y_{19} = y_{11} + y_{12} + y_{13} + y_{14} + y_{15} + y_{16} + y_{17} + y_{18}$

Use 5x2 subplot to fit all subfigures belong to a single figure. For more information about these functions, you can look Chapter 2: Sinusoids in your textbook.

1.4 Problem 4

rand generates uniformly distributed random number in [0,1] . Generate 41 random numbers following uniformly distributed random numbers.

Plot $y_{20} = z$, $y_{21} = z+x$, $y_{22} = z+\sin x$, $y_{23} = z \sin x$, $y_{24} = x \sin z$, $y_{25} = \sin(x+z)$, $y_{26} = z \sin 50x$, $y_{27} = \sin(x+50z)$, $y_{28} = \sin x/z$, $y_{29} = y_{21}+y_{22}+y_{23}+y_{24}+y_{25}+y_{26}+y_{27}+y_{28}$

Use 5x2 subplot to fit all subfigures belong to a single figure.

1.5 Problem 5

Starting with z (0,1) Gaussian(Normal) Random variable. (Use help menu for hist)

- Generate 10000 random variables with mean 0, variance 1; call it r1 vector
 - Generate 10000 random variables with mean 0, variance 4; call it r2 vector
 - Generate 10000 random variables with mean 0, variance 16; call it r3 vector
 - Generate 10000 random variables with mean 0, variance 256; call it r4 vector
- Plot hist(r1), hist(r2), hist(r3),hist(r4) on the same figure for comparison purposes.

1.6 Problem 6

Starting with z (0,1) Gaussian Random variable. (Use help menu for hist)

- Generate 10000 random variables with mean 10, variance 1; call it r6 vector
 - Generate 10000 random variables with mean 20, variance 4; call it r7 vector
 - Generate 10000 random variables with mean -10, variance 1; call it r8 vector
 - Generate 10000 random variables with mean -20, variance 4; call it r9 vector
- Plot hist(r6), hist(r7), hist(r8),hist(r9) on the same figure for comparison purposes.

1.7 Problem 7

Starting with z (0,1) uniformly distributed random variable.

- Generate 10000 random variables with mean 0, variance 1; call it r11 vector
 - Generate 10000 random variables with mean 0, variance 4; call it r21 vector
 - Generate 10000 random variables with mean 0, variance 16; call it r31 vector
 - Generate 10000 random variables with mean 0, variance 256; call it r41 vector
- Plot hist(r11), hist(r21), hist(r31),hist(r41) on the same figure for comparison purposes.

1.8 Problem 8

Starting with z (0,1) uniformly distributed random variable. (Use help menu for hist)

- Generate 10000 random variables with mean 10, variance 1; call it `r61` vector
 - Generate 10000 random variables with mean 20, variance 4; call it `r71` vector
 - Generate 10000 random variables with mean -10, variance 1; call it `r81` vector
 - Generate 10000 random variables with mean -20, variance 4; call it `r91` vector
- Plot `hist(r61)`, `hist(r71)`, `hist(r81)`, `hist(r91)` on the same figure for comparison purposes.

1.9 Problem 9

You are given an input csv file: `exampleSignal.csv`). Determine all the peaks for the provided .csv file. You can use built-in **findpeaks** method. Plot this signal on time domain and mark detected peaks by using MATLAB plot function properties. Examine this plot by looking at it and analyze the peaks that you have found or missed. Describe reasons why the algorithm missed some peaks by demonstrating examples in your report. Create a script called `problem10.m` for this question.

1.10 Problem 10

You are provided an image with the name `lena.png` which is commonly used in image processing studies. First read the image with **imread** function. After that, convert this rgb image into a grayscale image by using **rgb2gray** method. Then you will compute the mean, standard deviation, maximum (and location of maximum), minimum (and location of minimum) of the matrix you obtained from the image.

2 Report Preparation

Put the figures and explanations about questions into your report. Report must be a pdf file. Briefly describe what you have learnt from the above plots (plots from Questions 1-8) in your report. Furthermore, briefly describe what you have learnt about MATLAB. What were the challenges that you faced? What are the differences (advantages and disadvantages) between MATLAB and the other programming languages you have learned so far?

3 Submission and Grading

Prepare a report (pdf file) includes your code , explanations and comments of your code for each question. Compress the report and code files. Name it as "YourNumber_CmpE362_HW1.rar. Upload the file by using canvas before the deadline.

4 Notes

Deadline is strict. Do not send after deadline. When copying is detected, both parties will get zero.