



CORRESPONDENCE

ARCHIVE

Simplified Calculation of Body-Surface Area

October 22, 1987

N Engl J Med 1987; 317:1098

DOI: 10.1056/NEJM198710223171717

824 Citing Articles

TO THE EDITOR

Values for body-surface area are commonly used in the practice of internal medicine, particularly to calculate doses of chemotherapeutic agents and index cardiac output and stroke volume. Body-surface area (BSA) is generally calculated from height (Ht) and weight (Wt), according to equations such as the classic 1916 Du Bois formula¹: $BSA (m^2) = 0.007184 \times Ht (cm)^{0.725} \times Wt (kg)^{0.425}$. Although many nomograms based on such equations have been published, some are inaccurate,² and one is not always readily available when a determination must be made. At our institution, we use a simple, easy-to-remember modification of an equation by Gehan and George,³ which requires the use of a calculator

$$BSA (m^2) = \sqrt{\frac{Ht (in) \times Wt (lb)}{3131}}$$

or, in metric:

$$BSA (m^2) = \sqrt{\frac{Ht (cm) \times Wt (kg)}{3600}}$$

with a square-root key:

If a 73-inch-tall, 175-lb patient is used as an example, the keystroke sequence on most calculators would be: $73 \times 175 = \div 3131 = \sqrt{} = 2.02 m^2$.

Although a slight degree of accuracy has been lost in making the above equations easy to remember, deviations from accepted values derived from other formulas^{1, 3, 4} are generally less than 2 percent. We have made good clinical application of these equations and believe they may be of benefit to other physicians.

R.D. Mosteller, M.D.
Akron City Hospital, Akron, OH 44309

4 References ▾

Citing Articles (824) ▾

Close Citing Articles

More from the week of October 22, 1987

← →

HE BETH ISRAEL HOSPITAL, BOSTON	ORIGINAL ARTICLE
Expression	Intermittent Coronary Occlusion in Acute Myocardial Infarction
	D. Hackett and Others