

## Research Project Overview

# 1 Concept overview

Given

$$\sigma(z, t) = H_A \int_0^t G(t-s) \frac{d\epsilon}{ds}(z, s) ds = H_A G * \frac{d\epsilon}{dt} \quad (1)$$

where

$$G(t) = 1 + c \int_{\tau_1}^{\tau_2} g(t, \tau) dF(\tau) = G_\infty + G_d \mathbb{E}_F[g] \quad (2)$$

In other words,  $G_\infty \equiv 1, G_d \equiv c$  with

$$g(t, \tau) = \frac{e^{-t/\tau}}{\tau}, \quad \tau = \text{"relaxation time"} \quad (3)$$

Note that

$$\hat{g} = \frac{1}{1 + i\omega\tau} \quad (4)$$

Then,  $\sigma(z, t)$  can be expressed as

$$\sigma(z, t) = H_A G_\infty \epsilon|_0^t + H_A P \quad (5)$$

where  $P := G_d \mathbb{E}_F[g] * d\epsilon/dt$ , giving

$$\hat{P} = G_d \mathbb{E}_F[\hat{g}] \cdot \frac{d\hat{\epsilon}}{dt} = \underbrace{\int_{\tau_1}^{\tau_2} \frac{G_d}{1 + i\omega\tau} dF(\tau)}_{\text{"}\hat{G}_r(\omega)\text{"}} \cdot \frac{d\hat{\epsilon}}{dt} \quad (6)$$

"Can show that"  $P = \mathbb{E}_F[\mathcal{P}]$  where  $\mathcal{P}$  satisfies an (auxiliary) ODE:

$$\tau \dot{\mathcal{P}} + \mathcal{P} = G_d \frac{d\epsilon}{dt}, \text{ with } \tau \sim F(\tau). \quad (7)$$

# 2 Project overview

c	
Weeks	Planned Activities
1, 2	Show this problem is analogous to Debye Polarization in Electromagnetics (auxiliary ODE same, although PDEs differ)
3, 4	Implement polynomial chaos to solve random ODE for $\mathcal{P}$
5...	Rigorously prove approach is valid recover Gauss-Legendre as special case