



Review of Lecture 13

- Validation




$E_{\text{val}}(g^-)$ estimates $E_{\text{out}}(g)$

- Data contamination



\mathcal{D}_{val} slightly contaminated

- Cross validation



10-fold cross validation

Learning From Data


Yaser S. Abu-Mostafa
California Institute of Technology




Lecture 14: Support Vector Machines



Sponsored by Caltech's Provost Office, E&AS Division, and IST • Thursday, May 17, 2012



Outline


- Maximizing the margin
- The solution 
- Nonlinear transforms

Better linear separation

Linearly separable data

Different separating lines

Which is best?



Two questions:




1. Why is bigger margin better?



2. Which \mathbf{w} maximizes the margin?

Remember the growth function?


All dichotomies with any line:



Dichotomies with fat margin



Fat margins imply fewer dichotomies



Finding \mathbf{w} with large margin



Let \mathbf{x}_n be the nearest data point to the plane $\mathbf{w}^\top \mathbf{x} = 0$. How far is it?

2 preliminary technicalities:



1. Normalize \mathbf{w} :

$$|\mathbf{w}^\top \mathbf{x}_n| = 1$$



2. Pull out w_0 :

$\mathbf{w} = (w_1, \dots, w_d)$ apart from b

The plane is now $\boxed{\mathbf{w}^\top \mathbf{x} + b = 0}$ (no x_0)

Computing the distance


The distance between \mathbf{x}_n and the plane $\mathbf{w}^\top \mathbf{x} + b = 0$ where $|\mathbf{w}^\top \mathbf{x}_n + b| = 1$

The vector \mathbf{w} is \perp to the plane in the \mathcal{X} space:

Take \mathbf{x}' and \mathbf{x}'' on the plane

$$\mathbf{w}^\top \mathbf{x}' + b = 0 \quad \text{and} \quad \mathbf{w}^\top \mathbf{x}'' + b = 0$$

$$\implies \mathbf{w}^\top (\mathbf{x}' - \mathbf{x}'') = 0$$



and the distance is ...


Distance between \mathbf{x}_n and the plane:

Take any point \mathbf{x} on the plane

Projection of $\mathbf{x}_n - \mathbf{x}$ on \mathbf{w}



$$\hat{\mathbf{w}} = \frac{\mathbf{w}}{\|\mathbf{w}\|} \implies \text{distance} = |\hat{\mathbf{w}}^T (\mathbf{x}_n - \mathbf{x})|$$



$$\text{distance} = \frac{1}{\|\mathbf{w}\|} |\mathbf{w}^T \mathbf{x}_n - \mathbf{w}^T \mathbf{x}| = \frac{1}{\|\mathbf{w}\|} |\mathbf{w}^T \mathbf{x}_n + b - \mathbf{w}^T \mathbf{x} - b| = \frac{1}{\|\mathbf{w}\|} \quad \text{}$$

The optimization problem

$$\text{Maximize } \frac{1}{\|\mathbf{w}\|}$$

$$\text{subject to } \min_{n=1,2,\dots,N} |\mathbf{w}^\top \mathbf{x}_n + b| = 1$$



$$\text{Notice: } |\mathbf{w}^\top \mathbf{x}_n + b| = y_n (\mathbf{w}^\top \mathbf{x}_n + b)$$

$$\text{Minimize } \frac{1}{2} \mathbf{w}^\top \mathbf{w}$$



$$\text{subject to } y_n (\mathbf{w}^\top \mathbf{x}_n + b) \geq 1 \quad \text{for } n = 1, 2, \dots, N$$

Outline

- Maximizing the margin
- The solution
- Nonlinear transforms

Constrained optimization

Minimize $\frac{1}{2} \mathbf{w}^\top \mathbf{w}$

subject to $y_n (\mathbf{w}^\top \mathbf{x}_n + b) \geq 1 \quad \text{for } n = 1, 2, \dots, N$



$$\mathbf{w} \in \mathbb{R}^d, \quad b \in \mathbb{R}$$


Lagrange? inequality constraints \implies KKT

We saw this before

Remember regularization?

$$\text{Minimize } E_{\text{in}}(\mathbf{w}) = \frac{1}{N} (\mathbf{Z}\mathbf{w} - \mathbf{y})^\top (\mathbf{Z}\mathbf{w} - \mathbf{y})$$

$$\text{subject to: } \mathbf{w}^\top \mathbf{w} \leq C$$

∇E_{in} normal to constraint 


optimize


constrain

Regularization: E_{in}

$$\mathbf{w}^\top \mathbf{w}$$

SVM: $\mathbf{w}^\top \mathbf{w}$

$$E_{\text{in}} \text{ $$



Lagrange formulation



Minimize $\mathcal{L}(\mathbf{w}, b, \boldsymbol{\alpha}) = \frac{1}{2} \mathbf{w}^\top \mathbf{w} - \sum_{n=1}^N \alpha_n (y_n (\mathbf{w}^\top \mathbf{x}_n + b) - 1)$



w.r.t. \mathbf{w} and b and maximize w.r.t. each $\alpha_n \geq 0$



$$\nabla_{\mathbf{w}} \mathcal{L} = \mathbf{w} - \sum_{n=1}^N \alpha_n y_n \mathbf{x}_n = \mathbf{0}$$

$$\frac{\partial \mathcal{L}}{\partial b} = - \sum_{n=1}^N \alpha_n y_n = 0$$

Substituting ...

$$\mathbf{w} = \sum_{n=1}^N \alpha_n y_n \mathbf{x}_n \quad \text{and} \quad \sum_{n=1}^N \alpha_n y_n = 0$$

in the Lagrangian

$$\mathcal{L}(\mathbf{w}, b, \boldsymbol{\alpha}) = \frac{1}{2} \mathbf{w}^\top \mathbf{w} - \sum_{n=1}^N \alpha_n (y_n (\mathbf{w}^\top \mathbf{x}_n + b) - 1)$$

we get

$$\mathcal{L}(\boldsymbol{\alpha}) = \sum_{n=1}^N \alpha_n - \frac{1}{2} \sum_{n=1}^N \sum_{m=1}^N y_n y_m \alpha_n \alpha_m \mathbf{x}_n^\top \mathbf{x}_m$$

Maximize w.r.t. to $\boldsymbol{\alpha}$ subject to $\alpha_n \geq 0$ for $n = 1, \dots, N$ and $\sum_{n=1}^N \alpha_n y_n = 0$

The solution - quadratic programming

$$\min_{\alpha} \quad \frac{1}{2} \alpha^T \underbrace{\begin{bmatrix} y_1 y_1 \mathbf{x}_1^T \mathbf{x}_1 & y_1 y_2 \mathbf{x}_1^T \mathbf{x}_2 & \dots & y_1 y_N \mathbf{x}_1^T \mathbf{x}_N \\ y_2 y_1 \mathbf{x}_2^T \mathbf{x}_1 & y_2 y_2 \mathbf{x}_2^T \mathbf{x}_2 & \dots & y_2 y_N \mathbf{x}_2^T \mathbf{x}_N \\ \dots & \dots & \dots & \dots \\ y_N y_1 \mathbf{x}_N^T \mathbf{x}_1 & y_N y_2 \mathbf{x}_N^T \mathbf{x}_2 & \dots & y_N y_N \mathbf{x}_N^T \mathbf{x}_N \end{bmatrix}}_{\text{quadratic coefficients}} \alpha + \underbrace{(-\mathbf{1}^T)}_{\text{linear}} \alpha$$

subject to $\underbrace{\mathbf{y}^T \alpha}_{\text{linear constraint}} = 0$

$$\underbrace{0}_{\text{lower bounds}} \leq \alpha \leq \underbrace{\infty}_{\text{upper bounds}}$$

QP hands us α



Solution: $\alpha = \alpha_1, \dots, \alpha_N$

$$\Rightarrow \mathbf{w} = \sum_{n=1}^N \alpha_n y_n \mathbf{x}_n$$

KKT condition: For $n = 1, \dots, N$


$$\alpha_n (y_n (\mathbf{w}^\top \mathbf{x}_n + b) - 1) = 0$$



We saw this before!

$\alpha_n > 0 \Rightarrow \mathbf{x}_n$ is a support vector


$E_{\text{in}} = \text{const.}$



Support vectors


Closest \mathbf{x}_n 's to the plane: achieve the margin 

$$\implies y_n (\mathbf{w}^\top \mathbf{x}_n + b) = 1$$

$$\mathbf{w} = \sum_{\mathbf{x}_n \text{ is SV}} \alpha_n y_n \mathbf{x}_n \quad \text{$$

Solve for b using any SV: 

$$y_n (\mathbf{w}^\top \mathbf{x}_n + b) = 1$$




Outline

- Maximizing the margin
- The solution
- Nonlinear transforms



z instead of **x**

$$\mathcal{L}(\alpha) = \sum_{n=1}^N \alpha_n - \frac{1}{2} \sum_{n=1}^N \sum_{m=1}^N y_n y_m \alpha_n \alpha_m \mathbf{z}_n^\top \mathbf{z}_m$$





“Support vectors” in \mathcal{X} space

Support vectors live in \mathcal{Z} space

In \mathcal{X} space, “pre-images” of support vectors

The margin is maintained in \mathcal{Z} space

Generalization result



$$\mathbb{E}[E_{\text{out}}] \leq \frac{\mathbb{E}[\# \text{ of SV's}]}{N - 1}$$

