

Prime specification

Shigeo Hattori

November 10, 2019

First revision: November, 2019

bayship.org@gmail.com

<https://github.com/bayship-org/mathematics>

1 Prerequisite definitions

[GitHub:Minor_of_memBer.pdf](#)

In the article above, in its first two pages, all prerequisite definitions for this article are given. Especially, the second page of the above article defines that:

- A **memBer** as a member of a set.
- A deep member y of a memBer x is calculated **the deep number relative to x** .
- Two memBers (x, y) are said (x is a **minor** of y).

2 Notations

Definition 2.1.

"And" is also written as " $_{and}\wedge$ ".

"Or" is also written as " $_{or}\vee$ ".

$(x \text{ }_{and}\wedge y)$ is not commutative because y possibly depends on x .

$(x \text{ }_{or}\vee y)$ is not commutative because y possibly depends on $\neg x$. ■

3 Introduction

All definition D of variables is said a **variable context**.

More precisely, D must be a sequence of sub definitions of which each defines

1
2

exactly one variable. And all terms of D must be pairwise in order of dependencies, i.e., the dependent appears later.	3
All variable context V is said independent if all term is only dependent of terms in V .	4
All independent variable context V is also said an antecedent context .	5
All variable context V is said a consequent context of an antecedent context A if ((A followed by C) is independent as a new variable context).	6
	7
In the rest of this section, take $\forall(A, C)$ as an antecedent context A and a consequent context C of A .	8
	9
	10
Take $\forall x$ as a variable of A . Then x is said specified if x represents exactly one entity.	11
For example, $x := 1$ then x is said specified; $\forall x : \in \mathbb{N}$ then x is not specified; let x be a point then x is not specified.	12
	13
Take $\forall x$ as a variable of C . Then x is said specified relatively to A if (x represents exactly one entity if you assume that all variables of A are specified).	14
For example, let A define $\forall n : \in \mathbb{N}$ and C define $x := n + 1$. Then x is specified relatively to A because if n had been specified in A then x represented exactly one natural number.	15
	16
	17
	18
Take $\forall(x, y)$ as (a variable x of C) and (a deep member y of x such that y 's deep number relative to x is countable). Then y is said specified relatively to A if x is specified relatively to A .	19
	20
	21
	22
	23
	24
	25
	26
Take $\forall x$ as a variable of C such that x is specified relatively to A . Then x is said primary specified relatively to A if $\neg(*1 \text{ and } \wedge *2 \text{ and } \wedge *3)$.	27
	28
	29
	30
	31
	32
1. x^1 is a set.	33
2. There exists $\exists N : \subset \mathbb{N}$ as a countable index set.	34
3. There exists $\exists \{S\}_{i \in N}$ as a collection of sets bijectively indexed by N such that $(*a1 \text{ and } \wedge \dots \text{ and } \wedge *a3)$.	35
	36
a1. $x = \bigcup_{i \in N} S_i$	37

¹As a value, x is a set.

a2. Take $\forall(i, j) : \in N$ such that $i \neq j$. Then $S_i \cap S_j \neq \emptyset$. 38

a3. Take $\forall i : \in N$. Then C specifies S_i relatively to A . 39

40

For example, let A define (X, T, M) as the Euclidean space of 2-dimension where 41
 needless to say no absolute coordinate system is defined. 42

Then (C probably can define x to be prime specified) 43

as $x :=$ "the set of all topological equivalences of a line segment in (X, T, M) ". 44

4 Conjecture 45

Conjecture 4.1 (Conjecture). 46

Take $\forall(n, X, T, M)$ as the Euclidean space (X, T, M) of n -dimension where X 47
 is the space, T is the topology and M is the metric table. 48

As a remark, for the set of all orthogonal coordinate systems for the space, no 49
 special member is defined. 50

Let $A := (n, X, T, M)$. 51

Take all (C, x) as a consequent context C of A as the antecedent context and a 52
 variable x of C . 53

If $*1$ holds then x is a minor of T . 54

1. x is primary specified relatively to A . ■ 55

5 Examples 56

This section just gives examples of substituting actual values into variables of 57
 the main conjecture. 58

59

Definition 5.1 (Unknot). 60

Refer to the main conjecture for (n, X, T, M) . 61

Take $\forall k1$ as an unknot such that $\text{Space}(k1) \subset X$. 62

For the main conjecture, this example substitutes values 63
 into (n, x) as $(*1 \text{ and } \wedge \dots \text{ and } \wedge *7)$. 64

1. Let $n := 3$. 65

2. Let $K := \{k \mid ((T, k), (T, k1)) \text{ are isomorphic} \}$. 66

3. Take $\forall k : \in K$. 67

4.	$f1(k) := \{d \mid d \text{ is a proper knot diagram of } k\}.$	68
5.	$f2(k) := \{r \mid$	69
	$\exists d : \in f1(k) \text{ and } \wedge$	70
	$r \text{ is the number of crossings on } d$	71
	$\}.$	72
6.	$f(k) := \text{"the maximum number of } f2(k).$	73
7.	$x := \{k \mid f(k) = f(k1)\}.$	74
		75