

**POISON**

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

# Nufarm **AMINE** 625

SELECTIVE HERBICIDE

**ACTIVE CONSTITUENT:**

**625 g/L 2,4-D present as the dimethylamine and diethanolamine salts**

**GROUP I HERBICIDE**

For the control of broadleaf weeds in fallow before direct drilling or sowing of cereals and pastures; and in cereal crops, pastures, sugar cane, peanuts and non-agricultural areas as per the Directions for Use.

THIS IS A PHENOXY HERBICIDE THAT CAN CAUSE SEVERE DAMAGE TO NATIVE VEGETATION AND SUSCEPTIBLE CROPS SUCH AS COTTON, GRAPES, TOMATOES, OILSEED CROPS AND ORNAMENTALS.



**Nufarm**

Grow a better tomorrow.

**RESTRAINTS****GENERAL RESTRAINTS**

**DO NOT** exceed maximum application rate of 5.3 L/ha.

**DO NOT** apply by hand-held spraying equipment at rates exceeding 5.3 L/ha or 530 mL/100 L.

**DO NOT** apply by spraying equipment carried on the back of the user (Manually pressurised backpack sprayer).

**DO NOT** apply by aircraft at rates exceeding 3.6 L/ha.

**DO NOT** use open mixing/loading equipment if treating more than 50 hectares in one day.

**DO NOT** apply if heavy rains or storms are forecast within 3 days or if any rain is likely within 6 hours.

**DO NOT** irrigate to the point of runoff for at least 3 days after application.

**DO NOT** apply if crop or weeds are stressed due to dry or excessively moist conditions.

Additional USAGE restrictions apply in certain states (listed below) and in peanut, pasture and sugarcane situations (refer to Section 5 of the Directions for Use Tables).

Tasmania:

- **DO NOT** apply to fairways adjacent to natural water bodies.
- Only apply in no-till farming systems for the following situations:
  - Preparatory sprays in dryland cropping
  - Pre-emergent application in winter cereals, summer cereals and legumes

South Australia:

- Only apply in no-till farming systems for the following situations:
  - Preparatory sprays in dryland cropping
  - Pre-emergent application in winter cereals, summer cereals and legumes

Western Australia:

- Only apply in no-till farming systems for pre-emergent applications in winter cereals and legumes.

**SPRAY DRIFT RESTRAINTS**

**DO NOT** apply by vertical sprayer.

Specific definitions for terms used in this section of the label can be found at [www.apvma.gov.au/spraydrift](http://www.apvma.gov.au/spraydrift)

**DO NOT** allow **bystanders** to come into contact with the spray cloud.

**DO NOT** apply in a manner that may cause an unacceptable impact to **native vegetation, agricultural crops, landscaped gardens and aquaculture production**, or cause contamination of plant or livestock commodities, outside the application site from **spray drift**. The **buffer zones** in the relevant buffer zone tables below provide guidance but may not be sufficient in all situations. Wherever possible, correctly use application equipment designed to reduce spray drift and apply when the wind direction is away from these sensitive areas.

**DO NOT** apply unless the **wind speed** is between 3 and 20 kilometres per hour at the **application site** during the time of application.

**DO NOT** apply if there are **hazardous surface temperature inversion** conditions present at the **application site** during the time of application. **Surface temperature inversion conditions** exist most evenings one to two hours before sunset and persist until one to two hours after sunrise.

**BOOM SPRAYERS**

**DO NOT** apply by a boom sprayer unless the following requirements are met:

- Spray droplets are not smaller than a VERY COARSE **spray droplet size category**
- Minimum distances between the **application site** and downwind **sensitive areas** (see 'Mandatory buffer zones' section of the following table titled 'Buffer zones for boom sprayers') are observed.

**Buffer Zones for Boom Sprayers**

| Application rate | Boom Height above target canopy | Mandatory buffer zones (distances given in metres) |                       |                  |                  |                 |
|------------------|---------------------------------|----------------------------------------------------|-----------------------|------------------|------------------|-----------------|
|                  |                                 | Bystander Areas                                    | Natural Aquatic Areas | Pollinator Areas | Vegetation Areas | Livestock Areas |
| Up to 500 mL/ha  | 0.5 m or lower                  | 0                                                  | Not required          | 0                | Not required     | 0               |
|                  | 1.0 m or lower                  |                                                    | 30                    |                  | 25               |                 |
| Up to 1.8 L/ha   | 0.5 m or lower                  |                                                    | 30                    |                  | 25               |                 |
|                  | 1.0 m or lower                  |                                                    | 60                    |                  | 60               |                 |
| Up to 2.5 L/ha   | 0.5 m or lower                  |                                                    | 30                    |                  | 30               |                 |
|                  | 1.0 m or lower                  |                                                    | 80                    |                  | 75               |                 |
| Up to 3 L/ha     | 0.5 m or lower                  |                                                    | 35                    |                  | 35               |                 |
|                  | 1.0 m or lower                  |                                                    | 90                    |                  | 90               |                 |
| Up to 3.6 L/ha   | 0.5 m or lower                  |                                                    | 40                    |                  | 35               |                 |
|                  | 1.0 m or lower                  |                                                    | 110                   |                  | 110              |                 |
| Up to 4.3 L/ha   | 0.5 m or lower                  |                                                    | 45                    |                  | 40               |                 |
|                  | 1.0 m or lower                  |                                                    | 130                   |                  | 130              |                 |
| Up to 4.4 L/ha   | 0.5 m or lower                  |                                                    | 45                    |                  | 45               |                 |
|                  | 1.0 m or lower                  |                                                    | 130                   |                  | 130              |                 |
| Up to 4.8 L/ha   | 0.5 m or lower                  |                                                    | 50                    |                  | 45               |                 |
|                  | 1.0 m or lower                  |                                                    | 150                   |                  | 140              |                 |
| Up to 5.3 L/ha   | 0.5 m or lower                  |                                                    | 55                    |                  | 50               |                 |
|                  | 1.0 m or lower                  |                                                    | 160                   |                  | 160              |                 |

**AIRCRAFT**

**DO NOT** apply by aircraft unless the following requirements are met:

- Spray droplets are no smaller than a VERY COARSE **spray droplet size category**
- For maximum release heights above the target canopy of 3 metres or 25 per cent of wingspan or 25 per cent of rotor diameter whichever is the greatest, minimum distances between the application site and downwind sensitive areas (see 'Mandatory buffer zones' section of the following table titled 'Buffer zones for aircraft') are observed.

**Buffer Zones For Aircraft**

| Application rate | Aircraft Type | Mandatory buffer zones (distances given in meters) |                       |                  |                  |                 |
|------------------|---------------|----------------------------------------------------|-----------------------|------------------|------------------|-----------------|
|                  |               | Bystander Areas                                    | Natural Aquatic Areas | Pollinator Areas | Vegetation Areas | Livestock Areas |
| Up to 500 mL/ha  | Fixed Wing    | 0                                                  | 80                    | 0                | 80               | 0               |
|                  | Helicopter    |                                                    | 60                    |                  | 60               |                 |
| Up to 900 mL/ha  | Fixed Wing    |                                                    | 120                   |                  | 120              |                 |
|                  | Helicopter    |                                                    | 85                    |                  | 80               |                 |
| Up to 1.5 L/ha   | Fixed Wing    |                                                    | 170                   |                  | 160              |                 |
|                  | Helicopter    |                                                    | 120                   |                  | 120              |                 |
| Up to 1.7 L/ha   | Fixed Wing    |                                                    | 180                   |                  | 180              |                 |
|                  | Helicopter    |                                                    | 130                   |                  | 120              |                 |
| Up to 1.8 L/ha   | Fixed Wing    |                                                    | 190                   |                  | 180              |                 |
|                  | Helicopter    |                                                    | 130                   |                  | 130              |                 |
| Up to 2.4 L/ha   | Fixed Wing    |                                                    | 230                   |                  | 220              |                 |
|                  | Helicopter    |                                                    | 160                   |                  | 150              |                 |
| Up to 3.6 L/ha   | Fixed Wing    |                                                    | 325                   |                  | 300              |                 |
|                  | Helicopter    |                                                    | 200                   |                  | 200              |                 |
| Up to 4.8 L/ha   | Fixed Wing    |                                                    | 425                   |                  | 400              |                 |
|                  | Helicopter    |                                                    | 250                   |                  | 240              |                 |
| Up to 5.3 L/ha   | Fixed Wing    |                                                    | 475                   |                  | 450              |                 |
|                  | Helicopter    |                                                    | 275                   |                  | 275              |                 |

**SECTION 1. CEREALS (POST EMERGENCE AND HARVEST AID/SALVAGE SPRAY APPLICATION)**

| SITUATION/CROP               | WEEDS                                                                                            | STATE             | RATE                                           | CRITICAL COMMENTS                                                                                                                                                                                                                                                                                           |
|------------------------------|--------------------------------------------------------------------------------------------------|-------------------|------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Wheat                        | Refer Weed Table                                                                                 | NSW, ACT, SA only | 0.32 - 1.7 L/ha                                | Apply after the first node can be felt at the base of a tiller and before swelling of the head can be felt in a tiller (NSW, ACT, SA only).                                                                                                                                                                 |
|                              |                                                                                                  | Vic only          | 0.225 - 1.4 L/ha                               | Apply from tillering to boot stage (Vic only).                                                                                                                                                                                                                                                              |
|                              |                                                                                                  | Qld only          | 0.56 - 1.4 L/ha                                | Apply from mid tillering to before boot stage (Qld only).                                                                                                                                                                                                                                                   |
|                              |                                                                                                  | Tas only          | 0.8 - 1.4 L/ha                                 | Apply at 5 leaf to fully tillered (Tas only).                                                                                                                                                                                                                                                               |
| Barley                       |                                                                                                  | NSW, ACT, SA only | 0.32 - 1.4 L/ha                                |                                                                                                                                                                                                                                                                                                             |
|                              |                                                                                                  | Vic only          | 0.225 - 1.4 L/ha                               |                                                                                                                                                                                                                                                                                                             |
|                              |                                                                                                  | Qld only          | 0.56 - 1.4 L/ha                                |                                                                                                                                                                                                                                                                                                             |
|                              |                                                                                                  | Tas only          | 0.8 - 1.4 L/ha                                 |                                                                                                                                                                                                                                                                                                             |
| Cereal rye, Triticale        |                                                                                                  | NSW, ACT, SA only | 0.32 - 1.4 L/ha                                |                                                                                                                                                                                                                                                                                                             |
|                              |                                                                                                  | Vic only          | 0.225 - 1.4 L/ha                               |                                                                                                                                                                                                                                                                                                             |
| Oats                         |                                                                                                  | NSW, SA only      | 320 - 800 mL/ha                                |                                                                                                                                                                                                                                                                                                             |
|                              |                                                                                                  | Vic only          | 225 - 800 mL/ha                                |                                                                                                                                                                                                                                                                                                             |
| Cereals: Wheat, Oats, Barley | Cape tulip                                                                                       | WA only           | 0.64 - 1.3 L/ha                                | Apply from the 5 leaf stage up to jointing stage (Z15-33). Apply after the 6 leaf stage (Z16) for cranbrook, jacup, aroona and spear wheat and mortlock oats to avoid possible damage. <b>DO NOT</b> spray if lucerne is present.                                                                           |
|                              | Dock, Flatweed, Saffron thistle                                                                  |                   | 1.1 L/ha                                       |                                                                                                                                                                                                                                                                                                             |
|                              | Indian hedge mustard, London rocket, Lupin, Matricaria, Rapistrum, Wild radish                   |                   | 800 mL/ha                                      | WEED STAGE: 10 - 15 cm. Docks should be sprayed before 5 leaf stage. Cape tulip – low rate for cornils only.                                                                                                                                                                                                |
|                              | Wild turnip                                                                                      |                   | 640 mL/ha                                      |                                                                                                                                                                                                                                                                                                             |
|                              | Capeweed, Doublegee, Erodium, London rocket, Lupin, Mustard, Rapistrum, Wild radish, Wild turnip |                   | 200 mL/ha + 278 g/ha of a diuron 900WG product | Apply when crop has 4 - 5 leaves and most weeds have germinated and are in 2 - 5 leaf stage. Crop and weeds should be dry at time of application. Some temporary yellowing of crop may occur after application. Undersown sub-clovers may be slightly retarded.<br><b>DO NOT</b> apply to undersown medics. |

| SITUATION/CROP                                                                                                         | WEEDS                                                                                                         | STATE                  | RATE            | CRITICAL COMMENTS                                                                                                                                                                                                                                                                                                                                   |
|------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------|------------------------|-----------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Cereals: Wheat, Barley, Oats, Triticale, Cereal rye                                                                    | Volunteer canola ( <i>Brassica napus</i> ) including Roundup Ready* varieties                                 | All States             | 1 L/ha          | WEED STAGE: Up to 4 leaf.<br>CROP STAGE: 5 leaf to fully tillered.                                                                                                                                                                                                                                                                                  |
|                                                                                                                        |                                                                                                               |                        | 1.4 L/ha        | WEED STAGE: Up to 6 leaf.<br>CROP STAGE: 5 leaf to fully tillered.                                                                                                                                                                                                                                                                                  |
| Fallow, Stubble Spray prior to direct drilling or sowing -Winter Cereals, Grain legumes (peanuts -Qld only) and Canola | Refer Weed Table                                                                                              | Refer Weed Table       | 0.225 -1.7 L/ha | Observe plant back periods given in the table on this leaflet. Can be mixed with Glean*, Shirquat® 250, Revolver® or SpraySeed* where grasses are present. Select appropriate rate from the Weed Table. For Skeleton weed spraying should only be done 6 to 8 weeks before anticipated sowing date and subsequent cultivation limited to a minimum. |
| Millet                                                                                                                 | Refer Weed Table                                                                                              | NSW, SA, ACT, Vic only | 0.56 - 1.1 L/ha | Spray when secondary roots have developed, when fully tillered and before heads start to form at the base of the tillers. <b>DO NOT</b> use on panorama millet or panicum.                                                                                                                                                                          |
|                                                                                                                        |                                                                                                               | Qld only               | 560 - 900 mL/ha |                                                                                                                                                                                                                                                                                                                                                     |
| Sacaline, Broom millet, Millet                                                                                         | Cape tulip, Dock, Saffron thistle, Indian hedge mustard, London rocket, Lupin, Rapistrum, Radish, Wild turnip | WA only                | 1.1 L/ha        | Spray when crop is 10 - 30 cm high and secondary roots have developed and before tasseling. Apply as direct spray to weeds.                                                                                                                                                                                                                         |
| Harvest Aid or Salvage Spray - Winter Cereals                                                                          | Desiccate broadleaf weeds                                                                                     | All States             | 1.2 - 1.7 L/ha  | Apply after dough stage.                                                                                                                                                                                                                                                                                                                            |
| Common Stylo forage or seed crops                                                                                      | Refer to Weed Table                                                                                           | Qld only               | 800 mL/ha       | Apply post-emergence when weeds are 3 weeks old and crop is at least 3 weeks old.                                                                                                                                                                                                                                                                   |
| Caribbean Stylo forage or seed crops                                                                                   |                                                                                                               |                        | 0.8 - 1.6 L/ha  | Apply post-emergence when crop is 3 weeks old.                                                                                                                                                                                                                                                                                                      |

**SECTION 2. SUGARCANE, PEANUTS, BANANAS**

| SITUATION & CROP | WEEDS                                                                                                                                                 | RATE               | CRITICAL COMMENTS                                                                                                                                                                                                                                                                                                                                                                  |
|------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Sugarcane        | Bellvine                                                                                                                                              | 280 mL/100L water  | Apply in Spring, using directed spray.                                                                                                                                                                                                                                                                                                                                             |
|                  | Morning glory                                                                                                                                         | 0.56 - 1.1 L/ha    | Apply in Summer using high clearance tractor.                                                                                                                                                                                                                                                                                                                                      |
|                  | Pink convolvulus, Star of Bethlehem                                                                                                                   | 1.1 L/ha           | Apply in Autumn by aircraft.                                                                                                                                                                                                                                                                                                                                                       |
|                  | Bindy eye (Star burr), Blue top, Cobbler's pegs, Fleabanes, Jute, Leucas, Needle burr, Spear thistle, Water primrose, Ipomea vines, Convolvulus vines | 1.8 - 3.5 L/ha     | Add 300 mL of Shirwet® 600 to 100 L of spray mixture. Agitate well.<br><b>DO NOT</b> use on Q63, Q67, Q80 or Q96 Varieties.<br><b>DO NOT</b> use on Q63 or Q67 varieties (at any rate).<br><b>DO NOT</b> use above 1.1 L/ha on Q80, Q96 or H56 varieties.<br>Refer to local Sugar Research Australia (SRA) representative for further information on local variety susceptibility. |
|                  | Chinese mint, Blue snakeweed                                                                                                                          | 3.5 L/ha           |                                                                                                                                                                                                                                                                                                                                                                                    |
| Peanuts          | Broadleaf weeds; except Noogoora burr, Grasses except Mossman burr                                                                                    | 1.8 - 3.6 L/ha     | LOWER RATE: Apply as BAND SPRAY as soon as possible after planting in a 55 cm band.<br>HIGHER RATE: Apply as OVERALL SPRAY after planting and before crop emergence. Some crop damage may occur if heavy rainfalls between application and crop emergence.                                                                                                                         |
| Bananas          | To destroy Banana suckers                                                                                                                             | 160 mL/10 L water  | Inject at the rate of 15 mL per fully grown plant, 10 mL per medium sized plant and 5 mL for small suckers.                                                                                                                                                                                                                                                                        |
|                  |                                                                                                                                                       | 320 mL/100 L water | Allow suckers from corms of treated plants to form broad adult leaves, then spray. Isolated spots may require a second spray.                                                                                                                                                                                                                                                      |

**SECTION 3. PASTURES, NON-AGRICULTURAL, RIGHTS OF WAY, INDUSTRIAL, CONSERVATION TILLAGE, LAWNS**

| SITUATION & CROP                       | WEEDS                                                                                                                                                                                                                                                                                                                                   | STATE                       | RATE                           | CRITICAL COMMENTS                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              |
|----------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------|--------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Pastures and Non-Agricultural          | Refer Weed Table                                                                                                                                                                                                                                                                                                                        | NSW, Qld, SA, ACT, Tas only | 0.56 - 1.7 L/ha                | Pasture legumes including lucerne, clovers and medics may be damaged unless well protected by grasses. Spot spraying is preferred.                                                                                                                                                                                                                                                                                                                                                                                                             |
|                                        | Galvanised burr                                                                                                                                                                                                                                                                                                                         | NSW, ACT only               | 320 mL/100 L water             | Apply to young actively growing weeds. Ensure thorough and even coverage of plants.<br><b>Note:</b> Treated plants need to be burnt to destroy seeds.                                                                                                                                                                                                                                                                                                                                                                                          |
|                                        | Amsinckia, Docks, Bindweed, Caltrop, Flatweed, Spear thistle, Capeweed, Saffron thistle, Mustard, Wild radish, Wild turnip, Annual thistles, Paterson's curse, Heliotrope, Ragwort, Three cornered Jack (Double gee, Spiny emex)                                                                                                        | WA only                     | 1.1 - 2.4 L/ha                 | For pastures not containing legumes. Only seedling docks, spear thistle and saffron thistle will be controlled. SUMMER WEEDS: Use low rate for seedlings, 1.6 - 2.4 L/ha for larger plants. Stock poisoning may occur when grazed after spraying if large amounts present, particularly Heliotrope. WINTER WEEDS: Use low rate for seedlings, 1.6 - 2.4 L/ha for larger plants. If stock present, use spray/grazing rates.                                                                                                                     |
|                                        | Afghan melons                                                                                                                                                                                                                                                                                                                           |                             | 1.6 L/ha +1% crop oil          | Spray when plants are actively growing preferably before flowering or vining.                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
|                                        | Paddy melons                                                                                                                                                                                                                                                                                                                            |                             | 0.8 - 1.1 L/ha                 |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                |
|                                        | Prickly saltwort (Roly poly)                                                                                                                                                                                                                                                                                                            |                             | 1.6 L/ha                       | Spray when plants are small.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |
|                                        | Stinkwort                                                                                                                                                                                                                                                                                                                               |                             | 1.6 - 3.2 L/ha plus surfactant | Best results are obtained when plants are small. Use high rate on larger plants.                                                                                                                                                                                                                                                                                                                                                                                                                                                               |
|                                        | Dove weed                                                                                                                                                                                                                                                                                                                               |                             | 3.2 L/ha                       | Spray after good emergence of seedlings.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
| Pastures, Rights of Way and Industrial | Boxthorn, Boneseed, Hawthorn                                                                                                                                                                                                                                                                                                            | Vic, SA only                | Undiluted                      | Cut stump: Apply or paint undiluted Nufarm Amine 625 to freshly cut stumps                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
|                                        | Groundsel                                                                                                                                                                                                                                                                                                                               | NSW, Qld, ACT, SA only      | 320 mL/100 L water             | HIGH VOLUME: Thoroughly wet plants.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            |
|                                        |                                                                                                                                                                                                                                                                                                                                         | Tas only                    | 240 mL/15 L water              | CUT STUMP: Swab the cut stump immediately. Apply by a pouring can.                                                                                                                                                                                                                                                                                                                                                                                                                                                                             |
|                                        |                                                                                                                                                                                                                                                                                                                                         | Qld, NSW, ACT, SA, Tas only | 2.9 - 3.6 L/ha                 | AERIAL APPLICATION: Spray when Groundsel is actively growing.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
|                                        | Lantana                                                                                                                                                                                                                                                                                                                                 | NSW, Qld, ACT, SA only      | 320 mL/100 L water             | Use a coarse spray with sufficient pressure to penetrate canopy and wet stems as well as foliage. Spray at the end of a wet Summer (March to May). Defoliation should occur but respraying of new growth will be necessary in following Autumn. Broadcast grass seed and keep stock off following Summer to allow the pasture to establish. Damage may result to pasture legumes.                                                                                                                                                              |
|                                        | Mother of millions                                                                                                                                                                                                                                                                                                                      | NSW only                    | 400 mL/100 L water             | Hand gun only. A thorough coverage of leaves and plantlets is necessary. Use BS1000 at the rate of 1mL of surfactant per 1 L of mixture.                                                                                                                                                                                                                                                                                                                                                                                                       |
|                                        | Noogoora burr, Weir vine (Ipomea), Scarlet pimpernel (seedlings only), White eye (Mexican clover)                                                                                                                                                                                                                                       | Qld only                    | 160 mL/100 L water             | In all cases apply to young, actively growing weeds, ensuring thorough coverage.                                                                                                                                                                                                                                                                                                                                                                                                                                                               |
|                                        | Annual and Perennial pigweed, Artichoke thistle, Bathurst burr, Billygoat weed, Bluesnake weed, Burr medic, Clockweed*, Fleabanes, Galvanised burr, Hemlock, Hoary cress*, Kyalinga weed (Whisker Grass), Knobweed, Milky cotton bushes, Parthenium weed, Paterson's curse, Saffron thistle, Star burr, Thornapple, Variegated thistle* |                             | 320 mL/100 L water             | In all cases apply to young, actively growing weeds, ensuring thorough coverage.<br>* Spray rosette stage.<br>* Repeat spraying necessary.                                                                                                                                                                                                                                                                                                                                                                                                     |
|                                        | Rubber vine                                                                                                                                                                                                                                                                                                                             |                             | 160 mL/10 L water              | Apply to freshly cut stump.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |
|                                        | Sesbania pea                                                                                                                                                                                                                                                                                                                            |                             | 560 - 900 mL/ha                |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                |
|                                        | Water Hyacinth                                                                                                                                                                                                                                                                                                                          |                             | 3.5 - 5.3 L/ha                 | Apply to 2200 - 3300 L water/ha                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                |
|                                        | Wild tobacco tree                                                                                                                                                                                                                                                                                                                       |                             | 240 mL/15 L water              | <b>Cut Stump Treatment:</b> Swab cut stump within 1 hour of cutting. Apply by pouring can.                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
|                                        | <i>Pimelea</i> spp.                                                                                                                                                                                                                                                                                                                     | All states                  | 800 mL/ha + wetter             | Apply by boom spray at 1500 L/ha spray volume. To be applied when plant is green. <b>DO NOT</b> apply more than 2 applications per year with a minimum re-treatment interval of 21 days between consecutive applications. Amine 625 can be used to create and maintain hospital areas for livestock suffering from <i>Pimelea</i> poisoning. <i>Pimelea</i> may become more palatable after herbicide application - stock should be excluded from treated areas until sprayed <i>Pimelea</i> plants are leafless, seedless and obviously dead. |

| SITUATION & CROP                                                             | WEEDS                                                                                                                                                                                                                                                                                                                                                                                                                                              | STATE              | RATE                                      | CRITICAL COMMENTS                                                                                                                                                                                                                                                                                                                                                                                                                                                      |
|------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------|-------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
|                                                                              |                                                                                                                                                                                                                                                                                                                                                                                                                                                    |                    | 55 mL/100 L water + wetter                | Apply by spot spray. Thoroughly wet all foliage to the point of run-off (approximately 1500 L/ha spray volume).                                                                                                                                                                                                                                                                                                                                                        |
| Conservation Tillage - Direct Drilling, Surface Sowing or Fallow Maintenance | Charlock, Mustards, Shepherd's purse, Saffron, Slender, Spear and Variegated thistles*, Turnip weed, Wild radish, Wild turnip                                                                                                                                                                                                                                                                                                                      | All States         | 0.56 - 1.6 L/ha                           | Apply to actively growing young plants. Before sowing: Observe plant back periods given in the table on this leaflet.<br><br>* Warning: Treated plants may become toxic to stock.                                                                                                                                                                                                                                                                                      |
|                                                                              | Clover, Sorrel                                                                                                                                                                                                                                                                                                                                                                                                                                     |                    | 1.1 L/ha + 187 - 267 mL/ha of Kamba® 750  | Apply to actively growing plants in Autumn. <b>DO NOT</b> sow pasture seed for at least 30 days after application.                                                                                                                                                                                                                                                                                                                                                     |
| Lucerne stand reduction – Fallow commencement                                | Lucerne                                                                                                                                                                                                                                                                                                                                                                                                                                            | All states         | 2.4 L/ha                                  | Spray in spring when lucerne is actively growing using a minimum spray volume of 50 L/ha. Heavily graze lucerne in winter and early spring to reduce crown and root reserves. Allow lucerne to regrow to 15 - 30 cm before spraying. Successful lucerne stand reduction is more likely if >70 mm of rain falls in the 6 to 8 weeks prior to application. Add either 0.5% Uptake spray oil or a non-ionic surfactant. Maximum air temperature should not exceed 30° C.  |
| Pastures – Spray Graze Techniques                                            | # PRECAUTION. An increased quantity of poisonous plants may be eaten by stock using Spray-Graze eg. Caltrop, Capeweed, Paterson's curse and Variegated thistle and deaths could result from causes such as nitrate poisoning. With Paterson's curse, preferably graze stock soon destined for slaughter and avoid extended periods of grazing. Avoid grazing with young or breeding stock. <b>DO NOT</b> graze horses or pigs on Paterson's curse. |                    |                                           |                                                                                                                                                                                                                                                                                                                                                                                                                                                                        |
|                                                                              | Amsinckia, Thistles, Capeweed, Double gee, Mustard, Paterson's curse, Wild turnip, Wild radish, Docks, Geranium, Erodium                                                                                                                                                                                                                                                                                                                           | SA only            | 560 mL/ha                                 | Apply from 6 weeks after opening rains in Autumn until the end of August. Seven days after spraying stock paddock at 4-5 times normal rate, preferably with sheep. Maintain this level of grazing for 6 weeks or until pasture shows signs of over grazing. Then return to normal stocking levels. Use high stocking rates in following Spring to prevent weeds from flowering. Repeat treatments maybe required for 2-3 years for complete control.                   |
|                                                                              | Annual thistles, Capeweed, Double gee, Mustards, Paterson's curse, Turnip, Saffron thistle, Spear thistle, Geranium, Slender thistle                                                                                                                                                                                                                                                                                                               | Tas, Vic only      |                                           |                                                                                                                                                                                                                                                                                                                                                                                                                                                                        |
|                                                                              | Amsinckia, Docks (Seedling only), Capeweed, Double gee, Mustard, Wild radish, Wild turnip, Paterson's curse, Annual thistles                                                                                                                                                                                                                                                                                                                       | WA only            | 600 - 800 mL/ha                           |                                                                                                                                                                                                                                                                                                                                                                                                                                                                        |
|                                                                              | Spear thistle, Saffron thistle                                                                                                                                                                                                                                                                                                                                                                                                                     |                    | 1.2 L/ha                                  | Apply to Saffron thistle at the end of September when plants are running up to flower. Sub. clovers may be damaged at this rate and use is not recommended for all medic pastures.                                                                                                                                                                                                                                                                                     |
|                                                                              | Melons                                                                                                                                                                                                                                                                                                                                                                                                                                             |                    | 1.6 L/ha + 1% oil                         | Heavy stocking on young plants sprayed with 800mL/ha provides effective control.                                                                                                                                                                                                                                                                                                                                                                                       |
|                                                                              | Docks                                                                                                                                                                                                                                                                                                                                                                                                                                              | Vic only           | 1.1 L/ha                                  | Apply in September only and follow other recommendations above.                                                                                                                                                                                                                                                                                                                                                                                                        |
|                                                                              | Caltrop, Capeweed, Charlock, Mustards, Paterson's curse, Shepherd's purse, Saffron, Slender, Spear or Variegated thistle*, Turnip weed, Wild radish, Wild turnip                                                                                                                                                                                                                                                                                   | NSW, ACT only      | 0.28 - 1.1 L/ha                           | Spray actively growing 6-8 week old weeds. Introduce stock 7-10 days after spraying, preferably sheep (cattle are less effective). Stocking rate should be at least 5 times heavier than normal until weeds have been reduced, but before survival of desirable pasture species is threatened. Lucerne and medics may be damaged and should be grazed short before spraying. Other legumes may be affected.<br><br>*Warning: Treated plants may become toxic to stock. |
| Lawns, Playing fields                                                        | Refer Weed Table                                                                                                                                                                                                                                                                                                                                                                                                                                   | Qld, NSW, ACT only | 1.5 - 3 mL/1 L of water or 1.6 - 3.2 L/ha | Wet foliage thoroughly. <b>DO NOT</b> mow lawn for 1 week before and at least 1 week after application. <b>DO NOT</b> use on Buffalo grass (WA only).                                                                                                                                                                                                                                                                                                                  |
|                                                                              |                                                                                                                                                                                                                                                                                                                                                                                                                                                    | WA only            | 40 mL/10 - 15 L water/100 m <sup>2</sup>  |                                                                                                                                                                                                                                                                                                                                                                                                                                                                        |
| Hardwood and softwood plantations                                            | Broadleaf and woody weeds as per Weed Table including groundsel and <i>pinus</i> spp. wildlings.                                                                                                                                                                                                                                                                                                                                                   | All states         | Up to 1.6 L/ha                            | Apply a single pre-plant application and/or a maximum of 2 post-plant applications using shielded sprayers within the first 2 years following planting. Apply using aircraft (rotary wing only) or ground based equipment. <b>DO NOT</b> spray over or into watercourses. Amine 625 may be mixed with glyphosate for pre-plant applications.                                                                                                                           |

| SITUATION & CROP | WEEDS               | STATE | RATE            | CRITICAL COMMENTS                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            |
|------------------|---------------------|-------|-----------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Oil tea tree     | Refer to Weed Table |       | Up to 960 mL/ha | <p>Apply as a shielded spray. Avoid contact with foliage, green stems, exposed non-woody roots, desirable plants and trees as severe injury or destruction may result.</p> <p>Apply following harvest as a blanket spray only after:</p> <ul style="list-style-type: none"> <li>• All residual tea tree foliage has been removed by mechanical shaving or by using a burner,</li> <li>• No swollen buds are present on stumps. Note that buds can burst 8 days after harvest in summer, and</li> <li>• Surface of cut stumps are dry prior to spraying commences.</li> </ul> |

**SECTION 4. SPOT SPRAYING (ALL STATES)**

| SITUATION & CROP     | WEEDS               | CRITICAL COMMENTS                                                                                                                                                            |
|----------------------|---------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| High Volume Spraying | Refer to Weed Table | Add 1/10th of rate on weed table to 150 L of water. Each 150 L of mix will cover 1000 m <sup>2</sup> (1/10th ha). eg: If rate in weed table is 1.4 L use 140 mL/150 L water. |

**SECTION 5. USAGE RESTRICTIONS IN SUGARCANE, PASTURE AND PEANUT****Timing restriction for spraying in sugarcane**

| Situation                                          | Region                      | Timing Restriction (DO NOT APPLY DURING THE MONTHS) |                       |                                   |                                 |
|----------------------------------------------------|-----------------------------|-----------------------------------------------------|-----------------------|-----------------------------------|---------------------------------|
|                                                    |                             | Up to 1.2 L/ha                                      | Up to 1.8 L/ha        | Up to 2.6 L/ha                    | Up to 3.5 L/ha                  |
| <b>No trash blanket present during application</b> | Wet tropics & Baron (upper) | No timing restriction                               | No timing restriction | No timing restriction             | October to December             |
|                                                    | Burdekin & Baron (lower)    | No timing restriction                               | No timing restriction | October                           | September to October            |
|                                                    | Mackay/Whitsunday           | No timing restriction                               | October to November   | September to December             | August to December              |
|                                                    | Mary/Burnett                | No timing restriction                               | October to November   | April to May & August to December | April to January                |
|                                                    | Northern NSW & Rocky Point  | No timing restriction                               | No timing restriction | No timing restriction             | October to November             |
| <b>Trash blanket is present during application</b> | Wet tropics & Baron (upper) | No timing restriction                               | No timing restriction | No timing restriction             | November                        |
|                                                    | Burdekin & Baron (lower)    | No timing restriction                               | No timing restriction | October                           | October                         |
|                                                    | Mackay/Whitsunday           | No timing restriction                               | October               | October to November               | September to December           |
|                                                    | Mary/Burnett                | No timing restriction                               | October               | May & October to November         | April to May & July to December |
|                                                    | Northern NSW & Rocky Point  | No timing restriction                               | No timing restriction | No timing restriction             | October to November             |

**Application and timing restrictions for application to pastures**

| Situation                                               | State                 | DO NOT apply above maximum rate (L/ha) below |        |        |        |
|---------------------------------------------------------|-----------------------|----------------------------------------------|--------|--------|--------|
|                                                         |                       | Summer                                       | Autumn | Winter | Spring |
| <b>Pastures (prior to sowing, conservation tillage)</b> | Queensland & NT       | 5.1                                          | 5.1    | 5.1    | 5.1    |
|                                                         | New South Wales & ACT | 5.1                                          | 5.1    | 5.1    | 5.1    |
|                                                         | Victoria              | 0.5                                          | 1.6    | 5      | 1.6    |
|                                                         | Tasmania              | 0.5                                          | 1.2    | 3.5    | 1.6    |
|                                                         | South Australia       | 1.1                                          | 1.6    | 5      | 3.5    |
|                                                         | Western Australia     | 1.6                                          | 3.5    | 5      | 3.5    |
| <b>Pastures (established)</b>                           | Queensland & NT       | 5.3                                          | 5.3    | 5.3    | 5.3    |
|                                                         | New South Wales & ACT | 5.3                                          | 5.3    | 5.3    | 5.3    |
|                                                         | Victoria              | 1                                            | 2      | 5.3    | 3.6    |
|                                                         | Tasmania              | 0.7                                          | 1.7    | 5      | 3.1    |
|                                                         | South Australia       | 1.4                                          | 3.2    | 5.3    | 5.1    |
|                                                         | Western Australia     | 3.6                                          | 5      | 5.3    | 5.1    |

## Timing restrictions for spraying peanuts

| Situation                                     | Rate (L/ha)    | Region             | Timing Restriction: DO NOT APPLY DURING THE MONTHS |
|-----------------------------------------------|----------------|--------------------|----------------------------------------------------|
| Broadcast spraying, prior to sowing (peanuts) | Up to 1.4 L/ha | Cape York          | October and November                               |
|                                               |                | Northern Gulf      | October and November                               |
|                                               |                | Northern Territory | October and November                               |
|                                               |                | Wet Tropics        | No timing restrictions                             |
|                                               |                | Burdekin           | October                                            |
|                                               |                | Mackay/Whitsunday  | September to December                              |
|                                               |                | Mary/Burnett       | October to November                                |
|                                               |                | SE Queensland      | August to May                                      |
|                                               | Up to 1.7 L/ha | Cape York          | October and November                               |
|                                               |                | Northern Gulf      | October and November                               |
|                                               |                | Northern Territory | October and November                               |
|                                               |                | Wet Tropics        | No timing restrictions                             |
|                                               |                | Burdekin           | October                                            |
|                                               |                | Mackay/Whitsunday  | August to December                                 |
|                                               |                | Mary/Burnett       | September to November                              |
|                                               |                | SE Queensland      | <b>Use not supported</b>                           |
| Band spray post-sowing pre-emergence          | Up to 1.8 L/ha | Queensland dryland | No timing restrictions                             |
|                                               |                | Cape York          | No timing restrictions                             |
|                                               |                | Northern Gulf      | October and November                               |
|                                               |                | Northern Territory | October and November                               |
|                                               |                | Wet Tropics        | No timing restrictions                             |
|                                               |                | Burdekin           | No timing restrictions                             |
|                                               |                | Mackay/Whitsunday  | No timing restrictions                             |
|                                               |                | Mary/Burnett       | No timing restrictions                             |
|                                               |                | SE Queensland      | October to January                                 |
| Broadcast spray post-sowing pre-emergence     | Up to 3.6 L/ha | Queensland dryland | June to August                                     |
|                                               |                | Cape York          | October and November                               |
|                                               |                | Northern Gulf      | October and November                               |
|                                               |                | Northern Territory | October and November                               |
|                                               |                | Wet Tropics        | October and November                               |
|                                               |                | Burdekin           | September and October                              |
|                                               |                | Mackay/Whitsunday  | August to December                                 |
|                                               |                | Mary/Burnett       | April to January                                   |
|                                               |                | SE Queensland      | <b>Use not supported</b>                           |

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

IN TASMANIA, THIS PRODUCT MAY ONLY BE USED FROM 15 APRIL TO 15 SEPTEMBER UNLESS OTHERWISE PERMITTED BY THE REGISTRAR OF PESTICIDES.

## WITHHOLDING PERIODS:

## HARVEST:

CROP HARVEST WITHHOLDING PERIOD: NOT REQUIRED WHEN USED AS DIRECTED.

## GRAZING:

PASTURE, CEREAL CROPS, PIMELEA: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION.

LUCERNE (AFTER FALLOW COMMENCEMENT APPLICATION): DO NOT GRAZE, CUT OR CULTIVATE FOR AT LEAST 21 DAYS AFTER SPRAYING.

## WEED TABLE

**NOTE:** Where weeds are to be sprayed in a CROP or PASTURE use only the rates given for the particular crop or situation indicated under the Directions for Use table. In most cases this will give control, however some hard to kill weeds or those in advanced stages of growth may only be suppressed. The rates listed in the Weed Table below are for use where weeds only are present and no crop or pasture is involved. Use at these rates in a crop or pasture other than for spot spraying may cause damage. This product should only be used in those states where a rate or range of rates is indicated for the particular weeds listed.

| Weeds           | Application Rate per Hectare |              |           |        |     |       |                        | Critical Comments                           |
|-----------------|------------------------------|--------------|-----------|--------|-----|-------|------------------------|---------------------------------------------|
|                 | Crop                         |              |           |        |     |       | Pasture                |                                             |
|                 | Vic                          | NSW, ACT     | SA        | Qld    | Tas | WA    | NSW, SA, Qld, Tas only |                                             |
| Amaranthus spp. | -                            | 0.56 - 1.1 L | -         | 900 mL | -   | -     | -                      | Spray young plants.                         |
| Amsinckia       | -                            | -            | -         | -      | -   | 1.1 L | -                      |                                             |
| Apple of Peru   | -                            | 0.56 - 1.1 L | -         | 900 mL | -   | -     | -                      | Spray young plants. Susceptible when young. |
| Bathurst burr   | -                            | 0.8 -1.1 L   | 1.7-2.2 L | 900 mL | -   | -     | 0.8 - 1.1 L not SA     | Spray seedlings only.                       |

| Weeds                         | Application Rate per Hectare |              |              |             |       |              |                        | Critical Comments                                                                  |
|-------------------------------|------------------------------|--------------|--------------|-------------|-------|--------------|------------------------|------------------------------------------------------------------------------------|
|                               | Crop                         |              |              |             |       |              | Pasture                |                                                                                    |
|                               | Vic                          | NSW, ACT     | SA           | Qld         | Tas   | WA           | NSW, SA, Qld, Tas only |                                                                                    |
| Bellvine                      | -                            | 1.8 L        | -            | 1.8 L       | -     | -            | -                      | Spray before seeding. Advanced stages susceptible.                                 |
| Billygoat weed                | -                            | 3.4 L        | -            | 3.4 L       | -     | -            | -                      | Spray at young stage.                                                              |
| Bindweed                      | -                            | -            | -            | -           | -     | 1.1 L        | -                      |                                                                                    |
| Blackberry nightshade         | -                            | 0.56 - 1.1 L | -            | 900 mL      | -     | -            | -                      |                                                                                    |
| Blackeyed Susan               | -                            | 1.8 L        | -            | 1.8 L       | -     | -            | -                      | Apply at pre-flowering, preferably young stages.                                   |
| Blue snakeweed                | -                            | 1.8 L        | -            | 1.8 L       | -     | -            | -                      | Spray seedlings at young stages only.                                              |
| California burr               | -                            | 0.8 -1.1 L   | -            | 900 mL      | -     | -            | 0.8 - 1.1 L not SA     | Spray seedlings only                                                               |
| Cape tulip                    | -                            | -            | -            | -           | -     | 0.64 - 1.3 L | -                      | Low rate for cormils only                                                          |
| Capeweed                      | 1.4 L                        | -            | 1.7 L        | -           | 1.4 L | 1.1 L        | 1.7 - 2.8 L            | Spray seedlings to rosette stage.                                                  |
| Caltrop                       | -                            | 0.8 - 1.7 L  | -            | 900 mL      | -     | 1.1 L        | -                      | Moderately susceptible.                                                            |
| Castor oil plant              | -                            | 3.4 L        | -            | 3.4L        | -     | -            | -                      | Spray at young stage.                                                              |
| Charlock                      | 560-800 mL                   | 0.56 - 1.1 L | 560 mL       | -           | 1.4 L | -            | 0.8 - 1.1 L            | Spray at rosette stage.                                                            |
| Clover                        | -                            | 1.2 L        | -            | -           | -     | -            | -                      |                                                                                    |
| Cobbler's pegs                | -                            | 1.8 L        | -            | 1.8 L       | -     | -            | -                      | Apply at pre-flowering, preferably young stages.                                   |
| Common ice plant              | -                            | -            | 1.1 L        | -           | -     | -            | -                      |                                                                                    |
| Common sida                   | -                            | 1.8 L        | -            | 1.8 L       | -     | -            | -                      | Spray seedling or young stages only.                                               |
| Common sowthistle             | -                            | 1.8 L        | -            | 1.8 L       | 1.4 L | -            | -                      | Apply at pre-flowering, preferably young stages.                                   |
| Docks                         | 1.4 L                        | -            | 1.4 L        | 1.4 L       | 1.4 L | 1.1 L        | 3.2 L SA only          | Spray at multiple leaf stage. Effective only on seedlings.                         |
| Doveweed                      | -                            | -            | -            | -           | -     | 1.1 L        | -                      |                                                                                    |
| Fat hen                       | -                            | 0.56 - 1.7 L | -            | 900 mL      | 1.4 L | -            | -                      | Spray pre-flowering.                                                               |
| Flannel weed                  | -                            | 1.8 L        | -            | 1.8 L       | -     | -            | -                      | Spray seedling or young stages only.                                               |
| Flat weed                     | -                            | -            | -            | -           | -     | 1.1 L        | -                      |                                                                                    |
| Fumitory - red                | -                            | -            | 1.7 L        | -           | -     | -            | -                      |                                                                                    |
| Fumitory - white              | 800 mL                       | -            | 560mL        | -           | -     | -            | -                      | Spray at multiple leaf stage.                                                      |
| Heliotrope                    | -                            | -            | -            | -           | -     | 1.1 L        | -                      |                                                                                    |
| Hexham scent or Melilotus     | 1.4 L                        | -            | 1.1 L        | 1.4 L       | -     | -            | 1.1-1.7L               | Spray multiple leaf stage before seeding.                                          |
| Hoary cress                   | 0.9 - 1.4 L                  | 1.1 - 1.7 L  | 1.4 L        | 1.4 L       | -     | -            | 1.4 - 1.7 L            | Spray rosettes and pre-flowering.                                                  |
| Hogweed/Wireweed              | 1.4L                         | -            | -            | 1.4 L       | -     | -            | -                      | Spray at multiple leaf stage (Vic). Spray at seedling and young plant stage (Qld). |
| Horehound                     | -                            | -            | 1.4 L        | -           | -     | -            | 2.2 - 3.2 L SA only    | Spray seedlings.                                                                   |
| Indian hedge mustard          | -                            | -            | -            | -           | 1.4 L | 1.1 L        | -                      |                                                                                    |
| Khaki weed                    | -                            | -            | -            | -           | -     | -            | 1.1 - 2.2 L not SA     | Spray seedlings only.                                                              |
| Knobweed                      | -                            | -            | -            | 1.8 - 3.4 L | -     | -            | -                      | Lower rate for seedlings; higher rate for later stages.                            |
| Lincoln weed                  | -                            | -            | 1.7 L        | -           | -     | -            | -                      | Spray early rosettes.                                                              |
| London rocket                 | -                            | -            | -            | -           | -     | 1.1 L        | -                      |                                                                                    |
| Lupins                        | -                            | 0.8 - 1.7 L  | -            | -           | -     | 1.1 L        | -                      |                                                                                    |
| Melons - Camel, Paddy         | -                            | 0.56 - 1.1 L | -            | -           | -     | -            | -                      |                                                                                    |
| Mexican poppy                 | -                            | -            | -            | 1.4 L       | -     | -            | -                      | Spray seedlings – plants become more resistant with age.                           |
| Mintweed                      | -                            | 1.1 L        | -            | 900 mL      | -     | -            | -                      | Spray seedlings – resistant in later stages.                                       |
| Morning glory                 | -                            | 1.8 L        | -            | 1.8 L       | -     | -            | -                      | Spray at seedling to flowering stage.                                              |
| Mustards                      | 225-560 mL                   | 0.56 - 1.1 L | 0.56 - 1.4 L | 900 mL      | -     | 800 mL       | 0.56 - 1.1 L           | Spray at 2-4 leaf up to rosette stage.                                             |
| Needle burr                   | -                            | 1.8 L        | -            | 1.8 L       | -     | -            | -                      | Apply at pre-flowering, preferably young stages.                                   |
| New Zealand spinach           | -                            | 1.1 - 1.7 L  | -            | -           | -     | -            | -                      |                                                                                    |
| Noogoora burr                 | -                            | 0.8 - 1.1 L  | -            | 900 mL      | -     | -            | 0.8 - 1.1 L not SA     | Spray seedlings only.                                                              |
| Paterson's curse              | -                            | 1.1 - 1.7 L  | -            | 1.4 L       | -     | 1.3 L        | 1.7 - 2.2 L            | Spray rosettes or before plants have 10 leaves. Later stages harder to kill.       |
| Pinkburr (Pink flowered burr) | -                            | 1.8 L        | -            | 1.8 L       | -     | -            | -                      | Spray seedling or young stages only.                                               |
| Potato weed                   | -                            | 0.56 - 1.1 L | -            | 900 mL      | -     | -            | -                      |                                                                                    |

| Weeds                              | Application Rate per Hectare |              |        |        |        |        |                               | Critical Comments                                                                       |
|------------------------------------|------------------------------|--------------|--------|--------|--------|--------|-------------------------------|-----------------------------------------------------------------------------------------|
|                                    | Crop                         |              |        |        |        |        | Pasture                       |                                                                                         |
|                                    | Vic                          | NSW, ACT     | SA     | Qld    | Tas    | WA     | NSW, SA, Qld, Tas only        |                                                                                         |
| Purpletop                          | -                            | 3.4 L        | -      | 3.4 L  | -      | -      | -                             | Spray at young stage.                                                                   |
| Radish                             | -                            | -            | -      | -      | -      | 1.1 L  | -                             |                                                                                         |
| Ragwort                            | -                            | -            | -      | -      | 3.8 L  | 1.1 L  | -                             | Spray up to early rosette stage (Tas only).                                             |
| Rapistrum                          | -                            | -            | -      | -      | -      | 1.1 L  | -                             |                                                                                         |
| Rough poppy                        | -                            | 1.1 L        | -      | -      | -      | -      | -                             |                                                                                         |
| Safflower                          | -                            | 0.56 - 1.1 L | -      | -      | -      | -      | -                             |                                                                                         |
| Shepherd's purse                   | -                            | 1.1 - 1.7 L  | -      | -      | 1.4 L  | -      | 0.8 - 1.1 L                   | Spray young rosettes.                                                                   |
| Siratro (Purple bean)              | -                            | 1.8 L        | -      | 1.8 L  | -      | -      | -                             | Spray seedling or young stages only.                                                    |
| Skeleton weed                      | 1.4 L                        | 1.1 - 1.7 L  | 1.4 L  | -      | -      | -      | -                             | Spray rosettes before aerial growth commences                                           |
| Sorrel                             | 1.4 L                        | 1.7 L        | 1.4 L  | -      | -      | -      | -                             | Only moderately susceptible.                                                            |
| Speedwell - Ivy leaf               | -                            | -            | 1.1 L  | -      | -      | -      | -                             |                                                                                         |
| Spinyhead sida                     | -                            | 1.8 L        | -      | 1.8 L  | -      | -      | -                             | Spray seedling or young stages only                                                     |
| Starburr                           | -                            | 1.8 L        | -      | 1.8 L  | -      | -      | -                             | Spray before seeding, advanced stages susceptible.                                      |
| Spiny emex                         | -                            | -            | -      | 1.4 L  | -      | -      | -                             | Only young plants are susceptible.                                                      |
| Star of Bethlehem (Cupid's flower) | -                            | -            | -      | 1.8 L  | -      | -      | -                             | Spray before seeding, advanced stages susceptible                                       |
| Stinkwort                          | -                            | 0.8 - 1.4 L  | -      | -      | -      | -      | -                             |                                                                                         |
| Storksbill/Erodium                 | -                            | -            | -      | -      | 1.4 L  | -      | 1.6 - 3.2 L                   | Spray seedlings to young rosettes.                                                      |
| Sunflower (seedlings)              | 1.4 L                        | 0.56 - 1.4 L | -      | 900 mL | -      | -      | -                             |                                                                                         |
| Thistles: - Annual                 | -                            | -            | -      | -      | -      | 1.1 L  | -                             |                                                                                         |
| -Californian                       | -                            | -            | -      | -      | 2.7 L  | -      | 3.2 - 3.8 L                   | Repeated applications may be necessary. (NSW, ACT, Tas only)                            |
| -Saffron                           | 1.1 L                        | 0.56 - 1.7 L | 1.4 L  | 1.4 L  | 950 mL | 1.1 L  | 1.1 - 1.7 L                   | Low rate only sufficient to control weeds in crops at rosette stage when sprayed early. |
| -Slender/Shore                     | -                            | 0.8 - 1.7 L  | -      | -      | 1.4 L  | -      | 1.1 L                         | Suppression only.                                                                       |
| -Soldier                           | 1.4 L                        | -            | -      | -      | -      | -      | 1.1 - 1.6 L not NSW, ACT, Tas | Spray young rosette.                                                                    |
| -Spear                             | 560 mL                       | -            | -      | -      | 1.4 L  | -      | 1.1 - 1.6 L                   | Spray young rosettes.                                                                   |
| -Star                              | -                            | -            | -      | -      | -      | -      | 1.6 - 3.2 L SA only           | Use higher rate as flower stalk appears.                                                |
| -Variegated                        | -                            | 0.56 - 1.7 L | -      | 900 mL | 1.4 L  | -      | 1.1 - 1.7 L                   | Spray at rosette stage.                                                                 |
| Thornapple                         | -                            | 0.8 - 1.1 L  | -      | -      | -      | -      | 1.6 - 2.4 L not SA            | Spray seedlings only.                                                                   |
| Tridax (Tridax daisy)              | -                            | 1.8 L        | -      | 1.8 L  | -      | -      | -                             | Spray seedling or young stages only.                                                    |
| Turnip weed/Rapistrum              | -                            | 0.56 - 1.1 L | -      | 560 mL | -      | 800 mL | 0.56 - 1.1 L                  |                                                                                         |
| Vetches/Tares                      | 1.4 L                        | -            | 1.1 L  | -      | -      | -      | -                             | Spray at multiple leaf stage.                                                           |
| Ward's weed                        | -                            | -            | 1.1 L  | -      | -      | -      | -                             |                                                                                         |
| Wild cabbage                       | 1.4 L                        | -            | -      | -      | -      | -      | -                             | Spray multiple leaves.                                                                  |
| Wild poppy                         | 560 mL                       | -            | -      | -      | -      | -      | 1.1 - 1.7 L                   | Spray rosettes.                                                                         |
| Wild radish                        | 1.4 L                        | 1.4 - 1.7 L  | 1.4 L  | 900 mL | 1.4 L  | 800 mL | 0.8 - 1.1 L                   | Spray up to young rosette stage.                                                        |
| Wild turnip                        | 225 - 560 mL                 | 0.56 - 1.1 L | 320 mL | -      | 1.4 L  | 640 mL | 0.56 - 1.1 L                  | Spray 2-4 leaf up to rosette stage.                                                     |

**GENERAL INSTRUCTIONS**

Before opening, carefully read Directions for Use, Precautionary Statements, Safety Directions and First Aid Instructions.

**CROP SAFETY AND ESTABLISHMENT****Crop Tolerance**

| MAXIMUM TOLERANCE FOR USE IN WHEAT, BARLEY, OATS AND TRITICALE INCLUDING TOLERANCE FOR UNDERSOWN LEGUMES. |          |           |                 |                     |           |                           |                           |
|-----------------------------------------------------------------------------------------------------------|----------|-----------|-----------------|---------------------|-----------|---------------------------|---------------------------|
| Crop                                                                                                      | Qld      | NSW, ACT  | Vic             | Vic                 | SA        | WA                        | Tas                       |
|                                                                                                           |          |           | Early Tillering | Tillered Boot Stage |           | Tillered to Boot (Z15-35) | Tillered to Boot (Z15-35) |
| Cereal rye                                                                                                | -        | -         | -               | -                   | 1.4 L/ha  | -                         | -                         |
| Wheat                                                                                                     | 1.8 L/ha | 1.7 L/ha  | 225 mL/ha       | 1.4 L/ha            | 1.7 L/ha  | 1.7 L/ha                  | 1.7 L/ha                  |
| Barley                                                                                                    | 1.4 L/ha | 1.4 L/ha  | 225 mL/ha       | 1.4 L/ha            | 1.4 L/ha  | 1.4 L/ha                  | 1.4 L/ha                  |
| Oats                                                                                                      | -        | 800 mL/ha | 225 mL/ha       | 800 mL/ha           | 900 mL/ha | 1.3 L/ha                  | -                         |
| Triticale                                                                                                 | -        | 1.4 L/ha  | 225 mL/ha       | -                   | 1.4 L/ha  | 1.7 L/ha                  | 800 mL/ha                 |
| Undersown Clovers                                                                                         | -        | -         | 225 mL/ha       | -                   | -         | 680 mL/ha                 | 560 mL/ha                 |
| Undersown Medics                                                                                          | -        | -         | 225 mL/ha       | -                   | -         | NIL                       | 560 mL/ha                 |
| Undersown Lucerne                                                                                         | -        | -         | 225 mL/ha       | -                   | -         | NIL                       | -                         |

**Plant Back Periods (days) for Nufarm Amine 625 Selective Herbicide**

| Crop               | Rates           |                 |                |
|--------------------|-----------------|-----------------|----------------|
|                    | Up to 560 mL/ha | 0.56 - 1.1 L/ha | 1.1 - 1.7 L/ha |
| Balansa clover     | 7               | 7               | 10             |
| Barley %           | 1               | 1               | 3              |
| Chickpeas #        | 7               | 14              | 21             |
| Cotton             | 10              | 14              | 21             |
| Faba beans         | 7               | 7               | 10             |
| Field peas         | 7               | 14              | 14             |
| Lentils            | 7               | 7               | 10             |
| Linseed            | 7               | 7               | 14             |
| Lucerne            | 7               | 7               | 10             |
| Lupins +           | 7               | 14              | 21             |
| Medics             | 7               | 7               | 10             |
| Narbon beans       | 7               | 7               | 10             |
| Navybean           | 10              | 10              | 14             |
| Oats               | 3               | 3               | 7              |
| Perennial ryegrass | 7               | 7               | 10             |
| Persian clover     | 7               | 7               | 10             |
| Phalaris           | 7               | 7               | 10             |
| Canola/Rapeseed #  | 14              | 21              | 28             |
| Rice               | 7               | 7               | 14             |
| Safflower #        | 7               | 14              | 21             |
| Sorghum @          | 3               | 7               | 10             |
| Soybean            | 14              | 14              | 21             |
| Sub. clover        | 7               | 7               | 10             |
| Sunflower @        | 7               | 10              | 14             |
| Triticale %        | 1               | 3               | 7              |
| Vetch              | 7               | 7               | 10             |
| Wheat %            | 1               | 3               | 7              |
| White clover       | 7               | 7               | 10             |

**IMPORTANT: WHEN APPLIED TO DRY SOILS AT LEAST 15MM (1/2 INCH) OF RAIN MUST FALL PRIOR TO THE COMMENCEMENT OF THE PLANT BACK PERIOD.**

**NOTES:**

% In Queensland, no rainfall is required to fall prior to commencement of Plant Back Period for wheat, barley and triticale.

# In Queensland, planting of canola/rapeseed, chickpeas and safflower must be delayed for at least 14 days following rainfall of at least 15 mm.

@ In Central Queensland, when using 800 mL/ha or less of Nufarm Amine 625, the Plant Back Period for sorghum and sunflower is 1 day irrespective of rainfall.

+ In WA the Plant Back Period for lupins at all rates is 28 days.

**APPLICATION INFORMATION**

BOOM SPRAYING - Use 30 - 120 L/ha of water.

AERIAL SPRAYING - Use 10 - 90 L/ha of water.

**EQUIPMENT MAINTENANCE AND USAGE**

Equipment that has been used for this chemical should not be used for the application of other materials to sensitive plants, unless it has been well washed out with hot soapy water or 1% solution of ammonia, followed by several clear water rinses.

**SPRAY APPLICATIONS AND DRIFT RISK ASSESSMENT**

For aerial application it is recommended where possible for this product to be applied by an aerial applicator business that holds current accreditation for the Aerial Improvement Management System – “AIMS”, issued by the Aerial Agricultural Association of Australia Ltd.

**Checklist:**

- Have you cleaned/decontaminated your boom sprayer?
- Have you contacted your neighbour prior to spraying?
- Is your sprayer set-up correctly for the particular application?
- Check
  - boom calibration
  - at nozzle - nozzle choice
  - low drift/what spray quality
  - very coarse or larger spray quality?
  - boom height - speed of intended application
  - water volume
- You must check, determine and record the weather conditions immediately prior to, and immediately after the spray application is made.
- Record
  - Temperatures
  - Relative Humidity
  - Delta T
  - Wind speed
  - Is there a temperature inversion?
- Night Spraying - Extra care is required to ensure that inversion conditions are not present. Use smoke generator to determine wind direction and presence of inversion conditions.

For further information refer to [nufarm.com.au/spraywise](http://nufarm.com.au/spraywise)



[spraywisedecisions.com.au](http://spraywisedecisions.com.au) is an online weather forecasting program and is recommended for use when planning your pesticide application.

When spraying in or near areas with sensitive crops such as cotton, check online at [satacrop.com.au](http://satacrop.com.au) for the proximity of sensitive crops mapped by growers.

#### COMPATIBILITY

This product may be tank mixed with Atradex® WG, Kamba® 750, Diurex® WG, Glean®, Revolver®, Shirquat® 250, Spray•Seed®, Throttle®, Trooper 75-D.

#### RESISTANT WEEDS WARNING

| GROUP | I | HERBICIDE |
|-------|---|-----------|
|-------|---|-----------|

Nufarm Amine 625 Selective Herbicide is a member of the Phenoxy group of herbicides. The product has the disruptors of plant cell growth mode of action. For weed resistance management the product is a Group I herbicide. Some naturally occurring weed biotypes resistant to the product and other Group I herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by this product or other Group I herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Nufarm Australia Limited accepts no liability for any losses that may result from the failure of Nufarm Amine 625 Selective Herbicide to control resistant weeds.

#### PRECAUTIONS

##### RE-ENTRY PERIOD

**DO NOT** hand harvest sugar cane for at least 1 day after application.

If re-entering treated areas before the spray has dried, workers should wear overalls, elbow-length gloves and water-resistant footwear. Clothing Should be laundered after each days use.

##### PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

**DO NOT** spray cereals if lucerne is present.

**DO NOT** spray crops or weeds outside the stages indicated in "Critical Comments" as damage, loss of yield or inadequate weed control may result.

**DO NOT** apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants, crops, cropped lands or pastures.

Avoid spray drift onto susceptible crops such as cotton, tobacco, tomatoes, vines, lupins, fruit trees and ornamentals.

##### PROTECTION OF LIVESTOCK

Low hazard to bees. May be applied at any time as recommended in the Directions for Use.

##### PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

**Very toxic to aquatic life. DO NOT** contaminate streams, rivers or waterways with the chemical or used container.

##### STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. **DO NOT** store for prolonged periods in direct sunlight.

##### Non-Refillable containers

Triple-rinse containers before disposal. Add rinsings to spray tank. **DO NOT** dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. **DO NOT** burn empty containers or product.

##### Refillable containers

Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

#### SAFETY DIRECTIONS

Poisonous if absorbed by skin contact, inhaled or swallowed. Corrosive to the eyes and skin. Will irritate the nose and throat. Avoid contact with the eyes and skin. **DO NOT** inhale vapour or spray mist.

**Mixing and loading with open systems:** When opening the container and preparing spray or using undiluted concentrate, wear chemical resistant waterproof clothing over a layer of normal clothing and a washable hat, elbow-length chemical resistant gloves, impervious footwear and full facepiece respirator with organic vapour/gas cartridge or canister.

**Mixing and loading with closed systems:** When opening the container and preparing the spray, wear cotton overalls over normal clothing, buttoned to the neck and wrist, elbow length chemical resistant gloves and face shield or goggles.

**If applying by boomspray equipment with enclosed operator's cab and air filtration or aerial spraying equipment:** Wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow-length chemical resistant gloves.

**If applying by boomspray equipment with open operator's cab or hand-held spray equipment:** Wear chemical resistant waterproof clothing over a layer of normal clothing and a washable hat, elbow-length chemical resistant gloves and full facepiece respirator with organic vapour/gas cartridge or canister.

**After use:** If clothing becomes contaminated with product remove clothing immediately. If product on skin, immediately wash area with soap and water. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, respirator and if rubber wash with detergent and warm water, face shield or goggles and contaminated clothing.

#### FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766.

#### ADDITIONAL STATEMENTS (WHS REGULATIONS 2011)

**Causes serious eye damage. May cause an allergic skin reaction.** If medical advice is needed, have product container or label at hand. Wear protective gloves, clothing, eye and face protection. **DO NOT** eat, drink or smoke when using this product. Contaminated work clothing should not be allowed out of the

work place. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Immediately call a POISON CENTRE or doctor/physician. IF SWALLOWED: Call a POISON CENTRE or doctor/physician if you feel unwell. Rinse mouth. IF ON SKIN: Wash with plenty of soap and water. If skin irritation or rash occurs: Get medical advice / attention. Wash contaminated clothing before reuse.

**SAFETY DATA SHEET**

For further information refer to the Safety Data Sheet (SDS) which can be obtained from your supplier or from the Nufarm website – [nufarm.com.au](http://nufarm.com.au)

**In case of emergency: Phone 1800 033 498.** Ask for shift supervisor. Toll free 24 hours.

**CONDITIONS OF SALE**

"Any provisions or rights under the Competition and Consumer Act 2010 or relevant state legislation which cannot be excluded by those statutes or by law are not intended to be excluded by these conditions of sale. Subject to the foregoing, all warranties, conditions, rights and remedies, expressed or implied under common law, statute or otherwise, in relation to the sale, supply, use or application of this product, are excluded. Nufarm Australia Limited and/or its affiliates ("Nufarm") shall not accept any liability whatsoever (including consequential loss), or howsoever arising (including negligence) for any damage, injury or death connected with the sale, supply, use or application of this product except for liability which cannot be excluded by statute."

Nufarm Australia Limited  
ACN 004 377 780  
103-105 Pipe Road  
Laverton North Victoria 3026  
Tel: (03) 9282 1000

® All trade marks owned or used under license by Nufarm Australia Limited

\* Other trade marks.

**APVMA Approval No.: 56763/129694**