

# Lecture 3: Age & Stage structured population growth



Population: a set of individuals of the same species co-occurring in space and time.

### Characteristics of populations

Total number of individuals or population 'density'

Age/Stage/Size structure

Sex ratio

Spatial distribution

$$dN/dt = rN$$

$$N_t = N_0 e^{rt}$$

$$\ln(2)/r = t_{double}$$

$$dN/dt = rN \left( 1 - \frac{N}{K} \right)$$

$$dN/dt = rN \left( \frac{K - N}{K} \right)$$

$$N_t = N_0 \frac{K}{(K - N_0)e^{-rt} + N_0}$$

# Four important assumptions

## Four important assumptions

1. Per capita growth rate is a linear function of  $N$ .
2. Population growth rate responds instantly to changes in  $N$  (no time lags).
3. The external environment has no influence on population growth.
4. All individuals are equal (i.e., no age or size effects).

# Four important assumptions

1. Per capita growth rate is a linear function of  $N$ .
2. Population growth rate responds instantly to changes in  $N$  (no time lags).
3. The external environment has no influence on population growth.
4. All individuals are equal (i.e., no age or size effects).

These models work for 'simple' organisms like single-celled bacteria or protozoa.





But for most plants and animals, birth and death rates depend on age.



Northern leopard frog, *Rana pipiens*



A newborn elephant cannot reproduce immediately, but instead must wait ~10 years.





Hatchlings and seedlings often have higher mortality rates than adults.








A population that consisted of only old, post-reproductive chimpanzees will soon go extinct.



A population of only tadpoles cannot begin to grow until tadpoles become adults and can reproduce.







When using age classes, we treat all individuals in the same class as equivalents.



**The fecundity schedule:** the average number of female offspring born per unit time to an individual female of a particular age.

$b(x)$  or  $m(x)$   $\longrightarrow$  births or *m*aternity

**The fecundity schedule:** the average number of female offspring born per unit time to an individual female of a particular age.

$b(x)$  or  $m(x)$   $\longrightarrow$  births or maternity

$b(6) = 3$   $\longrightarrow$  A female of age 6 has,  
on average, 3 female offspring

$x$	$S(x)$	$b(x)$	$\frac{l(x)}{S(x)/S(0)}$	$\frac{g(x)}{l(x+1)/l(x)}$	$l(x)b(x)$	$l(x)b(x)x$	Initial estimate $e^{-rx}l(x)b(x)$
0	500	0					
1	400	2					
2	200	3					
3	50	1					
4	0	0					

# The survivorship schedule

$x$	$S(x)$	$b(x)$	$\frac{l(x)}{= S(x)/S(0)}$	$\frac{g(x)}{= l(x+1)/l(x)}$	$l(x)b(x)$	$l(x)b(x)x$	Initial estimate $e^{-rx}l(x)b(x)$
0	500	0					
1	400	2					
2	200	3					
3	50	1					
4	0	0					

The proportion of the original cohort that survives to the start of that age class

# The survivorship schedule

$x$	$S(x)$	$b(x)$	$\frac{l(x)}{= S(x)/S(0)}$	$\frac{g(x)}{= l(x+1)/l(x)}$	$l(x)b(x)$	$l(x)b(x)x$	Initial estimate $e^{-rx}l(x)b(x)$
0	500	0					
1	400	2					
2	200	3					
3	50	1					
4	0	0					

The proportion of the original cohort that survives to the start of that age class  
 The probability of surviving to that age class



# The survivorship schedule

$x$	$S(x)$	$b(x)$	$\frac{l(x)}{S(x)/S(0)}$	$\frac{g(x)}{= l(x+1)/l(x)}$	$l(x)b(x)$	$l(x)b(x)x$	Initial estimate $e^{-rx}l(x)b(x)$
0	500	0	1				
1	400	2	0.8				
2	200	3	0.4				
3	50	1	0.1				
4	0	0	0				

The proportion of the original cohort that survives to the start of that age class

The probability of surviving to that age class

# The survival probability

$x$	$S(x)$	$b(x)$	$\frac{l(x)}{S(x)/S(0)}$	$\frac{g(x)}{= l(x+1)/l(x)}$	$l(x)b(x)$	$l(x)b(x)x$	Initial estimate $e^{-rx}l(x)b(x)$
0	500	0	1				
1	400	2	0.8				
2	200	3	0.4				
3	50	1	0.1				
4	0	0	0				

The probability that an individual of age  $x$  survives to age  $x + 1$

# The survival probability

$x$	$S(x)$	$b(x)$	$\frac{l(x)}{S(x)/S(0)}$	$\frac{g(x)}{= l(x+1)/l(x)}$	$l(x)b(x)$	$l(x)b(x)x$	Initial estimate $e^{-rx}l(x)b(x)$
0	500	0	1	0.80			
1	400	2	0.8	0.50			
2	200	3	0.4	0.25			
3	50	1	0.1	0			
4	0	0	0				

The probability that an individual of age  $x$  survives to age  $x + 1$

# The net reproductive rate, $R_0$

$x$	$S(x)$	$b(x)$	$\frac{l(x)}{= S(x)/S(0)}$	$\frac{g(x)}{= l(x+1)/l(x)}$	$l(x)b(x)$	$l(x)b(x)x$	Initial estimate $e^{-rx}l(x)b(x)$
0	500	0	1	0.80			
1	400	2	0.8	0.50			
2	200	3	0.4	0.25			
3	50	1	0.1	0			
4	0	0	0				

The mean number of female offspring produced per female over her lifetime.

# The net reproductive rate, $R_0$

$x$	$S(x)$	$b(x)$	$\frac{l(x)}{= S(x)/S(0)}$	$\frac{g(x)}{= l(x+1)/l(x)}$	$l(x)b(x)$	$l(x)b(x)x$	Initial estimate $e^{-rx}l(x)b(x)$
0	500	0	1	0.80	0		
1	400	2	0.8	0.50	1.6		
2	200	3	0.4	0.25	1.2		
3	50	1	0.1	0	0.1		
4	0	0	0				

The mean number of female offspring produced per female over her lifetime.  
 The reproductive potential adjusted by the mortality schedule.



# The net reproductive rate, $R_0$

$x$	$S(x)$	$b(x)$	$\frac{l(x)}{= S(x)/S(0)}$	$\frac{g(x)}{= l(x+1)/l(x)}$	$l(x)b(x)$	$l(x)b(x)x$	Initial estimate $e^{-rx}l(x)b(x)$
0	500	0	1	0.80	0		
1	400	2	0.8	0.50	1.6		
2	200	3	0.4	0.25	1.2		
3	50	1	0.1	0	0.1		
4	0	0	0	$R_0 = \sum l(x)b(x) = 2.9$ offspring			

If  $R_0 > 1$ , there is a net surplus of offspring produced each generation, and the population increases exponentially.

# The net reproductive rate, $R_0$

$x$	$S(x)$	$b(x)$	$\frac{l(x)}{= S(x)/S(0)}$	$\frac{g(x)}{= l(x+1)/l(x)}$	$l(x)b(x)$	$l(x)b(x)x$	Initial estimate $e^{-rx}l(x)b(x)$
0	500	0	1	0.80	0		
1	400	2	0.8	0.50	1.6		
2	200	3	0.4	0.25	1.2		
3	50	1	0.1	0	0.1		
4	0	0	0	$R_0 = \sum l(x)b(x) = 2.9$ offspring			

If  $R_0 < 1$ , the population declines

# The net reproductive rate, $R_0$

$x$	$S(x)$	$b(x)$	$\frac{l(x)}{= S(x)/S(0)}$	$\frac{g(x)}{= l(x+1)/l(x)}$	$l(x)b(x)$	$l(x)b(x)x$	Initial estimate $e^{-rx}l(x)b(x)$
0	500	0	1	0.80	0		
1	400	2	0.8	0.50	1.6		
2	200	3	0.4	0.25	1.2		
3	50	1	0.1	0	0.1		
4	0	0	0	$R_0 = \sum l(x)b(x) = 2.9$ offspring			

If  $R_0 = 1$ , the population is stable.

# Generation time, $G$

$x$	$S(x)$	$b(x)$	$\frac{l(x)}{S(x)/S(0)}$	$\frac{g(x)}{l(x+1)/l(x)}$	$l(x)b(x)$	$l(x)b(x)x$	Initial estimate $e^{-rx}l(x)b(x)$
0	500	0	1	0.80	0		
1	400	2	0.8	0.50	1.6		
2	200	3	0.4	0.25	1.2		
3	50	1	0.1	0	0.1		
4	0	0	0				

Average age of the parents of all the offspring produced by a single cohort

# Generation time, $G$

$$\frac{\sum l(x)b(x)x}{\sum l(x)b(x)}$$

$x$	$S(x)$	$b(x)$	$\frac{l(x)}{S(x)/S(0)}$	$\frac{g(x)}{= l(x+1)/l(x)}$	$l(x)b(x)$	$l(x)b(x)x$	Initial estimate $e^{-rx}l(x)b(x)$
0	500	0	1	0.80	0	0	
1	400	2	0.8	0.50	1.6	1.6	
2	200	3	0.4	0.25	1.2	2.4	
3	50	1	0.1	0	0.1	0.3	
4	0	0	0				
$\Sigma = 4.3$							

Average age of the parents of all the offspring produced by a single cohort



# Generation time, $G$

$$\frac{\sum l(x)b(x)x}{\sum l(x)b(x)}$$

$x$	$S(x)$	$b(x)$	$\frac{l(x)}{= S(x)/S(0)}$	$\frac{g(x)}{= l(x+1)/l(x)}$	$l(x)b(x)$	$l(x)b(x)x$	Initial estimate $e^{-rx}l(x)b(x)$
0	500	0	1	0.80	0	0	
1	400	2	0.8	0.50	1.6	1.6	
2	200	3	0.4	0.25	1.2	2.4	
3	50	1	0.1	0	0.1	0.3	
4	0	0	0				
						$\Sigma = 4.3$	
						$G = 4.3/2.9 = 1.483$ years	

Average age of the parents of all the offspring produced by a single cohort

**Finally, estimating  $r$**

**Finally, estimating  $r$**

$$r \approx \frac{\ln(R_o)}{G}$$

# Finally, estimating $r$

$$r \approx \frac{\ln(R_0)}{G}$$

$$r \approx \frac{\ln(2.9)}{1.483} \approx 0.718 \text{ individuals per individual per year}$$

$$R_0 = \sum l(x)b(x) = 2.9 \text{ offspring}$$

$$G = 4.3/2.9 = 1.483 \text{ years}$$

# Finally, estimating $r$

$$r \approx \frac{\ln(R_0)}{G}$$

$$r \approx \frac{\ln(2.9)}{1.483} \approx 0.718 \text{ individuals per individual per year}$$

As generation time increases, rate of increase decreases; as number of offspring per female increases, rate of increase increases.

$$R_0 = \sum l(x)b(x) = 2.9 \text{ offspring}$$

$$G = 4.3/2.9 = 1.483 \text{ years}$$

# Finally, estimating $r$

$$r \approx \frac{\ln(R_0)}{G}$$

$$r \approx \frac{\ln(2.9)}{1.483} \approx 0.718 \text{ individuals per individual per year}$$

As generation time increases, rate of increase decreases; as number of offspring per female increases, rate of increase increases.

This is just an estimate, but a close estimate (usually within ~10%).

$x$	$S(x)$	$b(x)$	$\frac{l(x)}{=S(x)/S(0)}$	$\frac{g(x)}{=l(x+1)/l(x)}$	$l(x)b(x)$	$l(x)b(x)x$	Initial estimate $e^{-rx}l(x)b(x)$
0	500	0	1	0.80	0	0	0
1	400	2	0.8	0.50	1.6	1.6	0.780
2	200	3	0.4	0.25	1.2	2.4	0.285
3	50	1	0.1	0	0.1	0.3	0.012
4	0	0	0				$\Sigma = 1.077$

# The Euler equation (pronounced “Oiler”)

$$1 = \sum e^{-rx} l(x) b(x)$$

Unfortunately, there is no way to solve this equation except to plug in different values of  $r$ .



$x$	$S(x)$	$b(x)$	$\frac{l(x)}{= S(x)/S(0)}$	$\frac{g(x)}{= l(x+1)/l(x)}$	$l(x)b(x)$	$l(x)b(x)x$	Initial estimate $e^{-rx}l(x)b(x)$
0	500	0	1	0.80	0	0	0
1	400	2	0.8	0.50	1.6	1.6	0.780
2	200	3	0.4	0.25	1.2	2.4	0.285
3	50	1	0.1	0	0.1	0.3	0.012
4	0	0	0				$\Sigma = 1.077$

$$r \approx 0.718$$

$x$	$S(x)$	$b(x)$	$\frac{l(x)}{= S(x)/S(0)}$	$\frac{g(x)}{= l(x+1)/l(x)}$	$l(x)b(x)$	$l(x)b(x)x$	Initial estimate $e^{-rx}l(x)b(x)$
0	500	0	1	0.80	0	0	0
1	400	2	0.8	0.50	1.6	1.6	0.780
2	200	3	0.4	0.25	1.2	2.4	0.285
3	50	1	0.1	0	0.1	0.3	0.012
4	0	0	0				

$\Sigma = 1.077$

$$r \approx 0.718 \quad r_{(\text{Euler})} = 0.776$$

# Describing population age structure

Once we have  $r$  from the life tables, we can forecast the total population size by using any of the equations we've seen before.

# Describing population age structure

Once we have  $r$  from the life tables, we can forecast the total population size by using any of the equations we've seen before.

But we're also interested in age structure of the population.

$n_i(t)$  = number of individuals at time  $t$  in age class  $i$ .

# Describing population age structure

Once we have  $r$  from the life tables, we can forecast the total population size by using any of the equations we've seen before.

But we're also interested in age structure of the population.

$n_i(t)$  = number of individuals at time  $t$  in age class  $i$ .

$n_1(3) = 50$  : 50 individuals in age class 1 at the 3rd time step.

# Describing population age structure

Once we have  $r$  from the life tables, we can forecast the total population size by using any of the equations we've seen before.

But we're also interested in age structure of the population.

$n_i(t)$  = number of individuals at time  $t$  in age class  $i$ .

$n_1(3) = 50$  : 50 individuals in age class 1 at the 3rd time step.

Because there are  $k$  age classes, the age structure at time  $t$  is a vector of abundances.

# Describing population age structure

Once we have  $r$  from the life tables, we can forecast the total population size by using any of the equations we've seen before.

But we're also interested in age structure of the population.

$n_i(t)$  = number of individuals at time  $t$  in age class  $i$ .

$n_1(3) = 50$  : 50 individuals in age class 1 at the 3rd time step.

Because there are  $k$  age classes, the age structure at time  $t$  is a vector of abundances.

$$\mathbf{n}(t) = \begin{pmatrix} n_1(t) \\ n_2(t) \\ \vdots \\ n_k(t) \end{pmatrix}$$

# Describing population age structure

Once we have  $r$  from the life tables, we can forecast the total population size by using any of the equations we've seen before.

But we're also interested in age structure of the population.

$n_i(t)$  = number of individuals at time  $t$  in age class  $i$ .

$n_1(3) = 50$  : 50 individuals in age class 1 at the 3rd time step.

Because there are  $k$  age classes, the age structure at time  $t$  is a vector of abundances.

The population from our life table might look like this at  $t = 5$ .  
600 individuals in the 1st age class, but only 50 in the 4th.

$$\mathbf{n}_5 = \begin{pmatrix} 600 \\ 270 \\ 100 \\ 50 \end{pmatrix}$$



# Describing population age structure

Once we have  $r$  from the life tables, we can forecast the total population size by using any of the equations we've seen before.

But we're also interested in age structure of the population.

$n_i(t)$  = number of individuals at time  $t$  in age class  $i$ .

$n_1(3) = 50$  : 50 individuals in age class 1 at the 3rd time step.

Because there are  $k$  age classes, the age structure at time  $t$  is a vector of abundances.

The population from our life table might look like this at  $t = 5$ .  
600 individuals in the 1st age class, but only 50 in the 4th.

$$\mathbf{n}_5 = \begin{pmatrix} 600 \\ 270 \\ 100 \\ 50 \end{pmatrix}$$

Using the mortality and fertility schedules from the life table, we can prediction how the age structure changes from one time period  $\mathbf{n}(t)$  to the next,  $\mathbf{n}(t+1)$

# Calculating survival probabilities for age classes ( $P_i$ )

$x$	$S(x)$	$b(x)$	$\frac{l(x)}{= S(x)/S(0)}$	$\frac{g(x)}{= l(x+1)/l(x)}$	$l(x)b(x)$	$l(x)b(x)x$	Initial estimate $e^{-\sum l(x)b(x)}$
0	500	0	1	0.80	0	0	0
1	400	2	0.8	0.50	1.6	1.6	0.780
2	200	3	0.4	0.25	1.2	2.4	0.285
3	50	1	0.1	0	0.1	0.3	0.012
4	0	0	0				

$$P_i = \frac{l(i)}{l(i - 1)}$$

# Calculating survival probabilities for age classes ( $P_i$ )

$x$	$S(x)$	$b(x)$	$\frac{l(x)}{= S(x)/S(0)}$	$\frac{g(x)}{= l(x+1)/l(x)}$	$l(x)b(x)$	$l(x)b(x)x$	Initial estimate $e^{-\sum l(x)b(x)}$
0	500	0	1	0.80	0	0	0
1	400	2	0.8	0.50	1.6	1.6	0.780
2	200	3	0.4	0.25	1.2	2.4	0.285
3	50	1	0.1	0	0.1	0.3	0.012
4	0	0	0				

$$P_i = \frac{l(i)}{l(i - 1)}$$

$$n_{i+1}(t + 1) = P_i n_i(t)$$

# Calculating fertility for age classes ( $F_i$ )

$$n_{i+1}(t + 1) = P_i n_i(t)$$

Works for all age classes except the first. Because the number in the first age class depends on the reproduction of all age classes. So...

# Calculating fertility for age classes ( $F_i$ )

$$n_{i+1}(t+1) = P_i n_i(t)$$

Works for all age classes except the first. Because the number in the first age class depends on the reproduction of all age classes. So...

$$F_i = b(i)P_i$$

The fertility of age class  $i$  is the number of offspring produced, discounted by the survival probability for that age class.

# Calculating fertility for age classes ( $F_i$ )

$$n_{i+1}(t+1) = P_i n_i(t)$$

Works for all age classes except the first. Because the number in it depends on the reproduction of all age classes. So...

$$F_i = b(i)P_i$$

The fertility of age class  $i$  is the number of offspring produced, discounted by the survival probability for that age class.

$$n_1(t+1) = \sum_{i=1}^k F_i n_i(t)$$

Once the  $F_i$  is known for each age class, we multiply these fertilities by the number of individuals in each age class. This product is then summed over all age classes to calculate the number of new offspring.

# Calculating fertility for age classes ( $F_i$ )

$$n_{i+1}(t + 1) = P_i n_i(t)$$

Works for all age classes except the first. Because the number in it depends on the reproduction of all age classes. So...

$$F_i = b(i)P_i$$

The fertility of age class  $i$  is the number of offspring produced, discounted by the survival probability for that age class.

$$n_1(t + 1) = \sum_{i=1}^k F_i n_i(t)$$

Once the  $F_i$  is known for each age class, we multiply these fertilities by the number of individuals in each age class. This product is then summed over all age classes to calculate the number of new offspring.

Now... let's calculate the number of individuals in each age class for a single time step.

Now... let's calculate the number of individuals in each age class for a single time step.

$$n_1(t+1) = F_1 n_1(t) + F_2 n_2(t) + F_3 n_3(t) + F_4 n_4(t)$$



Now... let's calculate the number of individuals in each age class for a single time step.

$$n_1(t+1) = F_1n_1(t) + F_2n_2(t) + F_3n_3(t) + F_4n_4(t)$$

$$n_2(t+1) = P_1n_1(t)$$

Now... let's calculate the number of individuals in each age class for a single time step.

$$n_1(t+1) = F_1n_1(t) + F_2n_2(t) + F_3n_3(t) + F_4n_4(t)$$

$$n_2(t+1) = P_1n_1(t)$$

$$n_3(t+1) = P_2n_2(t)$$

Now... let's calculate the number of individuals in each age class for a single time step.

$$n_1(t+1) = F_1n_1(t) + F_2n_2(t) + F_3n_3(t) + F_4n_4(t)$$

$$n_2(t+1) = P_1n_1(t)$$

$$n_3(t+1) = P_2n_2(t)$$

$$n_4(t+1) = P_3n_3(t)$$

# The Leslie matrix

$$n_1(t+1) = F_1 n_1(t) + F_2 n_2(t) + F_3 n_3(t) + F_4 n_4(t)$$

$$n_2(t+1) = P_1 n_1(t)$$

$$n_3(t+1) = P_2 n_2(t)$$

$$n_4(t+1) = P_3 n_3(t)$$

# The Leslie matrix

$$n_1(t+1) = F_1n_1(t) + F_2n_2(t) + F_3n_3(t) + F_4n_4(t)$$

$$n_2(t+1) = P_1n_1(t)$$

$$n_3(t+1) = P_2n_2(t)$$

$$n_4(t+1) = P_3n_3(t)$$

A Leslie matrix describes the changes in population size due to mortality and reproduction.

# The Leslie matrix

$$n_1(t+1) = F_1n_1(t) + F_2n_2(t) + F_3n_3(t) + F_4n_4(t)$$

$$n_2(t+1) = P_1n_1(t)$$

$$n_3(t+1) = P_2n_2(t)$$

$$n_4(t+1) = P_3n_3(t)$$

A Leslie matrix describes the changes in population size due to mortality and reproduction.

If there are  $k$  age classes, the Leslie matrix is a  $k \times k$  square matrix.

$$\mathbf{A} = \begin{bmatrix} F_1 & F_2 & F_3 & F_4 \\ P_1 & 0 & 0 & 0 \\ 0 & P_2 & 0 & 0 \\ 0 & 0 & P_3 & 0 \end{bmatrix}$$

# The Leslie matrix

$$n_1(t+1) = F_1n_1(t) + F_2n_2(t) + F_3n_3(t) + F_4n_4(t)$$

$$n_2(t+1) = P_1n_1(t)$$

$$n_3(t+1) = P_2n_2(t)$$

$$n_4(t+1) = P_3n_3(t)$$

A Leslie matrix describes the changes in population size due to mortality and reproduction.

If there are  $k$  age classes, the Leslie matrix is a  $k \times k$  square matrix.

Each **column** is the age at time  $t$

Each **row** is the age at time  $t+1$

$$\mathbf{A} = \begin{bmatrix} F_1 & F_2 & F_3 & F_4 \\ P_1 & 0 & 0 & 0 \\ 0 & P_2 & 0 & 0 \\ 0 & 0 & P_3 & 0 \end{bmatrix}$$

# The Leslie matrix

$$n_1(t+1) = F_1n_1(t) + F_2n_2(t) + F_3n_3(t) + F_4n_4(t)$$

$$n_2(t+1) = P_1n_1(t)$$

$$n_3(t+1) = P_2n_2(t)$$

$$n_4(t+1) = P_3n_3(t)$$

A Leslie matrix describes the changes in population size due to mortality and reproduction.

If there are  $k$  age classes, the Leslie matrix is a  $k \times k$  square matrix.

Each **column** is the age at time  $t$

Each **row** is the age at time  $t+1$

The first row are the fertilities - contributions to newborn of each age class.

$$\mathbf{A} = \begin{bmatrix} F_1 & F_2 & F_3 & F_4 \\ P_1 & 0 & 0 & 0 \\ 0 & P_2 & 0 & 0 \\ 0 & 0 & P_3 & 0 \end{bmatrix}$$



# The Leslie matrix

$$n_1(t+1) = F_1n_1(t) + F_2n_2(t) + F_3n_3(t) + F_4n_4(t)$$

$$n_2(t+1) = P_1n_1(t)$$

$$n_3(t+1) = P_2n_2(t)$$

$$n_4(t+1) = P_3n_3(t)$$

A Leslie matrix describes the changes in population size due to mortality and reproduction.

If there are  $k$  age classes, the Leslie matrix is a  $k \times k$  square matrix.

Each **column is the age at time  $t$**

Each **row is the age at time  $t + 1$**

The first row are the fertilities - contributions to newborn of each age class.

The survival probabilities are always in the subdiagonal.

They represent transitions from one age class to the next.

$$\mathbf{A} = \begin{bmatrix} F_1 & F_2 & F_3 & F_4 \\ P_1 & 0 & 0 & 0 \\ 0 & P_2 & 0 & 0 \\ 0 & 0 & P_3 & 0 \end{bmatrix}$$

# The Leslie matrix

$$n_1(t+1) = F_1n_1(t) + F_2n_2(t) + F_3n_3(t) + F_4n_4(t)$$

$$n_2(t+1) = P_1n_1(t)$$

$$n_3(t+1) = P_2n_2(t)$$

$$n_4(t+1) = P_3n_3(t)$$

$$\mathbf{n}(t + 1) = \mathbf{A}\mathbf{n}(t)$$

A Leslie matrix describes the changes in population size due to mortality and reproduction.

If there are  $k$  age classes, the Leslie matrix is a  $k \times k$  square matrix.

Each **column is the age at time  $t$**

Each **row is the age at time  $t + 1$**

The first row are the fertilities - contributions to newborn of each age class.

The survival probabilities are always in the subdiagonal.

They represent transitions from one age class to the next.

0's: those transitions aren't possible.

$$\mathbf{A} = \begin{bmatrix} F_1 & F_2 & F_3 & F_4 \\ P_1 & 0 & 0 & 0 \\ 0 & P_2 & 0 & 0 \\ 0 & 0 & P_3 & 0 \end{bmatrix}$$

# The Leslie matrix

$x$	$i$ (age class)	$S(x)$	$b(x)$	$l(x)$	$Pi =$ $l(i)/l(i-1)$	$Fi =$ $b(i)Pi$
0		500	0	1		
1	1	400	2	0.8	0.80	1.6
2	2	200	3	0.4	0.50	1.5
3	3	50	1	0.1	0.25	0.25
4	4	0	0	0	0	0

# The Leslie matrix

$x$	$i$ (age class)	$S(x)$	$b(x)$	$l(x)$	$P_i = l(i)/l(i-1)$	$F_i = b(i)P_i$
0		500	0	1		
1	1	400	2	0.8	0.80	1.6
2	2	200	3	0.4	0.50	1.5
3	3	50	1	0.1	0.25	0.25
4	4	0	0	0	0	0

$$A = \begin{bmatrix} 1.6 & 1.5 & 0.25 & 0 \\ 0.8 & 0 & 0 & 0 \\ 0 & 0.5 & 0 & 0 \\ 0 & 0 & 0.25 & 0 \end{bmatrix}$$

# The Leslie matrix

$$\mathbf{A} = \begin{bmatrix} 1.6 & 1.5 & 0.25 & 0 \\ 0.8 & 0 & 0 & 0 \\ 0 & 0.5 & 0 & 0 \\ 0 & 0 & 0.25 & 0 \end{bmatrix}$$

# The Leslie matrix

Let's compare two populations.

$$\mathbf{A} = \begin{bmatrix} 1.6 & 1.5 & 0.25 & 0 \\ 0.8 & 0 & 0 & 0 \\ 0 & 0.5 & 0 & 0 \\ 0 & 0 & 0.25 & 0 \end{bmatrix} \quad \mathbf{n}_0 = \begin{pmatrix} 200 \\ 0 \\ 0 \\ 0 \end{pmatrix}$$

$$\mathbf{A} = \begin{bmatrix} 1.6 & 1.5 & 0.25 & 0 \\ 0.8 & 0 & 0 & 0 \\ 0 & 0.5 & 0 & 0 \\ 0 & 0 & 0.25 & 0 \end{bmatrix} \quad \mathbf{n}_0 = \begin{pmatrix} 50 \\ 50 \\ 50 \\ 50 \end{pmatrix}$$

## APPENDIX: BASIC MATRIX MANIPULATIONS

### Matrix Multiplication

A matrix is a table of numbers, and the mathematical manipulation of matrices derives from analysis of systems of linear equations. Thus, for example, if we have the system

$$Y_1 = a_1X_1 + b_1X_2 + c_1X_3, \quad (\text{A1a})$$

$$Y_2 = a_2X_1 + b_2X_2 + c_2X_3, \quad (\text{A1b})$$

and

$$Y_3 = a_3X_1 + b_3X_2 + c_3X_3, \quad (\text{A1c})$$

simply as a matter of convenience we can group the  $X_i$ s to the right and more easily visualize the structure of the system as follows:

$$\begin{vmatrix} Y_1 \\ Y_2 \\ Y_3 \end{vmatrix} = \begin{vmatrix} a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{vmatrix} \begin{vmatrix} X_1 \\ X_2 \\ X_3 \end{vmatrix}.$$

Here we have three matrices: first a matrix with a single column (the  $Y$ s), second a square matrix with three columns (sometimes referred to as the detached coefficient matrix), and third a matrix with a single column (the  $X$ s). Sometimes the various matrices are simply referred to using a single letter, but it is customary when speaking of matrices to put them in boldface type, so the above equation could be

$$\mathbf{Y} = \mathbf{AX}. \quad (\text{A2})$$



# The Leslie matrix


Let's compare two populations.

$$\mathbf{A} = \begin{bmatrix} 1.6 & 1.5 & 0.25 & 0 \\ 0.8 & 0 & 0 & 0 \\ 0 & 0.5 & 0 & 0 \\ 0 & 0 & 0.25 & 0 \end{bmatrix} \quad \mathbf{n}_0 = \begin{pmatrix} 200 \\ 0 \\ 0 \\ 0 \end{pmatrix}$$
$$\begin{aligned} 1.6 \cdot 200 + 1.5 \cdot 0 + 0.25 \cdot 0 + 0 \cdot 0 &= 320 \\ 0.8 \cdot 200 + 0 \cdot 0 + 0 \cdot 0 + 0 \cdot 0 &= 160 \\ 0 \cdot 200 + 0.5 \cdot 0 + 0 \cdot 0 + 0 \cdot 0 &= 0 \\ 0 \cdot 200 + 0 \cdot 0 + 0.25 \cdot 0 + 0 \cdot 0 &= 0 \end{aligned}$$


$$\mathbf{A} = \begin{bmatrix} 1.6 & 1.5 & 0.25 & 0 \\ 0.8 & 0 & 0 & 0 \\ 0 & 0.5 & 0 & 0 \\ 0 & 0 & 0.25 & 0 \end{bmatrix} \quad \mathbf{n}_0 = \begin{pmatrix} 50 \\ 50 \\ 50 \\ 50 \end{pmatrix}$$
$$\begin{aligned} 1.6 \cdot 50 + 1.5 \cdot 50 + 0.25 \cdot 50 + 0 \cdot 50 &= 205 \\ 0.8 \cdot 50 + 0 \cdot 50 + 0 \cdot 50 + 0 \cdot 50 &= 40 \\ 0 \cdot 50 + 0.5 \cdot 50 + 0 \cdot 50 + 0 \cdot 50 &= 25 \\ 0 \cdot 50 + 0 \cdot 50 + 0.25 \cdot 50 + 0 \cdot 50 &= 12.5 \end{aligned}$$







If a population is growing with constant birth and death rates, it will quickly converge on a stable age distribution, with relative numbers in each age class staying about the same.



# Stage- and size-structured population growth

Sometimes size or life stage is more important than age.



Remember each column represents the stage at time  $t$  and each row represents the stage at time  $t + 1$ .

	egg	larva	adult
egg	0	0	$F_{ae}$
larva	$P_{el}$	$P_{ll}$	0
adult	0	$P_{la}$	$P_{aa}$

Remember each column represents the stage at time  $t$  and each row represents the stage at time  $t + 1$ .

	egg	larva	adult
egg	0	0	$F_{ae}$
larva	$P_{el}$	$P_{ll}$	0
adult	0	$P_{la}$	$P_{aa}$

The first row represents the fertilities.

Remember each column represents the stage at time  $t$  and each row represents the stage at time  $t + 1$ .

	egg	larva	adult
egg	0	0	$F_{ae}$
larva	$P_{el}$	$P_{ll}$	0
adult	0	$P_{la}$	$P_{aa}$

The first row represents the fertilities.

The other entries are *transition probabilities*.


	size 1	size 2	size 3	size 4	size 5
size 1	$P_{11}$	$F_{21}$	$F_{31}$	$F_{41}$	$F_{51}$
size 2	$P_{12}$	$P_{22}$	0	0	0
size 3	0	$P_{23}$	$P_{33}$	0	0
size 4	0	0	$P_{34}$	$P_{44}$	0
size 5	0	0	0	$P_{45}$	$P_{55}$

	egg	larva	adult
egg	0	0	$F_{ae}$
larva	$P_{el}$	$P_{ll}$	0
adult	0	$P_{la}$	$P_{aa}$







	egg	larva	adult
egg	0	0	$F_{ae}$
larva	$P_{el}$	$P_{ll}$	0
adult	0	$P_{la}$	$P_{aa}$



	egg	larva	adult
egg	0	0	$F_{ae}$
larva	$P_{el}$	$P_{ll}$	0
adult	0	$P_{la}$	$P_{aa}$



	egg	larva	adult
egg	0	0	$F_{ae}$
larva	$P_{el}$	$P_{ll}$	0
adult	0	$P_{la}$	$P_{aa}$



	egg	larva	adult
egg	0	0	$F_{ae}$
larva	$P_{el}$	$P_{ll}$	0
adult	0	$P_{la}$	$P_{aa}$

