ALICE MUON Software for run 3

Sean Murray
Physics
University of Cape Town

July 6, 2016

Outline

ALICE

(Illustrating BEAMER's \pause command.)

A couple of years ago, a fifth-grade teacher asked me to explain to her the reasoning behind the "invert and multiply" rule for dividing fractions, e.g.

ALICE

(Illustrating BEAMER's \pause command.)

A couple of years ago, a fifth-grade teacher asked me to explain to her the reasoning behind the "invert and multiply" rule for dividing fractions, e.g.

$$\frac{4}{5} \div \frac{2}{3} = \frac{4}{5} \times \frac{3}{2}$$

ALICE

(Illustrating BEAMER's \pause command.)

A couple of years ago, a fifth-grade teacher asked me to explain to her the reasoning behind the "invert and multiply" rule for dividing fractions, e.g.

$$\frac{4}{5} \div \frac{2}{3} = \frac{4}{5} \times \frac{3}{2}$$

Let's try to find answers understandable by fifth graders (at least the more patient ones).

ALICE MUON Arm

Here let's just use intuition, understandable by fifth graders.

ALICE MUON Arm

Here let's just use intuition, understandable by fifth graders.
If we give 1/3 of a cookie to each person, how many people can we feed with 1 cookie?

ALICE MUON Arm

Here let's just use intuition, understandable by fifth graders.
If we give 1/3 of a cookie to each person, how many people can we feed with 1 cookie?
Obviously, the answer is 3.
So we've derived the "invert and multiply" rule in a special case:

$$1 \div \frac{1}{3} = 3$$

Detector Structure

But what if we give 2/3 of a cookie, not 1/3, to each person?
We're giving $2\times$ as much per person.
So we can feed only 1/2 as many people.
So we feed $\frac{1}{2} \times 3 = \frac{3}{2}$.[1]
So we've derived the "invert and multiply" rule in another case:

$$1 \div \frac{2}{3} = \frac{3}{2}$$

[1] One person gets only a half share.

Detector Structure 2

Now, suppose we have only 4/5 of a cookie.
Then we can feed only 4/5 as many people, i.e.

$$\frac{4}{5} \times \frac{3}{2} \textit{ people}$$

Detector Structure 2

Now, suppose we have only 4/5 of a cookie.
Then we can feed only 4/5 as many people, i.e.

$$\frac{4}{5} \times \frac{3}{2} \textit{ people}$$

So we've derived the "invert and multiply" rule in the general case:

$$\frac{4}{5} \div \frac{2}{3} = \frac{4}{5} \times \frac{3}{2}$$

Outline

Current MUON Software

(Illustrating BEAMER's \uncover command.)

Theorem

The angles in a triangle sum to 180°*.*

Current MUON Software

(Illustrating BEAMER's \uncover command.)

Theorem

The angles in a triangle sum to 180°*.*

Plan: Extend AC past C to D. Draw CE parallel to AB.



Proof.

1. u = y

□

Proof.

1. u = y Alternate angles of a transveral.

□

Proof.

1. u = y — Alternate angles of a transveral.
2. v = x

□

Proof.

1. u = y	Alternate angles of a transveral.
2. v = x	Consecutive interior angles of a transveral

□

Proof.

1. $u = y$ — Alternate angles of a transveral.
2. $v = x$ — Consecutive interior angles of a transveral
3. $z+u+v = 180°$

□

Proof.

1. $u = y$	Alternate angles of a transveral.
2. $v = x$	Consecutive interior angles of a transveral
3. $z+u+v = 180°$	ACD is a straight line.

□

Proof.

1. $u = y$	Alternate angles of a transveral.
2. $v = x$	Consecutive interior angles of a transveral
3. $z+u+v = 180°$	ACD is a straight line.
4. $z+y+x = 180°$	

□

Proof.

1. $u = y$	Alternate angles of a transveral.
2. $v = x$	Consecutive interior angles of a transveral
3. $z+u+v = 180°$	ACD is a straight line.
4. $z+y+x = 180°$	Substitution from Steps 1 and 2.

□

Outline

What is a cluster finder

- This tour just scratches the surface.

What is a cluster finder

- This tour just scratches the surface.
- BEAMER has enough features to fill a 210-page user manual!

What is a cluster finder

- This tour just scratches the surface.
- BEAMER has enough features to fill a 210-page user manual!
- Advanced example: `http://latex-beamer.sourceforge.net/beamerexample1.pdf`.