


Recall: If an $n \times n$ matrix A is invertible then:

- the equation $A\mathbf{x} = \mathbf{b}$ has a unique solution for each $\mathbf{b} \in \mathbb{R}^n$
- the linear transformation $T_A: \mathbb{R}^n \rightarrow \mathbb{R}^n$, $T_A(\mathbf{v}) = A\mathbf{v}$ has an inverse function.

Determinants recognize which matrices are invertible:



Example: Determinant for a 1×1 matrix.

$$A = \begin{bmatrix} a \end{bmatrix}$$

Example: Determinant for a 2×2 matrix.

$$A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

Definition

If A is an $n \times n$ matrix then for $1 \leq i, j \leq n$ the (i, j) -minor of A is the matrix A_{ij} obtained by deleting the i^{th} row and j^{th} column of A .

Example.

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix}$$

Definition

Let A be an $n \times n$ matrix

$$A = \begin{bmatrix} a_{11} & \dots & a_{1n} \\ \vdots & & \vdots \\ a_{n1} & \dots & a_{nn} \end{bmatrix}$$

1) If $n = 1$, i.e. $A = [a_{11}]$, then $\det A = a_{11}$

2) If $n > 1$ then

$$\begin{aligned} \det A = & (-1)^{1+1} a_{11} \cdot \det A_{11} \\ & + (-1)^{1+2} a_{12} \cdot \det A_{12} \\ & \dots \quad \dots \quad \dots \quad \dots \\ & + (-1)^{1+n} a_{1n} \cdot \det A_{1n} \end{aligned}$$

Example. ($n = 2$)

$$A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$$

Note

If A is a 2×2 matrix

$$A = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix}$$

then $\det A = a_{11} \cdot a_{22} - a_{12} \cdot a_{21}$

Example. (n=3)

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix}$$

Example (n=4)

$$A = \begin{bmatrix} 1 & 0 & 2 & 0 \\ 0 & 4 & 0 & 1 \\ 2 & 1 & 6 & 1 \\ 3 & 5 & 7 & 0 \end{bmatrix}$$

Note. In order to compute the determinant of an $n \times n$ matrix in this way we need to compute:

| | |
|-----------------------------------|---|
| n | determinants of $(n-1) \times (n-1)$ matrices |
| $n(n-1)$ | determinants of $(n-2) \times (n-2)$ matrices |
| $n(n-1)(n-2)$ | determinants of $(n-3) \times (n-3)$ matrices |
| \dots | \dots |
| $n(n-1)(n-2) \cdot \dots \cdot 3$ | determinants of 2×2 matrices |

E.g. for a 25×25 matrix we would need to compute

$$25 \cdot 24 \cdot 23 \cdot \dots \cdot 3 = 7,755,605,021,665,492,992,000,000$$

determinants of 2×2 matrices.

Next: How to compute determinants faster.