

Recall:

If  $A = [v_1 \ \dots \ v_n]$  is an  $m \times n$  matrix then:

- 1)  $\text{Col}(A) = \text{Span}(v_1, \dots, v_n)$
- 2)  $\text{Nul}(A) = \{v \in \mathbb{R}^m \mid Av = 0\}$

$\text{Col}(A)$  is a subspace of  $\mathbb{R}^m$ .

e.g.  $A = \begin{bmatrix} 1 & 3 & 5 \\ 2 & 4 & 6 \end{bmatrix} \Big\}^{m=2} \quad \text{Col}(A) = \text{Span}\left(\begin{bmatrix} 1 \\ 2 \end{bmatrix}, \begin{bmatrix} 3 \\ 4 \end{bmatrix}, \begin{bmatrix} 5 \\ 6 \end{bmatrix}\right) \subseteq \mathbb{R}^2$

$\text{Nul}(A)$  is a subspace of  $\mathbb{R}^n$ .

e.g.  $\text{Nul}\left(\begin{bmatrix} 1 & 3 & 5 \\ 2 & 4 & 6 \end{bmatrix}\right) = \left\{ \begin{bmatrix} c_1 \\ c_2 \\ c_3 \end{bmatrix} \mid \begin{bmatrix} 1 & 3 & 5 \\ 2 & 4 & 6 \end{bmatrix} \cdot \begin{bmatrix} c_1 \\ c_2 \\ c_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \right\} \subseteq \mathbb{R}^3$

## Construction of a basis of $\text{Col}(A)$

Example:

$$A = \begin{bmatrix} 1 & 3 & 5 \\ 2 & 4 & 6 \end{bmatrix} \quad \text{Col}(A) = \text{Span}\left(\begin{bmatrix} 1 \\ 2 \end{bmatrix}, \begin{bmatrix} 3 \\ 4 \end{bmatrix}, \begin{bmatrix} 5 \\ 6 \end{bmatrix}\right) \subseteq \mathbb{R}^2.$$

### Lemma

Let  $V$  be a vector space, and let  $v_1, \dots, v_p \in V$ . If a vector  $v_i$  is a linear combination of the other vectors then

$$\text{Span}(v_1, \dots, v_p) = \text{Span}(v_1, \dots, v_{i-1}, v_{i+1}, \dots, v_p)$$

**Upshot.** One can construct a basis of a vector space  $V$  as follows:

- Start with a set of vectors  $\{v_1, \dots, v_p\}$  such that  $\text{Span}(v_1, \dots, v_p) = V$ .
- Keep removing vectors without changing the span, until you get a linearly independent set.

**Example.** Find a basis of  $\text{Col}(A)$  where  $A$  is the following matrix:

$$A = \begin{matrix} & \begin{matrix} v_1 & v_2 & v_3 & v_4 & v_5 & v_6 \end{matrix} \\ \begin{matrix} \textcircled{1} & 0 & 2 & 0 & 1 & 0 \\ 0 & \textcircled{1} & 3 & 0 & -1 & 0 \\ 0 & 0 & 0 & \textcircled{1} & 3 & 0 \\ 0 & 0 & 0 & 0 & 0 & \textcircled{1} \\ 0 & 0 & 0 & 0 & 0 & 0 \end{matrix} \end{matrix}$$

matrix in the reduced  
row echelon form.

Solution

$$\text{Col}(A) = \text{Span}(v_1, v_2, \cancel{v_3}, v_4, \cancel{v_5}, v_6) \subseteq \mathbb{R}^5$$

Note:  $v_3 = 2v_1 + 3v_2$

$$v_5 = v_1 - v_2 + 3v_4$$

This gives:

$$\text{Col}(A) = \text{Span}(v_1, v_2, v_4, v_6)$$

Note: The set  $\{v_1, v_2, v_4, v_6\}$  is linearly independent,  
so it is a basis of  $\text{Col}(A)$ .

In general: If  $A$  is a matrix in the reduced echelon form then the set of all columns of  $A$  which contain leading ones is a basis of  $\text{Col}(A)$ .

**Example.** Find a basis of  $\text{Col}(A)$  where  $A$  is the following matrix:

$$A = \begin{matrix} & \begin{matrix} v_1 & v_2 & v_3 & v_4 & v_5 \end{matrix} \\ \begin{bmatrix} -3 & 6 & -1 & 1 & -7 \\ 1 & -2 & 2 & 3 & -1 \\ 2 & -4 & 5 & 8 & -4 \end{bmatrix} \end{matrix}$$

Solution:

$$\text{Col}(A) = \text{Span}(v_1, v_2, v_3, v_4, v_5)$$

$$\begin{matrix} \begin{matrix} v_1 & v_2 & v_3 & v_4 & v_5 \\ \begin{bmatrix} -3 & 6 & -1 & 1 & -7 \\ 1 & -2 & 2 & 3 & -1 \\ 2 & -4 & 5 & 8 & -4 \end{bmatrix} \end{matrix} & \xrightarrow{\text{row red}} & \begin{matrix} w_1 & w_2 & w_3 & w_4 & w_5 \\ \begin{bmatrix} 1 & -2 & 0 & -1 & 3 \\ 0 & 0 & 1 & 2 & -2 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix} \end{matrix} \end{matrix}$$

$$\begin{cases} v_1 = -2v_2 \\ v_4 = -v_1 + 2v_3 \\ v_5 = 3v_1 - 2v_3 \end{cases}$$

$$\begin{cases} w_2 = -2w_1 \\ w_4 = -w_1 + 2w_3 \\ w_5 = 3w_1 - 2w_3 \end{cases}$$

This gives:  $\text{Col}(A) = \text{Span}(v_1, v_3)$ .

Check: the set  $\{v_1, v_3\}$  is linearly independent, so it is a basis of  $\text{Col}(A)$ .

In general: If  $A$  is a matrix then the set of pivot columns of  $A$  is a basis of  $\text{Col}(A)$ .