

Example.

$$A = \begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix} \text{ is invertible, } A^{-1} = \begin{bmatrix} \frac{1}{2} & \frac{1}{2} \\ -\frac{1}{2} & \frac{1}{2} \end{bmatrix}$$

Matrix inverses and matrix equations



Proposition

If A is an invertible matrix then for any vector \mathbf{b} the equation $A\mathbf{x} = \mathbf{b}$ has exactly one solution.

Example. Solve the following matrix equation:



$$\begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix} \cdot \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$$

Matrix inverses and matrix transformations





Example.

$$A = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$$



Example.

$$A = \begin{bmatrix} 1 & 2 \\ 2 & 4 \end{bmatrix}$$



Upshot. If an $m \times n$ matrix A is invertible then the matrix transformation $T_A: \mathbb{R}^n \rightarrow \mathbb{R}^m$ must be one-to-one and onto.

Recall: If A be is $m \times n$ matrix then the matrix transformation $T_A: \mathbb{R}^n \rightarrow \mathbb{R}^m$ is:

- onto if and only if A has a pivot position in every row
- one-to-one if and only if A has a pivot position in every column.

Theorem

If A is not a square matrix then it is not invertible.

If A is a square matrix then the following conditions are equivalent:

- 1) A is an invertible matrix.
- 2) The matrix A has a pivot position in every row and column.
- 3) The reduced row echelon form of A is the identity matrix I_n .

Proposition

If A is an $n \times n$ invertible matrix then

$$A^{-1} = [\mathbf{w}_1 \quad \mathbf{w}_2 \quad \dots \quad \mathbf{w}_n]$$

where \mathbf{w}_i is the solution of $A\mathbf{x} = \mathbf{e}_i$.


Example.

$$A = \begin{bmatrix} 1 & -1 \\ -1 & 1 \end{bmatrix}$$

Simplification:
How to solve several matrix equations with the same
coefficient matrix at the same time

$$Ax = \mathbf{b}_1, Ax = \mathbf{b}_2, \dots, Ax = \mathbf{b}_n$$

matrix of equations



$$\left[A \mid \mathbf{b}_1 \ \mathbf{b}_2 \ \dots \ \mathbf{b}_n \right]$$

augmented matrix



$$\left[\begin{array}{c|c} & \end{array} \right]$$

reduced matrix



solutions

Example. Solve the vector equations $A\mathbf{x} = \mathbf{e}_1$ and $A\mathbf{x} = \mathbf{e}_2$ where

$$A = \begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix} \quad \mathbf{e}_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \quad \mathbf{e}_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

Summary:
How to invert a matrix

Example: $A = \begin{bmatrix} 1 & 1 \\ 1 & 2 \end{bmatrix}$

1) Augment A by the identity matrix.

2) Reduce the augmented matrix.

2) If after the row reduction the matrix on the left is the identity matrix, then A is invertible and

$$A^{-1} = \text{the matrix on the right}$$

Otherwise A is not invertible.

Properties of matrix inverses

1) If A is invertible then A^{-1} is invertible and

$$(A^{-1})^{-1} = A$$

2) If A, B are invertible then AB is invertible and

$$(AB)^{-1} = B^{-1}A^{-1}$$

3) If A is invertible then A^T is invertible and

$$(A^T)^{-1} = (A^{-1})^T$$

Ciphers.

Cipher is an algorithm for encrypting and decrypting data to conceal its meaning.

Basic working scheme of ciphers

