

Recall: An $n \times n$ matrix A defines a linear transformation



$$T_A: \mathbb{R}^n \rightarrow \mathbb{R}^n$$

given by $T_A(\mathbf{v}) = A\mathbf{v}$.

Next goal: Understand this linear transformation better.



Example.

$$A = \begin{bmatrix} 2 & 0 \\ 0 & 3 \end{bmatrix}$$



Example.

$$A = \begin{bmatrix} 2 & 1 \\ 1 & 2 \end{bmatrix}$$



Definition

Let A be an $n \times n$ matrix. If $\mathbf{v} \in \mathbb{R}^n$ is a non-zero vector and λ is a scalar such that

$$A\mathbf{v} = \lambda\mathbf{v}$$

then we say that

- λ is an *eigenvalue* of A
- \mathbf{v} is an *eigenvector* of A corresponding to λ .

Example.

$$A = \begin{bmatrix} 2 & 0 \\ 0 & 3 \end{bmatrix}$$

Example.

$$A = \begin{bmatrix} 2 & 1 \\ 1 & 2 \end{bmatrix}$$

Computation of eigenvalues

Recall: $I_n = n \times n$ identity matrix:

$$I_n = \begin{bmatrix} 1 & 0 & \dots & 0 \\ 0 & 1 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 1 \end{bmatrix}$$

Proposition

If A be an $n \times n$ matrix then $\lambda \in \mathbb{R}$ is an eigenvalue of A if and only if the matrix equation

$$(A - \lambda I_n)\mathbf{x} = \mathbf{0}$$

has a non-trivial solution.

Proposition

If B is an $n \times n$ matrix then equation

$$Bx = 0$$

has a non-trivial solution if and only if the matrix B is not invertible.

Proposition

If A be an $n \times n$ matrix then $\lambda \in \mathbb{R}$ is an eigenvalue of A if and only if

$$\det(A - \lambda I_n) = 0$$

Example. Find all eigenvalues of the following matrix:

$$A = \begin{bmatrix} 2 & 2 & 1 \\ 1 & 3 & 1 \\ 1 & 2 & 2 \end{bmatrix}$$

Definition

If A is an $n \times n$ matrix then

$$P(\lambda) = \det(A - \lambda I_n)$$

is a polynomial of degree n . $P(\lambda)$ is called the *characteristic polynomial* of the matrix A .

Upshot

If A is a square matrix then

$$\text{eigenvalues of } A = \text{roots of } P(\lambda)$$

Example.

$$A = \begin{bmatrix} 2 & 2 & 1 \\ 1 & 3 & 1 \\ 1 & 2 & 2 \end{bmatrix}$$

Corollary

An $n \times n$ matrix can have at most n distinct eigenvalues.

Computation of eigenvectors

Proposition

If λ is an eigenvalue of an $n \times n$ matrix A then

$$\left\{ \begin{array}{l} \text{eigenvectors of } A \\ \text{corresponding to } \lambda \end{array} \right\} = \left\{ \begin{array}{l} \text{vectors in} \\ \text{Nul}(A - \lambda I_n) \end{array} \right\}$$

Corollary/Definition

If A is an $n \times n$ matrix and λ is an eigenvalue of A then the set of all eigenvectors corresponding to λ is a subspace of \mathbb{R}^n .

This subspace is called the *eigenspace* of A corresponding to λ .

Proposition

If λ is an eigenvalue of an $n \times n$ matrix A then

$$\left\{ \begin{array}{l} \text{eigenspace of } A \\ \text{corresponding to } \lambda \end{array} \right\} = \text{Nul}(A - \lambda I_n)$$

Example. Consider the following matrix:

$$A = \begin{bmatrix} 2 & 2 & 1 \\ 1 & 3 & 1 \\ 1 & 2 & 2 \end{bmatrix}$$

Recall that eigenvalues of A are $\lambda_1 = 1$ and $\lambda_2 = 5$. Compute bases of eigenspaces of A corresponding to these eigenvalues.

Solution.

$$\underline{\lambda_1 = 1}$$

$$\underline{\lambda_2 = 5}$$