

Definition

If

$$\mathbf{u} = \begin{bmatrix} a_1 \\ \vdots \\ a_n \end{bmatrix} \quad \mathbf{v} = \begin{bmatrix} b_1 \\ \vdots \\ b_n \end{bmatrix}$$

are vectors in \mathbb{R}^n then the *inner product* (or *dot product*) of \mathbf{u} and \mathbf{v} is the number

$$\mathbf{u} \cdot \mathbf{v} = a_1 b_1 + \dots + a_n b_n$$

Properties of the dot product:

- 1) $\mathbf{u} \cdot \mathbf{v} = \mathbf{v} \cdot \mathbf{u}$
- 2) $(\mathbf{u} + \mathbf{v}) \cdot \mathbf{w} = \mathbf{u} \cdot \mathbf{w} + \mathbf{v} \cdot \mathbf{w}$
- 3) $(c\mathbf{u}) \cdot \mathbf{v} = c(\mathbf{u} \cdot \mathbf{v})$
- 4) $\mathbf{u} \cdot \mathbf{u} \geq 0$ and $\mathbf{u} \cdot \mathbf{u} = 0$ if and only if $\mathbf{u} = \mathbf{0}$.

Definition

If $\mathbf{u} \in \mathbb{R}^n$ then the *length* (or the *norm*) of \mathbf{u} is the number

$$\|\mathbf{u}\| = \sqrt{\mathbf{u} \cdot \mathbf{u}}$$

Note. If $\mathbf{u} = \begin{bmatrix} a_1 \\ \vdots \\ a_n \end{bmatrix}$ then $\|\mathbf{u}\| = \sqrt{a_1^2 + \dots + a_n^2}$.

Properties of the norm:

- 1) $\|\mathbf{u}\| \geq 0$ and $\|\mathbf{u}\| = 0$ if and only if $\mathbf{u} = \mathbf{0}$.
- 2) $\|c\mathbf{u}\| = |c| \cdot \|\mathbf{u}\|$


Definition

A vector $\mathbf{u} \in \mathbb{R}^n$ is an *unit vector* if $\|\mathbf{u}\| = 1$.

Definition

If $\mathbf{u}, \mathbf{v} \in \mathbb{R}^n$ then the *distance* between \mathbf{u} and \mathbf{v} is the number

$$\text{dist}(\mathbf{u}, \mathbf{v}) = \|\mathbf{u} - \mathbf{v}\|$$



Note. If $\mathbf{u} = \begin{bmatrix} a_1 \\ \vdots \\ a_n \end{bmatrix}$, $\mathbf{v} = \begin{bmatrix} b_1 \\ \vdots \\ b_n \end{bmatrix}$ then

$$\text{dist}(\mathbf{u}, \mathbf{v}) = \sqrt{(a_1 - b_1)^2 + \dots + (a_n - b_n)^2}$$

Definition

Vectors $\mathbf{u}, \mathbf{v} \in \mathbb{R}^n$ are *orthogonal* if $\mathbf{u} \cdot \mathbf{v} = 0$.

Pythagorean Theorem

Vectors \mathbf{u}, \mathbf{v} are orthogonal if and only if

$$\|\mathbf{u}\|^2 + \|\mathbf{v}\|^2 = \|\mathbf{u} + \mathbf{v}\|^2$$