


Recall:

1)  $\text{Span}(\mathbf{v}_1, \dots, \mathbf{v}_p) = \left\{ \begin{array}{l} \text{the set of all} \\ \text{linear combinations} \\ c_1\mathbf{v}_1 + \dots + c_p\mathbf{v}_p \end{array} \right\}$


2) A set of vectors  $\{\mathbf{v}_1, \dots, \mathbf{v}_p\}$  is linearly independent if the equation

$$x_1\mathbf{v}_1 + \dots + x_p\mathbf{v}_p = \mathbf{0}$$


has only one, trivial solution  $x_1 = 0, \dots, x_p = 0$ .




## Linear independence vs. Span




$\{u\}$  linearly independent



$\{u\}$  linearly dependent



$\{u, v\}$  linearly independent



$\{u, v\}$  linearly dependent

### Theorem

If  $\{v_1, \dots, v_p\}$  is a linearly dependent set of vectors in then for some  $v_i$  we have

$$\text{Span}(v_1, \dots, v_p) = \text{Span}(v_1, \dots, v_{i-1}, v_{i+1}, \dots, v_p)$$

**Example.**

$$v_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \quad v_2 = \begin{bmatrix} 2 \\ 0 \end{bmatrix} \quad v_3 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$