



MTH 309T LINEAR ALGEBRA

EXAM 1

October 3, 2019

Name:

Yanyan Li

UB Person Number:

5	0	2	7	5	1	3	0
0	0	0	0	0	0	0	0
1	1	1	1	1	1	1	1
2	2	2	2	2	2	2	2
3	3	3	3	3	3	3	3
4	4	4	4	4	4	4	4
5	5	5	5	5	5	5	5
6	6	6	6	6	6	6	6
7	7	7	7	7	7	7	7
8	8	8	8	8	8	8	8
9	9	9	9	9	9	9	9

Instructions:

- Textbooks, calculators and any other electronic devices are not permitted. You may use one sheet of notes.
- For full credit solve each problem fully, showing all relevant work.

1	2	3	4	5	6	7	TOTAL	GRADE

--

--

--

--

--

--

--

0

nan

1

2

3

4

5

6

7

TOTAL

GRADE



1. (20 points) Consider the following vectors in \mathbb{R}^3 :

$$v_1 = \begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix}, \quad v_2 = \begin{bmatrix} -1 \\ 1 \\ -3 \end{bmatrix}, \quad v_3 = \begin{bmatrix} 1 \\ 2 \\ 0 \end{bmatrix}, \quad w = \begin{bmatrix} -2 \\ 2 \\ b \end{bmatrix}$$

a) Find all values of b such that $w \in \text{Span}(v_1, v_2, v_3)$.

b) Is the set $\{v_1, v_2, v_3\}$ linearly independent? Justify your answer.

$$a) \left[\begin{array}{ccc|c} 1 & -1 & 1 & -2 \\ 0 & 1 & 2 & 2 \\ 2 & -3 & 0 & b \end{array} \right] \xrightarrow{\text{row reduce}} \left[\begin{array}{ccc|c} 1 & 0 & 3 & 0 \\ 0 & 1 & 2 & 2 \\ 0 & 0 & 0 & 2 + \frac{b}{3} \end{array} \right]$$

$$\therefore w \in \text{Span}(v_1, v_2, v_3)$$

$$\therefore 2 + \frac{b}{3} = 0$$

$$\therefore b = -6$$

$$b) \left[\begin{array}{ccc|c} 1 & -1 & 1 & -2 \\ 0 & 1 & 2 & 2 \\ 2 & -3 & 0 & b \end{array} \right] \xrightarrow{\text{row reduce}} \left[\begin{array}{ccc|c} 1 & 0 & 3 & 0 \\ 0 & 1 & 2 & 2 \\ 2 & -3 & 0 & b \end{array} \right] \rightarrow \left[\begin{array}{ccc|c} 1 & 0 & 3 & 0 \\ 0 & 1 & 2 & 2 \\ 0 & 1 & 2 & b \end{array} \right] \rightarrow \left[\begin{array}{ccc|c} 1 & 0 & 3 & 0 \\ 0 & 1 & 2 & 2 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

there is no pivot position in last column.

Thus, the set is linearly dependent.



2. (10 points) Consider the following matrix:

$$A = \begin{bmatrix} 1 & -1 & 2 \\ 1 & 0 & 1 \\ 0 & 2 & -1 \end{bmatrix}$$

Compute A^{-1} .

$$AA^{-1} = I$$

$$\left[\begin{array}{ccc|ccc} 1 & -1 & 2 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 2 & -1 & 0 & 0 & 1 \end{array} \right] \longrightarrow \left[\begin{array}{ccc|ccc} 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & -1 & 1 & 1 & -1 & 0 \\ 0 & 2 & -1 & 0 & 0 & 1 \end{array} \right] \xrightarrow{x_2} \longrightarrow$$

$$\left[\begin{array}{ccc|ccc} 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 0 & 1 & 2 & -2 & 1 \\ 0 & 2 & -1 & 0 & 0 & 1 \end{array} \right] \longrightarrow \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & -2 & 3 & -1 \\ 0 & 0 & 1 & 2 & -2 & 1 \\ 0 & 2 & -1 & 0 & 0 & 1 \end{array} \right] \xrightarrow{x_2} \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & -2 & 3 & -1 \\ 0 & 2 & 0 & 2 & -2 & 2 \\ 0 & 0 & 1 & -2 & -2 & 1 \end{array} \right]$$

$$\longrightarrow \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & -2 & 3 & -1 \\ 0 & 1 & 0 & 1 & -1 & 1 \\ 0 & 0 & 1 & -2 & -2 & 1 \end{array} \right]$$

$$A^{-1} = \begin{bmatrix} -2 & 3 & -1 \\ 1 & -1 & 1 \\ 2 & -2 & 1 \end{bmatrix}$$



3. (10 points) Let A be the same matrix as in Problem 2, and let

$$B = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 4 \\ 3 & 2 & 1 \end{bmatrix}$$

Find a matrix C such that $A^T C = B$ (where A^T is the transpose of A).

$$A = \begin{bmatrix} 1 & -1 & 2 \\ 1 & 0 & 1 \\ 0 & 2 & -1 \end{bmatrix} \Rightarrow A^T = \begin{bmatrix} 1 & 1 & 0 \\ -1 & 0 & 2 \\ 2 & 1 & -1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 1 & 0 \\ -1 & 0 & 2 \\ 2 & 1 & -1 \end{bmatrix} C = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 4 \\ 3 & 2 & 1 \end{bmatrix}$$

$$\left[\begin{array}{ccc|ccc} 1 & 1 & 0 & 1 & 2 & 3 \\ -1 & 0 & 2 & 4 & 5 & 4 \\ 2 & 1 & -1 & 3 & 2 & 1 \end{array} \right] \xrightarrow{\text{rref}} \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & 8 & 5 & 0 \\ 0 & 1 & 0 & -7 & -3 & 3 \\ 0 & 0 & 1 & 6 & 5 & 2 \end{array} \right]$$

$$C = \begin{bmatrix} 8 & 5 & 0 \\ -7 & -3 & 3 \\ 6 & 5 & 2 \end{bmatrix}$$



4. (20 points) Let $T: \mathbb{R}^2 \rightarrow \mathbb{R}^3$ be a linear transformation given by

$$T\left(\begin{bmatrix} x_1 \\ x_2 \end{bmatrix}\right) = \begin{bmatrix} x_1 - 2x_2 \\ x_1 + x_2 \\ x_1 - 3x_2 \end{bmatrix}$$

a) Find the standard matrix of T .

b) Find all vectors u satisfying $T(u) = \begin{bmatrix} 1 \\ 10 \\ -2 \end{bmatrix}$.

$$a) T(e_1) = T\left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}\right) = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$$

$$T(e_2) = T\left(\begin{bmatrix} 0 \\ 1 \end{bmatrix}\right) = \begin{bmatrix} -2 \\ 1 \\ -3 \end{bmatrix}$$

$$\text{the standard matrix} = \begin{bmatrix} 1 & -2 \\ 1 & 1 \\ 1 & -3 \end{bmatrix}$$

$$b) \begin{bmatrix} 1 & -2 \\ 1 & 1 \\ 1 & -3 \end{bmatrix} u = \begin{bmatrix} 1 \\ 10 \\ -2 \end{bmatrix}$$

$$\left[\begin{array}{cc|c} 1 & -2 & 1 \\ 1 & 1 & 10 \\ 1 & -3 & -2 \end{array} \right] \xrightarrow{\text{ref}} \left[\begin{array}{cc|c} 1 & 0 & 7 \\ 0 & 1 & 3 \\ 0 & 0 & 0 \end{array} \right] \Rightarrow \begin{matrix} x_1 = 7 \\ x_2 = 3 \end{matrix} \Rightarrow v = \begin{bmatrix} 7 \\ 3 \end{bmatrix}$$



5. (20 points) For each matrix A given below determine if the matrix transformation $T_A: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ given by $T_A(v) = Av$ is one-to-one or not. If T_A is not one-to-one, find two vectors v_1 and v_2 such that $T_A(v_1) = T_A(v_2)$.

a) $A = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 3 & 4 & 4 \end{bmatrix}$

b) $A = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 3 & 4 & 2 \end{bmatrix}$

a) $\left[\begin{array}{ccc|c} 1 & 1 & 0 & 0 \\ 0 & 2 & 4 & 0 \\ 3 & 4 & 4 & 0 \end{array} \right] \rightarrow \left[\begin{array}{ccc|c} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{array} \right] \quad \text{Null}(A) = \{0\}$

every column has pivot position

Thus, It's one-to-one

b) $\left[\begin{array}{ccc|c} 1 & 1 & 0 & 0 \\ 0 & 2 & 4 & 0 \\ 3 & 4 & 2 & 0 \end{array} \right] \rightarrow \left[\begin{array}{ccc|c} 1 & 0 & -2 & 0 \\ 0 & 1 & 2 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$

the third column has no pivot position,

Thus, It's not one-to-one

$v_1 = \begin{bmatrix} 2 \\ -2 \\ 1 \end{bmatrix} \quad v_2 = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$

$T_A(v_1) = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 3 & 4 & 2 \end{bmatrix} \begin{bmatrix} 2 \\ -2 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$

$T_A(v_2) = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 3 & 4 & 2 \end{bmatrix} \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$



6. (10 points) For each of the statements given below decide if it is true or false. If it is true explain why. If it is false give a counterexample.

a) If u, v, w are vectors in \mathbb{R}^3 such that $w + u \in \text{Span}(u, v)$ then $w \in \text{Span}(u, v)$.

True

$$w + u = c_1 u + c_2 v \quad (c_1, c_2 \in \mathbb{R})$$

$$w = (c_1 - 1)u + c_2 v$$

$$w \in \text{Span}(u, v)$$

b) If u, v, w are vectors in \mathbb{R}^3 such that the set $\{u, v, w\}$ is linearly independent then the set $\{u, v\}$ must be linearly independent.

True.

$\{u, v, w\}$ is linearly independent means they're not multiple of each other.

then $\{u, v\}$: u, v won't be multiple of each other

then they're linearly independent



7. (10 points) For each of the statements given below decide if it is true or false. If it is true explain why. If it is false give a counterexample.

a) If A is a 2×2 matrix and u, v are vectors in \mathbb{R}^2 such that Au, Av are linearly dependent then u, v also must be linearly dependent.

True

b) If $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is a linear transformation and $u, v, w \in \mathbb{R}^2$ are vectors such that u is in $\text{Span}(v, w)$ then $T(u)$ must be in $\text{Span}(T(v), T(w))$.

True

$$u = c_1 v + c_2 w$$

$$T(u) = T(c_1 v + c_2 w)$$

$\because T$ is Linear transformation

$$\therefore T(c_1 v) + T(c_2 w) = T(c_1 v + c_2 w)$$

$$\therefore T(u) = T(c_1 v) + T(c_2 w)$$

$$\therefore T(u) \in \text{Span}(T(v), T(w))$$