



MTH 309T LINEAR ALGEBRA

EXAM 1

October 3, 2019

Name:

Michael Leishear

UB Person Number:

5	0	2	6	4	1	5	8
0	0	0	0	0	0	0	0
1	1	1	1	1	1	1	1
2	2	2	2	2	2	2	2
3	3	3	3	3	3	3	3
4	4	4	4	4	4	4	4
5	5	5	5	5	5	5	5
6	6	6	6	6	6	6	6
7	7	7	7	7	7	7	7
8	8	8	8	8	8	8	8
9	9	9	9	9	9	9	9

Instructions:

- Textbooks, calculators and any other electronic devices are not permitted. You may use one sheet of notes.
- For full credit solve each problem fully, showing all relevant work.

1	2	3	4	5	6	7	TOTAL	GRADE

8

6

5

15

12

2

2

--

--

50

D+

1

2

3

4

5

6

7

PIAZZA

HILL

TOTAL

GRADE



1. (20 points) Consider the following vectors in \mathbb{R}^3 :

$$v_1 = \begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix}, \quad v_2 = \begin{bmatrix} -1 \\ 1 \\ -3 \end{bmatrix}, \quad v_3 = \begin{bmatrix} 1 \\ 2 \\ 0 \end{bmatrix}, \quad w = \begin{bmatrix} -2 \\ 2 \\ b \end{bmatrix}$$

- a) Find all values of b such that $w \in \text{Span}(v_1, v_2, v_3)$.
 b) Is the set $\{v_1, v_2, v_3\}$ linearly independent? Justify your answer.

$$\left[\begin{array}{ccc|c} 1 & -1 & 1 & -2 \\ 0 & 1 & 2 & 2 \\ 2 & -3 & 0 & b \end{array} \right] \xrightarrow{R_3 - R_2} \left[\begin{array}{ccc|c} 1 & -1 & 1 & -2 \\ 0 & 1 & 2 & 2 \\ 0 & 3 & 0 & 2b \end{array} \right]$$

Row multiplication is not an elementary row operation!

$$\left[\begin{array}{ccc|c} 1 & -1 & 1 & -2 \\ 0 & 1 & 2 & 2 \\ 0 & 0 & -6 & 2b-6 \end{array} \right] \xrightarrow{\cdot -\frac{1}{6}} \left[\begin{array}{ccc|c} 1 & -1 & 1 & -2 \\ 0 & 1 & 2 & 2 \\ 0 & 0 & 1 & -\frac{1}{3}b+1 \end{array} \right]$$

$$\left[\begin{array}{ccc|c} 1 & 0 & 3 & 0 \\ 0 & 1 & 2 & 2 \\ 0 & 0 & 1 & -\frac{1}{3}b+1 \end{array} \right] \xrightarrow{-3 \cdot R_3} \left[\begin{array}{ccc|c} 1 & 0 & 0 & -b+3 \\ 0 & 1 & 2 & 2 \\ 0 & 0 & 1 & -\frac{1}{3}b+1 \end{array} \right] \xrightarrow{R_2 - 2 \cdot R_3} \left[\begin{array}{ccc|c} 1 & 0 & 0 & -b+3 \\ 0 & 1 & 0 & \frac{2}{3}b \\ 0 & 0 & 1 & -\frac{1}{3}b+1 \end{array} \right]$$

$$\left[\begin{array}{ccc|c} 1 & -1 & 1 & 0 \\ 0 & 1 & 2 & 0 \\ 2 & -3 & 0 & 0 \end{array} \right] \xrightarrow{R_3 + 3 \cdot R_2} \left[\begin{array}{ccc|c} 1 & -1 & 1 & 0 \\ 0 & 1 & 2 & 0 \\ 0 & 0 & 6 & 0 \end{array} \right]$$

$$\left[\begin{array}{ccc|c} 1 & -1 & 1 & 0 \\ 0 & 1 & 2 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

$$x_1 = x_2 - x_3$$

$$x_2 = -3x_3$$

It is not linearly independent because there are infinite solutions.

$$x_3 \begin{bmatrix} -2 \\ -3 \\ 1 \end{bmatrix}$$



2. (10 points) Consider the following matrix:

$$A = \begin{bmatrix} 1 & -1 & 2 \\ 1 & 0 & 1 \\ 0 & 2 & -1 \end{bmatrix}$$

Compute A^{-1} .

$$\left[\begin{array}{ccc|ccc} 1 & -1 & 2 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 2 & -1 & 0 & 0 & 1 \end{array} \right] \rightarrow \left[\begin{array}{ccc|ccc} 1 & -1 & 2 & 1 & 0 & 0 \\ 0 & 2 & -1 & 0 & 0 & 1 \\ 1 & 0 & 1 & 0 & 1 & 0 \end{array} \right] R2 \cdot \frac{1}{2}$$

$$\left[\begin{array}{ccc|ccc} 1 & -1 & 2 & 1 & 0 & 0 \\ 0 & 1 & -\frac{1}{2} & 0 & 0 & \frac{1}{2} \\ 1 & 0 & 1 & 0 & 1 & 0 \end{array} \right] R3 - R2 \rightarrow \left[\begin{array}{ccc|ccc} 1 & -1 & 2 & 1 & 0 & 0 \\ 0 & 1 & -\frac{1}{2} & 0 & 0 & \frac{1}{2} \\ 0 & 0 & 1 & 0 & 1 & 0 \end{array} \right] R1 + \frac{1}{2}R2$$

$$\left[\begin{array}{ccc|ccc} 1 & -1 & 2 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & \frac{1}{2} \\ 0 & 0 & 1 & 0 & 1 & 0 \end{array} \right] R1 + R2 \rightarrow \left[\begin{array}{ccc|ccc} 1 & 0 & 2 & 1 & 0 & \frac{1}{2} \\ 0 & 1 & 0 & 0 & 0 & \frac{1}{2} \\ 0 & 0 & 1 & 0 & 1 & 0 \end{array} \right] R1 - 2R3$$

$$\left[\begin{array}{ccc|ccc} 1 & 0 & 0 & 1 & 0 & \frac{1}{2} \\ 0 & 1 & 0 & 0 & 0 & \frac{1}{2} \\ 0 & 0 & 1 & 0 & 1 & 0 \end{array} \right] A^{-1} = \begin{bmatrix} 1 & 0 & \frac{1}{2} \\ 0 & 0 & \frac{1}{2} \\ 0 & 0 & 0 \end{bmatrix}$$



3. (10 points) Let A be the same matrix as in Problem 2, and let

$$B = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 4 \\ 3 & 2 & 1 \end{bmatrix}$$

Find a matrix C such that $A^T C = B$ (where A^T is the transpose of A).

$$A^T = \begin{bmatrix} 1 & 1 & 0 \\ -1 & 0 & 2 \\ 2 & 1 & -1 \end{bmatrix} \cdot C = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 4 \\ 3 & 2 & 1 \end{bmatrix} \quad \checkmark$$

$$C = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 4 \\ 3 & 2 & 1 \end{bmatrix} \cdot \begin{bmatrix} 1 & 1 & 0 \\ -1 & 0 & 2 \\ 2 & 1 & -1 \end{bmatrix}^{-1}$$

Simpler: $(A^T)^{-1} = (A^{-1})^T$,
then use problem 2.

$$\left[\begin{array}{ccc|ccc} 1 & 1 & 0 & 1 & 0 & 0 \\ -1 & 0 & 2 & 0 & 1 & 0 \\ 2 & 1 & -1 & 0 & 0 & 1 \end{array} \right] \xrightarrow{R_2+R_1} \left[\begin{array}{ccc|ccc} 1 & 1 & 0 & 1 & 0 & 0 \\ 0 & 1 & 2 & 0 & 1 & 0 \\ -1 & 0 & 2 & 0 & 0 & 1 \end{array} \right] \xrightarrow{R_3+R_2} \left[\begin{array}{ccc|ccc} 1 & 1 & 0 & 1 & 0 & 0 \\ 0 & 1 & 2 & 0 & 1 & 0 \\ 0 & 1 & 4 & 0 & 1 & 0 \end{array} \right] \xrightarrow{R_3-R_2} \left[\begin{array}{ccc|ccc} 1 & 1 & 0 & 1 & 0 & 0 \\ 0 & 1 & 2 & 0 & 1 & 0 \\ 0 & 0 & 2 & 0 & 0 & 0 \end{array} \right]$$

$$\left[\begin{array}{ccc|ccc} 1 & 1 & 0 & 1 & 0 & 0 \\ 0 & 1 & 2 & 0 & 1 & 0 \\ 0 & 0 & 2 & 0 & 0 & 0 \end{array} \right] \xrightarrow{R_3-R_2} \left[\begin{array}{ccc|ccc} 1 & 1 & 0 & 1 & 0 & 0 \\ 0 & 1 & 2 & 0 & 1 & 0 \\ 0 & 0 & -2 & 0 & -1 & 0 \end{array} \right] \xrightarrow{R_3 \cdot -\frac{1}{2}} \left[\begin{array}{ccc|ccc} 1 & 1 & 0 & 1 & 0 & 0 \\ 0 & 1 & 2 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & \frac{1}{2} & 0 \end{array} \right] \xrightarrow{-4 \cdot R_3} \left[\begin{array}{ccc|ccc} 1 & 1 & 0 & 1 & 0 & 0 \\ 0 & 1 & 2 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & \frac{1}{2} & 0 \end{array} \right]$$

$$\left[\begin{array}{ccc|ccc} 1 & 1 & 0 & 1 & 0 & 0 \\ 0 & 1 & 2 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & \frac{1}{2} & 0 \end{array} \right] \xrightarrow{-R_2} \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & 1 & -1 & 0 \\ 0 & 1 & 2 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & \frac{1}{2} & 0 \end{array} \right]$$

$$\frac{1}{2} \begin{bmatrix} 3 \\ 1 \\ 1 \end{bmatrix} - \frac{1}{2} \begin{bmatrix} 3 \\ 4 \\ 1 \end{bmatrix} = \begin{bmatrix} -1 \\ -4 \\ 0 \end{bmatrix}$$

$$C = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 4 \\ 3 & 2 & 1 \end{bmatrix} \cdot \begin{bmatrix} 3 & -4 & 0 \\ -2 & 4 & 0 \\ \frac{1}{2} & -\frac{1}{2} & 0 \end{bmatrix}$$

$$3 \begin{bmatrix} \frac{1}{4} \\ \frac{1}{3} \\ \frac{1}{2} \end{bmatrix} + -4 \begin{bmatrix} \frac{1}{4} \\ \frac{1}{3} \\ \frac{1}{2} \end{bmatrix} + 0 \begin{bmatrix} \frac{1}{4} \\ \frac{1}{3} \\ \frac{1}{2} \end{bmatrix} = \begin{bmatrix} -1 \\ -4 \\ 0 \end{bmatrix}$$

$$-2 \begin{bmatrix} \frac{2}{5} \\ \frac{2}{5} \\ \frac{2}{5} \end{bmatrix} + 4 \begin{bmatrix} \frac{2}{5} \\ \frac{2}{5} \\ \frac{2}{5} \end{bmatrix} + 0 = \begin{bmatrix} 4 \\ 16 \\ 4 \end{bmatrix}$$



4. (20 points) Let $T: \mathbb{R}^2 \rightarrow \mathbb{R}^3$ be a linear transformation given by

$$T\left(\begin{bmatrix} x_1 \\ x_2 \end{bmatrix}\right) = \begin{bmatrix} x_1 - 2x_2 \\ x_1 + x_2 \\ x_1 - 3x_2 \end{bmatrix}$$

a) Find the standard matrix of T .

b) Find all vectors u satisfying $T(u) = \begin{bmatrix} 1 \\ 10 \\ -2 \end{bmatrix}$.

$$e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \quad T(e_1) = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} \quad T(e_2) = \begin{bmatrix} -2 \\ 1 \\ -3 \end{bmatrix}$$

$$e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

$$\text{Standard matrix} = \begin{bmatrix} 1 & -2 \\ 1 & 1 \\ 0 & -3 \end{bmatrix} \quad \checkmark$$

$$\left[\begin{array}{cc|c} 1 & -2 & 1 \\ 1 & 1 & 10 \\ 0 & -3 & -2 \end{array} \right] \xrightarrow{R_2 - \frac{1}{3}R_3} \left[\begin{array}{cc|c} 1 & -2 & 1 \\ 0 & 1 & \frac{20}{3} \\ 0 & -3 & -2 \end{array} \right] \xrightarrow{R_3 + 3R_2}$$

$$\left[\begin{array}{cc|c} 1 & -2 & 1 \\ 0 & 1 & \frac{20}{3} \\ 0 & 0 & 18 \end{array} \right] \xrightarrow{R_1 + 2R_2} \left[\begin{array}{cc|c} 1 & 0 & \frac{10}{3} \\ 0 & 1 & \frac{20}{3} \\ 0 & 0 & 18 \end{array} \right]$$

This would mean: no solutions

$$u = \begin{bmatrix} 7 \\ 3 \end{bmatrix}$$



5. (20 points) For each matrix A given below determine if the matrix transformation $T_A: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ given by $T_A(v) = Av$ is one-to-one or not. If T_A is not one-to-one, find two vectors v_1 and v_2 such that $T_A(v_1) = T_A(v_2)$.

a) $A = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 3 & 4 & 4 \end{bmatrix}$

b) $A = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 3 & 4 & 2 \end{bmatrix}$

$$\left[\begin{array}{ccc|c} 1 & 1 & 0 & 0 \\ 0 & 2 & 4 & 0 \\ 3 & 4 & 4 & 0 \end{array} \right] \xrightarrow{\cdot \frac{1}{2}} \left[\begin{array}{ccc|c} 1 & 1 & 0 & 0 \\ 0 & 1 & 2 & 0 \\ 3 & 4 & 4 & 0 \end{array} \right] \xrightarrow{R_3 - R_2} \left[\begin{array}{ccc|c} 1 & 1 & 0 & 0 \\ 0 & 1 & 2 & 0 \\ 0 & 4 & 6 & 0 \end{array} \right] \xrightarrow{R_3 - 4R_2}$$

$$\left[\begin{array}{ccc|c} 1 & 1 & 0 & 0 \\ 0 & 1 & 2 & 0 \\ 0 & 0 & -2 & 0 \end{array} \right]$$

a) ~~not one to one~~

← if not one-to-one, then what are v_1 and v_2 ?

$$\left[\begin{array}{ccc|c} 1 & 1 & 0 & 0 \\ 0 & 2 & 4 & 0 \\ 3 & 4 & 2 & 0 \end{array} \right] \xrightarrow{\cdot \frac{1}{2}} \left[\begin{array}{ccc|c} 1 & 1 & 0 & 0 \\ 0 & 1 & 2 & 0 \\ 3 & 4 & 2 & 0 \end{array} \right]$$

$$\left[\begin{array}{ccc|c} 1 & 1 & 0 & 0 \\ 0 & 1 & 2 & 0 \\ 3 & 4 & 2 & 0 \end{array} \right] \xrightarrow{R_3 - 4R_2}$$

$$\left[\begin{array}{ccc|c} 1 & 1 & 0 & 0 \\ 0 & 1 & 2 & 0 \\ 3 & 0 & -2 & 0 \end{array} \right] \xrightarrow{R_3 - R_2}$$

$$\left[\begin{array}{ccc|c} 1 & 1 & 0 & 0 \\ 0 & 1 & 2 & 0 \\ 0 & 0 & -4 & 0 \end{array} \right] \xrightarrow{\cdot -\frac{1}{4}}$$

$$\left[\begin{array}{ccc|c} 1 & 1 & 0 & 0 \\ 0 & 1 & 2 & 0 \\ 0 & 0 & 1 & 0 \end{array} \right]$$

b) ~~one to one~~ ✓



6. (10 points) For each of the statements given below decide if it is true or false. If it is true explain why. If it is false give a counterexample.

a) If u, v, w are vectors in \mathbb{R}^3 such that $w + u \in \text{Span}(u, v)$ then $w \in \text{Span}(u, v)$.

True, $v_0 + u$ is in the span.
~~if $w + u \in \text{Span}(u, v)$ then $w \in \text{Span}(u, v)$~~
 ?

b) If u, v, w are vectors in \mathbb{R}^3 such that the set $\{u, v, w\}$ is linearly independent then the set $\{u, v\}$ must be linearly independent.

~~false~~, $\{u, v\}$ could have a different solution with nothing trivial



7. (10 points) For each of the statements given below decide if it is true or false. If it is true explain why. If it is false give a counterexample.

a) If A is a 2×2 matrix and u, v are vectors in \mathbb{R}^2 such that Au, Av are linearly dependent then u, v also must be linearly dependent.

~~False~~ false[✓], u, v can be linearly independent before being multiplied by A .
why?

b) If $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is a linear transformation and $u, v, w \in \mathbb{R}^2$ are vectors such that u is in $\text{Span}(v, w)$ then $T(u)$ must be in $\text{Span}(T(v), T(w))$.

True[✓], $T(u)$ is in $\text{span}(T(v), T(w))$ if u is in $\text{span}(v, w)$.
why?