



MTH 309T LINEAR ALGEBRA

EXAM 1

October 3, 2019

Name:

Dorian McMath

UB Person Number:

5	0	2	3	0	4	7	3
0	0	0	0	0	0	0	0
1	1	1	1	1	1	1	1
2	2	2	2	2	2	2	2
3	3	3	3	3	3	3	3
4	4	4	4	4	4	4	4
5	5	5	5	5	5	5	5
6	6	6	6	6	6	6	6
7	7	7	7	7	7	7	7
8	8	8	8	8	8	8	8
9	9	9	9	9	9	9	9

Instructions:

- Textbooks, calculators and any other electronic devices are not permitted. You may use one sheet of notes.
- For full credit solve each problem fully, showing all relevant work.

1	2	3	4	5	6	7	TOTAL	GRADE

--

--

--

--

--

--

--

0

nan

1

2

3

4

5

6

7

TOTAL

GRADE



1. (20 points) Consider the following vectors in \mathbb{R}^3 :

$$v_1 = \begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix}, \quad v_2 = \begin{bmatrix} -1 \\ 1 \\ -3 \end{bmatrix}, \quad v_3 = \begin{bmatrix} 1 \\ 2 \\ 0 \end{bmatrix}, \quad w = \begin{bmatrix} -2 \\ 2 \\ b \end{bmatrix}$$

a) Find all values of b such that $w \in \text{Span}(v_1, v_2, v_3)$.

b) Is the set $\{v_1, v_2, v_3\}$ linearly independent? Justify your answer.

$$\left[\begin{array}{ccc|c} 1 & -1 & 1 & -2 \\ 0 & 1 & 2 & 2 \\ 2 & -3 & 0 & b \end{array} \right] \rightarrow \left[\begin{array}{ccc|c} 1 & -1 & 1 & -2 \\ 0 & 1 & 2 & 2 \\ 0 & -1 & -2 & 4+b \end{array} \right] \rightarrow \left[\begin{array}{ccc|c} 1 & -1 & 1 & -2 \\ 0 & 1 & 2 & 2 \\ 0 & 0 & 0 & 5+b \end{array} \right]$$

$$\rightarrow \left[\begin{array}{ccc|c} 1 & 0 & 3 & 0 \\ 0 & 1 & 2 & 2 \\ 0 & 0 & 0 & 5+b \end{array} \right]$$

free
var

~~must equal~~

a) $b = -5$

b) The set is linearly dependent because
there is not a pivot column in every column



2. (10 points) Consider the following matrix:

$$A = \begin{bmatrix} 1 & -1 & 2 \\ 1 & 0 & 1 \\ 0 & 2 & -1 \end{bmatrix}$$

Compute A^{-1} .

$$\left[\begin{array}{ccc|ccc} 1 & -1 & 2 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 2 & -1 & 0 & 0 & 1 \end{array} \right] \rightarrow \left[\begin{array}{ccc|ccc} 2 & -2 & 4 & 2 & 0 & 0 \\ 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 2 & -1 & 0 & 0 & 1 \end{array} \right] \rightarrow \left[\begin{array}{ccc|ccc} 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & -2 & 2 & 2 & -2 & 0 \\ 0 & 2 & -1 & 0 & 0 & 1 \end{array} \right]$$

$$\left[\begin{array}{ccc|ccc} 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & -2 & 2 & 2 & -2 & 0 \\ 0 & 0 & 1 & 2 & -2 & 1 \end{array} \right] \rightarrow \left[\begin{array}{ccc|ccc} 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & -1 & -1 & 1 & 1 & 0 \\ 0 & 0 & 1 & 2 & -2 & 1 \end{array} \right] \rightarrow \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & -2 & 3 & -1 \\ 0 & 1 & 0 & 1 & -1 & 1 \\ 0 & 0 & 1 & 2 & -2 & 1 \end{array} \right]$$

$$A^{-1} = \begin{bmatrix} -2 & 3 & -1 \\ 1 & -1 & 1 \\ 2 & -2 & 1 \end{bmatrix}$$



3. (10 points) Let A be the same matrix as in Problem 2, and let

$$B = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 4 \\ 3 & 2 & 1 \end{bmatrix}$$

Find a matrix C such that $A^T C = B$ (where A^T is the transpose of A).

$$A^T = \begin{bmatrix} 1 & 1 & 0 \\ -1 & 0 & 2 \\ 2 & 1 & -1 \end{bmatrix}$$

$$\left[\begin{array}{ccc|ccc} 1 & 1 & 0 & 1 & 2 & 3 \\ -1 & 0 & 2 & 4 & 5 & 4 \\ 2 & 1 & -1 & 3 & 2 & 1 \end{array} \right] \rightarrow \left[\begin{array}{ccc|ccc} 1 & 0 & 1 & 2 & 3 & 0 \\ -1 & 0 & 2 & 4 & 5 & 4 \\ 0 & 1 & -3 & 5 & 7 & 7 \end{array} \right]$$

$$\left[\begin{array}{ccc|ccc} -1 & 0 & 2 & 4 & 5 & 4 \\ 0 & 1 & 2 & 5 & 7 & 7 \\ 0 & 1 & -3 & 5 & 7 & 7 \end{array} \right] \rightarrow \left[\begin{array}{ccc|ccc} -1 & 0 & 2 & 4 & 5 & 4 \\ 0 & 1 & 2 & 5 & 7 & 7 \\ 0 & 0 & -5 & 1 & 0 & 0 \end{array} \right] \rightarrow \left[\begin{array}{ccc|ccc} 1 & 0 & -2 & -4 & -5 & -4 \\ 0 & 1 & 2 & 5 & 7 & 7 \\ 0 & 0 & -5 & 1 & 0 & 0 \end{array} \right]$$

$$\left[\begin{array}{ccc|ccc} 1 & 0 & 0 & -4 & -5\frac{1}{3} & -4 \\ 0 & 1 & 0 & 5 & 7\frac{1}{3} & 7 \\ 0 & 0 & 1 & 0 & -\frac{1}{3} & 0 \end{array} \right] \quad \begin{aligned} -\frac{15}{3} - \frac{1}{3} &= -\frac{16}{3} \\ \frac{21}{3} + \frac{1}{3} &= \frac{22}{3} \end{aligned}$$

$$C = \begin{bmatrix} -4 & -\frac{16}{3} & -4 \\ 5 & \frac{22}{3} & 7 \\ 0 & -\frac{1}{3} & 0 \end{bmatrix}$$



4. (20 points) Let $T: \mathbb{R}^2 \rightarrow \mathbb{R}^3$ be a linear transformation given by

$$T\left(\begin{bmatrix} x_1 \\ x_2 \end{bmatrix}\right) = \begin{bmatrix} x_1 - 2x_2 \\ x_1 + x_2 \\ x_1 - 3x_2 \end{bmatrix} \quad 3 \times 2 \quad 2 \times 2$$

a) Find the standard matrix of T .

b) Find all vectors u satisfying $T(u) = \begin{bmatrix} 1 \\ 10 \\ -2 \end{bmatrix}$.

$$\begin{bmatrix} 1 & -2 & 0 \\ 1 & 2 & 0 \\ 1 & -3 & 0 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & -2 & 0 \\ 0 & 4 & 0 \\ 0 & -1 & 0 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & -2 & 0 \\ 0 & -1 & 0 \\ 0 & 4 & 0 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & -2 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

~~$$\begin{bmatrix} 1 & -2 & 0 \\ 1 & 2 & 0 \\ 1 & -3 & 0 \end{bmatrix}$$~~

~~$$\begin{bmatrix} 1 & -2 & 0 \\ 1 & 2 & 0 \\ 1 & -3 & 0 \end{bmatrix}$$~~

a) $\begin{bmatrix} 1 & -2 \\ 1 & 1 \\ 1 & -3 \end{bmatrix}$

b) $u = \begin{bmatrix} 3 \\ 2 \\ -3 \end{bmatrix}$

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} \cdot \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} =$$



5. (20 points) For each matrix A given below determine if the matrix transformation $T_A: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ given by $T_A(v) = Av$ is one-to-one or not. If T_A is not one-to-one, find two vectors v_1 and v_2 such that $T_A(v_1) = T_A(v_2)$.

a) $A = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 3 & 4 & 4 \end{bmatrix}$

$$\begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 0 & 1 & 4 \end{bmatrix}$$

$$\downarrow \begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 1 & 4 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{matrix} \text{leading ones} \\ \text{pivot columns} \end{matrix}$$

* a) one to one

b) $A = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 3 & 4 & 2 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 0 & 1 & 2 \end{bmatrix}$

~~PO~~

$$\downarrow \begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 2 \\ 0 & 1 & 2 \end{bmatrix}$$

→ not one to one *


$$\begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{bmatrix} \begin{matrix} 1 \\ 1 \\ 0 \end{matrix}$$

b) $v_1 = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}$
 $v_2 = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}$



6. (10 points) For each of the statements given below decide if it is true or false. If it is true explain why. If it is false give a counterexample.

a) If u, v, w are vectors in \mathbb{R}^3 such that $w + u \in \text{Span}(u, v)$ then $w \in \text{Span}(u, v)$.



w could be projected outside the span when not added to u

b) If u, v, w are vectors in \mathbb{R}^3 such that the set $\{u, v, w\}$ is linearly independent then the set $\{u, v\}$ must be linearly independent.

True because ~~the~~ ~~the~~ you will still have ~~every~~ every column as a pivot column



7. (10 points) For each of the statements given below decide if it is true or false. If it is true explain why. If it is false give a counterexample.

a) If A is a 2×2 matrix and u, v are vectors in \mathbb{R}^2 such that Au, Av are linearly dependent then u, v also must be linearly dependent.

~~True~~ ~~False~~ ~~True~~ ~~False~~ False

this may simply mean that the matrix A is linearly dependent not necessarily u or v

b) If $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is a linear transformation and $u, v, w \in \mathbb{R}^2$ are vectors such that u is in $\text{Span}(v, w)$ then $T(u)$ must be in $\text{Span}(T(v), T(w))$.

~~True because~~

True because if the same transformation is performed on some u that's in the span of some v and w , ~~it will~~ u will be moved into that new span of $T(v), T(w)$ by the transformation $T(u)$.