



# MTH 309T LINEAR ALGEBRA

## EXAM 1

October 3, 2019

Name:

Emily Blanchard

UB Person Number:

5	0	2	2	8	8	3	7
0	0	0	0	0	0	0	0
1	1	1	1	1	1	1	1
2	2	2	2	2	2	2	2
3	3	3	3	3	3	3	3
4	4	4	4	4	4	4	4
5	5	5	5	5	5	5	5
6	6	6	6	6	6	6	6
7	7	7	7	7	7	7	7
8	8	8	8	8	8	8	8
9	9	9	9	9	9	9	9

Instructions:

- Textbooks, calculators and any other electronic devices are not permitted. You may use one sheet of notes.
- For full credit solve each problem fully, showing all relevant work.

1 2 3 4 5 6 7 TOTAL GRADE

--	--	--	--	--	--	--	--	--

10

--

--

--

--

--

--

10

nan

1

2

3

4

5

6

7

TOTAL

GRADE



1. (20 points) Consider the following vectors in  $\mathbb{R}^3$ :

$$v_1 = \begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix}, \quad v_2 = \begin{bmatrix} -1 \\ 1 \\ -3 \end{bmatrix}, \quad v_3 = \begin{bmatrix} 1 \\ 2 \\ 0 \end{bmatrix}, \quad w = \begin{bmatrix} -2 \\ 2 \\ b \end{bmatrix}$$

a) Find all values of  $b$  such that  $w \in \text{Span}(v_1, v_2, v_3)$ .

b) Is the set  $\{v_1, v_2, v_3\}$  linearly independent? Justify your answer.

$$\left[ \begin{array}{ccc|c} 1 & -1 & 1 & -2 \\ 0 & 1 & 2 & 2 \\ 2 & -3 & 0 & b \end{array} \right] \xrightarrow{R_3 - 2R_2} \left[ \begin{array}{ccc|c} 1 & -1 & 1 & -2 \\ 0 & 1 & 2 & 2 \\ 0 & -1 & -2 & 4+b \end{array} \right] \xrightarrow{R_3 + R_2} \left[ \begin{array}{ccc|c} 1 & 0 & 3 & 0 \\ 0 & 1 & 2 & 2 \\ 0 & -1 & -2 & 4+b \end{array} \right] \xrightarrow{R_3 + R_2} \left[ \begin{array}{ccc|c} 1 & 0 & 3 & 0 \\ 0 & 1 & 2 & 2 \\ 0 & 0 & 0 & 6+b \end{array} \right] \Rightarrow b = 6$$

b) The set  $\{v_1, v_2, v_3\}$  is linearly dependent because

$x_3$  is a free variable meaning there are infinitely many solutions.



2. (10 points) Consider the following matrix:

$$A = \begin{bmatrix} 1 & -1 & 2 \\ 1 & 0 & 1 \\ 0 & 2 & -1 \end{bmatrix}$$

Compute  $A^{-1}$ .

$$\cdot (-1) \left( \begin{array}{ccc|ccc} 1 & -1 & 2 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 2 & -1 & 0 & 0 & 1 \end{array} \right)$$

↓

$$\cdot (-1) \left( \begin{array}{ccc|ccc} 1 & -1 & 2 & 1 & 0 & 0 \\ 0 & 1 & -1 & -1 & 1 & 0 \\ 0 & 2 & -1 & 0 & 0 & 1 \end{array} \right)$$

↓

$$\cdot (-1) \left( \begin{array}{ccc|ccc} 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & -1 & -1 & 1 & 0 \\ 0 & 2 & -1 & 0 & 0 & 1 \end{array} \right)$$

↓

$$\cdot (-1) \left( \begin{array}{ccc|ccc} 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & -1 & -1 & 1 & 0 \\ 0 & 0 & 1 & 2 & -2 & 1 \end{array} \right)$$

↓

$$\cdot (-1) \left( \begin{array}{ccc|ccc} 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 & -1 & 1 \\ 0 & 0 & 1 & 2 & -2 & 1 \end{array} \right)$$

↓

$$\begin{bmatrix} 1 & 0 & 0 & -2 & 3 & -1 \\ 0 & 1 & 0 & 1 & -1 & 1 \\ 0 & 0 & 1 & 2 & -2 & 1 \end{bmatrix}$$

⇒

$$A^{-1} = \begin{bmatrix} -2 & 3 & -1 \\ 1 & -1 & 1 \\ 2 & -2 & 1 \end{bmatrix}$$

check:  $\begin{bmatrix} 1 & -1 & 2 \\ 1 & 0 & 1 \\ 0 & 2 & -1 \end{bmatrix} \cdot \begin{bmatrix} -2 & 3 & -1 \\ 1 & -1 & 1 \\ 2 & -2 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \quad \checkmark$



3. (10 points) Let  $A$  be the same matrix as in Problem 2, and let

$$B = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 4 \\ 3 & 2 & 1 \end{bmatrix} \quad A = \begin{bmatrix} 1 & -1 & 2 \\ 1 & 0 & 1 \\ 0 & 2 & -1 \end{bmatrix}$$

$$Ax = b \\ x = A^{-1} \cdot b$$

Find a matrix  $C$  such that  $A^T C = B$  (where  $A^T$  is the transpose of  $A$ ).

$$A^T C = B \Rightarrow C = (A^T)^{-1} \cdot B \quad (A^T)^{-1} = (A^{-1})^T$$

$$A^{-1} = \begin{bmatrix} -2 & 3 & -1 \\ 1 & -1 & 1 \\ 2 & -2 & 1 \end{bmatrix} \quad (A^{-1})^T = \begin{bmatrix} -2 & 1 & 2 \\ 3 & -1 & -2 \\ -1 & 1 & 1 \end{bmatrix}$$

$$(A^{-1})^T \cdot B = \begin{bmatrix} -2 & 1 & 2 \\ 3 & -1 & -2 \\ -1 & 1 & 1 \end{bmatrix} \cdot \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 4 \\ 3 & 2 & 1 \end{bmatrix} = \begin{bmatrix} -2+4+6 & -4+5+4 & -6+4+2 \\ 3-4-6 & 6-5-4 & 9-4-2 \\ -1+4+3 & -2+5+2 & -3+4+1 \end{bmatrix}$$

$$C = \begin{bmatrix} 8 & 5 & 0 \\ -7 & -3 & 3 \\ 6 & 5 & 2 \end{bmatrix}$$



4. (20 points) Let  $T: \mathbb{R}^2 \rightarrow \mathbb{R}^3$  be a linear transformation given by

$$T\left(\begin{bmatrix} x_1 \\ x_2 \end{bmatrix}\right) = \begin{bmatrix} x_1 - 2x_2 \\ x_1 + x_2 \\ x_1 - 3x_2 \end{bmatrix}$$

a) Find the standard matrix of  $T$ .

b) Find all vectors  $u$  satisfying  $T(u) = \begin{bmatrix} 1 \\ 10 \\ -2 \end{bmatrix}$ .

a)

$$A = [T(e_1) \quad T(e_2)] \quad T(e_1) = T\left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}\right) = \begin{bmatrix} 1 - 2(0) \\ 1 + 0 \\ 1 - 3(0) \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$$

$$T(e_2) = T\left(\begin{bmatrix} 0 \\ 1 \end{bmatrix}\right) = \begin{bmatrix} 0 - 2(1) \\ 0 + 1 \\ 0 - 3(1) \end{bmatrix} = \begin{bmatrix} -2 \\ 1 \\ -3 \end{bmatrix}$$

$$\text{standard matrix of } T = A = \begin{bmatrix} 1 & -2 \\ 1 & 1 \\ 1 & -3 \end{bmatrix}$$

b)  $A \cdot u = T(u)$

$$\begin{bmatrix} 1 & -2 \\ 1 & 1 \\ 1 & -3 \end{bmatrix} \cdot \begin{bmatrix} u_1 \\ u_2 \end{bmatrix} = \begin{bmatrix} 1 \\ 10 \\ -2 \end{bmatrix}$$

$$\left[ \begin{array}{cc|c} 1 & -2 & 1 \\ 1 & 1 & 10 \\ 1 & -3 & -2 \end{array} \right] \xrightarrow{R_2 - R_1, R_3 - R_1} \left[ \begin{array}{cc|c} 1 & -2 & 1 \\ 0 & 3 & 9 \\ 0 & -1 & -3 \end{array} \right] \xrightarrow{R_2 \leftrightarrow R_3} \left[ \begin{array}{cc|c} 1 & -2 & 1 \\ 0 & -1 & -3 \\ 0 & 3 & 9 \end{array} \right] \xrightarrow{R_2 \cdot (-1)} \left[ \begin{array}{cc|c} 1 & -2 & 1 \\ 0 & 1 & 3 \\ 0 & 3 & 9 \end{array} \right] \xrightarrow{R_3 - 3R_2} \left[ \begin{array}{cc|c} 1 & -2 & 1 \\ 0 & 1 & 3 \\ 0 & 0 & 0 \end{array} \right]$$

$$\left[ \begin{array}{cc|c} 1 & 0 & 7 \\ 0 & 1 & 3 \\ 0 & 0 & 0 \end{array} \right] \xrightarrow{R_1 + 2R_2} \left[ \begin{array}{cc|c} 1 & 0 & 7 \\ 0 & 1 & 3 \\ 0 & 0 & 0 \end{array} \right]$$

$$x_1 = 7$$

$$x_2 = 3$$

$$u = \begin{bmatrix} 7 \\ 3 \end{bmatrix}$$

$$\text{check: } T\left(\begin{bmatrix} 7 \\ 3 \end{bmatrix}\right) = \begin{bmatrix} 7 - 2(3) \\ 7 + 3 \\ 7 - 3(3) \end{bmatrix} = \begin{bmatrix} 1 \\ 10 \\ -2 \end{bmatrix} \quad \checkmark$$



5. (20 points) For each matrix  $A$  given below determine if the matrix transformation  $T_A: \mathbb{R}^3 \rightarrow \mathbb{R}^3$  given by  $T_A(v) = Av$  is one-to-one or not. If  $T_A$  is not one-to-one, find two vectors  $v_1$  and  $v_2$  such that  $T_A(v_1) = T_A(v_2)$ .

a)  $A = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 3 & 4 & 4 \end{bmatrix}$

b)  $A = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 3 & 4 & 2 \end{bmatrix}$

a)  $\begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 3 & 4 & 4 \end{bmatrix} \xrightarrow{(-3)} \begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 0 & 1 & 4 \end{bmatrix} \xrightarrow{\cdot \frac{1}{2}} \begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 2 \\ 0 & 1 & 4 \end{bmatrix} \xrightarrow{(-1)} \begin{bmatrix} 1 & 0 & -2 \\ 0 & 1 & 2 \\ 0 & 1 & 4 \end{bmatrix} \xrightarrow{(-1)} \begin{bmatrix} 1 & 0 & -2 \\ 0 & 1 & 2 \\ 0 & 0 & 2 \end{bmatrix} \xrightarrow{\cdot \frac{1}{2}} \begin{bmatrix} 1 & 0 & -2 \\ 0 & 1 & 2 \\ 0 & 0 & 1 \end{bmatrix} \xrightarrow{(-2)} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$

Since  $A$  has a pivot position in every column,  $T_A(v)$  is one-to-one.

b)  $\begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 3 & 4 & 2 \end{bmatrix} \xrightarrow{(-3)} \begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 4 \\ 0 & 1 & 2 \end{bmatrix} \xrightarrow{\cdot \frac{1}{2}} \begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 2 \\ 0 & 1 & 2 \end{bmatrix} \xrightarrow{(-1)} \begin{bmatrix} 1 & 0 & -2 \\ 0 & 1 & 2 \\ 0 & 1 & 2 \end{bmatrix} \xrightarrow{(-1)} \begin{bmatrix} 1 & 0 & -2 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{bmatrix}$

Since  $A$  does not have a pivot position in every column,  $T_A(v)$  is not one-to-one.

$T_A(v_1) = T_A(v_2)$

Let  $T_A(v_1) = T_A(v_2) = \begin{bmatrix} 1 \\ 2 \\ 0 \end{bmatrix}$

$\begin{bmatrix} x_1 & x_2 & x_3 \\ 1 & 0 & -2 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{bmatrix} \begin{cases} x_1 = 1 + 2x_3 \\ x_2 = 2 - 2x_3 \\ x_3 = x_3 \end{cases}$

if  $x_3 = 1$ :

$x_1 = 1 + 2(1) = 3$   
 $x_2 = 2 - 2(1) = 0$   
 $x_3 = 1$

if  $x_3 = -1$ :

$x_1 = 1 + 2(-1) = -1$   
 $x_2 = 2 - 2(-1) = 4$   
 $x_3 = -1$

$v_1 = \begin{bmatrix} 1 \\ 3 \\ 0 \end{bmatrix} \quad v_2 = \begin{bmatrix} -1 \\ 4 \\ -1 \end{bmatrix}$



6. (10 points) For each of the statements given below decide if it is true or false. If it is true explain why. If it is false give a counterexample.

a) If  $u, v, w$  are vectors in  $\mathbb{R}^3$  such that  $w + u \in \text{Span}(u, v)$  then  $w \in \text{Span}(u, v)$ .

True. Vector  $u$  must be in the  $\text{Span}(u, v)$ , and given that  $w + u \in \text{Span}(u, v)$  then  $w + u$  must also be in the  $\text{Span}(u, v)$ .

b) If  $u, v, w$  are vectors in  $\mathbb{R}^3$  such that the set  $\{u, v, w\}$  is linearly independent then the set  $\{u, v\}$  must be linearly independent.

True. In order for  $\{u, v, w\}$  to be linearly independent, it must have a leading one in every column. This means  $\{u, v\}$  also has a leading one in every column, so  $\{u, v\}$  is linearly independent.





7. (10 points) For each of the statements given below decide if it is true or false. If it is true explain why. If it is false give a counterexample.

a) If  $A$  is a  $2 \times 2$  matrix and  $u, v$  are vectors in  $\mathbb{R}^2$  such that  $Au, Av$  are linearly dependent then  $u, v$  also must be linearly dependent.

$\{Au, Av\}$  linearly dependent

False. Multiplying matrix  $A$  by vectors  $u$  and  $v$  does not necessarily preserve linear dependence.

b) If  $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$  is a linear transformation and  $u, v, w \in \mathbb{R}^2$  are vectors such that  $u$  is in  $\text{Span}(v, w)$  then  $T(u)$  must be in  $\text{Span}(T(v), T(w))$ .

True. If vector  $u \in \text{span}(v, w)$  and transformation  $T$  is applied to  $u, v, w$  then  $T(u) \in \text{span}(T(v), T(w))$ .